



Synflorix

Pneumococcal polysaccharide and Non-Typeable *Haemophilus influenzae* (NTHi) protein D conjugate vaccine, adsorbed Suspension for injection

QUALITATIVE AND QUANTITATIVE COMPOSITION

One dose (0.5 mL) contains 1 microgram of polysaccharide for serotypes 1^{1,2}, 5^{1,2}, 6B^{1,2}, 7F^{1,2}, 9V^{1,2}, 14^{1,2} and 23F^{1,2}, and 3 micrograms for serotypes 4^{1,2}, 18C^{1,3} and 19F^{1,4}.

¹adsorbed on aluminium phosphate

0.5 milligram Al³⁺

²conjugated to protein D (derived from NTHi) carrier protein

~13 micrograms

³conjugated to tetanus toxoid carrier protein

~8 micrograms

⁴conjugated to diphtheria toxoid carrier protein

~5 micrograms

Synflorix is a turbid white suspension. Upon storage, a fine white deposit with a clear colourless supernatant can be observed.

CLINICAL INFORMATION

Indications

Active immunization against invasive disease (including sepsis, meningitis, bacteraemic pneumonia and bacteremia) and acute otitis media caused by *Streptococcus pneumoniae* serotype 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F in infants and children from 2 months up to 5 years of age. See section *Warnings and Precautions* and *Pharmacodynamics* for information on protection against specific pneumococcal serotypes.

The use of **Synflorix** should be determined on the basis of official recommendations taking into consideration the impact of invasive disease in different age groups as well as the variability of serotype epidemiology in different geographical areas.

Dosage and Administration

Official recommendations should be taken into account when immunising with **Synflorix**.

Infants from 2 months to 6 months of age:

3-dose primary series

The recommended immunization series to ensure optimal protection consists of four doses, each of 0.5 mL. The primary infant series consists of three doses with the first dose usually given at 2 months of age and with an interval of at least 1 month between doses. A booster dose is recommended at least 6 months after the last primary dose (see section *Pharmacodynamics*).

2-dose primary series

Alternatively, when **Synflorix** is given as part of a routine infant immunization programme, a series consisting of 3 doses, each of 0.5 mL may be given. The first dose may be administered from the age of 2 months, with a second dose 2 months later. A booster dose is recommended at least 6 months after the last primary dose (see section *Pharmacodynamics*).

Preterm infants born after at least 27 weeks of gestational age

The recommended immunization series consists of 4 doses, each of 0.5 mL. The primary infant series consists of 3 doses with the first dose usually given at 2 months of age and with an interval of at least 1 month between doses. A booster dose is recommended at least 6 months after the last primary dose (see section *Pharmacodynamics*).

Previously unvaccinated older infants and children:

- 7-11 months of age: 2 doses of 0.5 mL with an interval of at least 1 month between doses. A third dose is recommended in the second year of life with an interval of at least 2 months.
- 12 months - 5 years of age: 2 doses of 0.5 mL with an interval of at least 2 months between doses.

Special populations:

The safety and immunogenicity of **Synflorix** in children with underlying immunocompromising conditions predisposed to invasive pneumococcal diseases (such as SCD, or HIV infection) have been studied in children below 2 years of age. Limited data are available for children with asplenia or splenic dysfunction ages 2 to 5 years.

Synflorix may be given according to the above mentioned schedules, except that a 3-dose schedule should be given as primary vaccination in infants starting vaccination from 2 months to 6 months of age (see *Warnings and Precautions* and *Pharmacodynamics*).

It is recommended that subjects who receive a first dose of **Synflorix** complete the full vaccination course with **Synflorix**.

The vaccine should be given by intramuscular injection. The preferred sites are anterolateral aspect of the thigh in infants or the deltoid muscle of the upper arm in children.

Contraindications

Synflorix should not be administered to subjects with known hypersensitivity to any component of the vaccine (see *Qualitative and Quantitative Composition* and *List of Excipients*).

Warnings and Precautions

It is good clinical practice to precede vaccination by a review of the medical history (especially with regard to previous vaccination and possible occurrence of undesirable events) and a clinical examination.

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of a rare anaphylactic event following the administration of the vaccine.

As with other vaccines, the administration of **Synflorix** should be postponed in subjects suffering from acute severe febrile illness. However, the presence of a minor infection, such as a cold, should not result in the deferral of vaccination.

Synflorix should under no circumstances be administered intravascularly or intradermally. No data are available on subcutaneous administration of **Synflorix**.

Syncpe (fainting) can occur following, or even before, any vaccination as a psychogenic response to the needle injection. It is important that procedures are in place to avoid injury from faints.

As for other vaccines administered intramuscularly, **Synflorix** should be given with caution to individuals with thrombocytopenia or any coagulation disorder since bleeding may occur following an intramuscular administration to these subjects.

There is insufficient evidence that **Synflorix** provides protection against pneumococcal serotypes not contained in the vaccine or against non-typeable *Haemophilus influenzae*. **Synflorix** does not provide protection against other microorganism.

As with any vaccine, **Synflorix** may not protect all vaccinated individuals against invasive pneumococcal disease or otitis media caused by the serotypes in the vaccine. Protection against otitis media caused by pneumococcal serotypes in the vaccine is expected to be substantially lower than protection against invasive disease. In addition, as otitis media is caused by many microorganism other than the *Streptococcus pneumoniae* serotypes represented in the vaccine, the overall protection against otitis media is expected to be limited.

Synflorix will not protect against pneumococcal serogroups other than those included in the vaccine. Although antibody response to diphtheria toxoid, tetanus toxoid and Protein D (protein D is highly conserved in all *Haemophilus influenzae* strains including NTHi) occurs, immunization with **Synflorix** does not substitute routine immunization with diphtheria, tetanus or *Haemophilus influenzae* type b (Hib) vaccines. Official recommendations for the immunizations against diphtheria, tetanus and Hib should also be followed.

As with any vaccine, a protective immune response may not be elicited in all vaccinees.

Safety and immunogenicity data are available for HIV infected infants, children with SCD and children with splenic dysfunction (see *Adverse Reactions* and *Pharmacodynamics*). Safety and immunogenicity data for **Synflorix** are not available for individuals in other specific immunocompromised groups and vaccination should be considered on an individual basis.

Children with impaired immune responsiveness, whether due to the use of immunosuppressive therapy, a genetic defect, HIV infection, or other causes, may have reduced antibody response to active immunization.

For children at high-risk for pneumococcal disease (such as children with SCD, asplenia, HIV infection, chronic illness or those who have other immunocompromising conditions), the appropriate-for-age **Synflorix** vaccination series should be given (see *Dosage and Administration*). The use of pneumococcal conjugate vaccine does not replace the use of 23-valent pneumococcal polysaccharide vaccines which should be given according to local recommendations in those children.

Prophylactic administration of antipyretics before or immediately after vaccines administration can reduce the incidence and intensity of post-vaccination febrile reactions. Data however, suggest that the use of prophylactic paracetamol might reduce the immune response to pneumococcal vaccines. The clinical relevance of this observation remains unknown.

The use of prophylactic antipyretic medicinal products is recommended:

- For all children receiving **Synflorix** simultaneously with vaccines containing whole cell pertussis because of higher rate of febrile reactions (see section *Adverse Reactions*)
- For children with seizure disorders or with a prior history of febrile seizures

Antipyretic treatment should be initiated according to local treatment guidelines.

The potential risk of apnoea and the need for respiratory monitoring for 48-72h should be considered when administering the primary immunization series to very premature infants (born ≤ 28 weeks of gestation) and particularly for those with a previous history of respiratory immaturity. As the benefit of vaccination is high in this group of infants, vaccination should not be withheld or delayed.

Interactions

Synflorix can be given concomitantly with any of the following monovalent or combination vaccines [including DTPa-HBV-IPV/Hib and DTPw-HBV/Hib]: diphtheria-tetanus-acellular pertussis vaccine (DTPa), hepatitis B vaccine (HBV), inactivated polio vaccine (IPV), *Haemophilus influenzae* type b vaccine (Hib), diphtheria-tetanus-whole cell pertussis vaccine (DTPw), measles-mumps-rubella vaccine (MMR), varicella vaccine, meningococcal serogroup C conjugate vaccine (CRM₁₉₇ and TT conjugates), meningococcal serogroup A, C, W-135 and Y conjugate vaccine (MenACWY-TT), oral polio vaccine (OPV) and rotavirus vaccine. Different injectable vaccines should always be given at different injection sites.

Clinical studies demonstrated that the immune responses and the safety profiles of the co-administered vaccines were unaffected, with the exception of the inactivated poliovirus type 2 response (seroprotection ranging from 78% to 100% across studies) and MenACWY-TT vaccine when co-administered with a booster dose of **Synflorix** following a 3 doses primary series (lower antibody geometric mean concentration (GMC) and opsonophagocytic assay geometric mean titre (OPA GMT) for pneumococcal serotype 18C only). Enhancement of antibody response to Hib-TT conjugate, diphtheria and tetanus antigens was observed. The clinical relevance of the above observations is unknown.

As with other vaccines it may be expected that in patients receiving immunosuppressive treatment an adequate response may not be elicited.

Adverse Reactions

Clinical trials involved the administration of approximately 64,000 doses of **Synflorix** to approximately 22,500 healthy children and 137 preterm infants as primary vaccination. Approximately 19,500 healthy children and 116 preterm infants received a booster dose of **Synflorix** in the second year of life. Safety was also assessed in approximately 400 children from 2 to 5 years old. In all trials, **Synflorix** was administered concurrently with the recommended childhood vaccines.

An increase in reactogenicity was reported after booster vaccination compared to the doses of the primary courses with **Synflorix**.

No increase in the incidence or severity of the adverse reactions was seen with subsequent doses of the primary vaccination series.

Reactogenicity was higher in children receiving whole cell pertussis vaccines concomitantly.

Adverse reactions reported (for all age groups) are listed according to the following frequency:

Very common:	≥1/10
Common:	≥1/100 to <1/10
Uncommon:	≥1/1,000 to <1/100
Rare:	≥1/10,000 to <1/1,000
Very rare:	<1/10,000

Frequency	Adverse reactions
Clinical trials	
Very common	Appetite lost, irritability, drowsiness, pain, redness, swelling at the injection site, fever ≥38°C rectally (age <2 years)
Common	Injection site reactions like injection site induration, fever >39°C rectally (age <2 years)
Uncommon	Crying abnormal, apnoea in very premature infants (≤28 weeks of gestation) (see <i>Warnings and Precautions</i>), diarrhoea, vomiting, rash, injection site reactions like injection site haematoma, haemorrhage and nodule
Rare	Allergic reactions (such as allergic dermatitis, atopic dermatitis, eczema), convulsions (including febrile convulsions), urticaria ⁽¹⁾
Very rare	Angioedema, Kawasaki disease
<i>Adverse reactions additionally reported after booster vaccination of primary series and/or catch-up vaccination:</i>	
Common	Fever ≥38°C rectally (age 2 to 5 years)
Uncommon	Injection site reactions ⁽²⁾ like pruritus, diffuse swelling of the injected limb, sometimes involving the adjacent joint; age <2 years: fever >40°C rectally; age 2 to 5 years: headache, nausea and fever >39°C rectally
Post-marketing experience	
Rare	Hypotonic-hyporesponsive episode
Very rare	Anaphylaxis

⁽¹⁾ Uncommon following catch-up vaccination in children 12 to 23 months of age.

⁽²⁾ Following booster vaccination, children >12 months of age are more likely to experience injection site reactions when compared to infants during the primary series.

Safety of **Synflorix** was assessed in 83 HIV positive infants, 101 HIV negative infants born from an HIV positive mother and 150 children with SCD. Results suggest comparable reactogenicity and safety profile of **Synflorix** between these high risk groups and healthy children.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamics

Pharmaco-therapeutic group: pneumococcal vaccines, ATC code: J07AL52.

1. Efficacy and effectiveness in clinical trials

In a large-scale phase III/IV, double-blind, cluster-randomized, controlled, clinical trial in Finland (FinIP), children received either **Synflorix** or control vaccines according to a 3+1 or 2+1 infant schedule (3-4-5 months of age or 3-5 months of age with booster at 11 months). In the catch-up cohorts, children 7-11 months of age or 12-18 months of age at first vaccine dose received either **Synflorix** or control vaccines according to the appropriate-for-age **Synflorix** vaccination schedule.

In a large-scale phase III, randomized, double-blind clinical trial (Clinical Otitis Media and Pneumonia Study - COMPAS), infants aged 6-16 weeks received either **Synflorix** or the control vaccine according to a 3+1 schedule (2-4-6 months of age with booster at 15-18 months).

1.1 Invasive Pneumococcal Disease (IPD)

Infant cohort below 7 months of age at enrolment

Vaccine effectiveness (in FinIP) or efficacy (in COMPAS) was demonstrated in preventing culture-confirmed IPD due to vaccine serotypes (Table 1).

Table 1: Prevention of IPD in infants receiving at least one dose of *Synflorix* (Infant total vaccinated cohort)

Type of IPD	FinIP						COMPAS		
	No. of cases			VE (95% CI)		No. of cases		VE (95% CI)	
	<i>Synflorix</i> 3+1 schedule	<i>Synflorix</i> 2+1 schedule	Control ⁽²⁾	3+1 schedule	2+1 schedule	<i>Synflorix</i> 3+1 schedule	Control	3+1 schedule	
	N 10,273	N 10,054	N 10,200			N 11,798	N 11,799		
Vaccine serotype ⁽¹⁾	0	1	12	100% ⁽³⁾ (82.8; 100)	91.8% ⁽⁴⁾ (58.3; 99.6)	0	18	100% (77.3;100)	
Serotype 6B	0	0	5	100% (54.9; 100)	100% (54.5; 100)	0	2	-	
Serotype 14	0	0	4	100% (39.6; 100)	100% (43.3; 100)	0	9	100% (49.5;100)	
Any serotype	0	2	14	100% ⁽⁵⁾ (85.6;100)	85.8% ⁽⁵⁾ (49.1;97.8)	7	21	66.7% (21.8;85.9)	

IPD: Invasive Pneumococcal Disease

VE: Vaccine effectiveness (FinIP) or efficacy (COMPAS)

N: number of subjects per group; CI: Confidence Interval

⁽¹⁾ In FinIP, the other vaccine serotypes causing IPD were 7F (1 case in the *Synflorix* 2+1 clusters), 18C, 19F and 23F (1 case of each in the control clusters). In COMPAS, serotypes 5 (2 cases), 18C (4 cases) and 23F (1 case) were also detected in control group in addition to serotypes 6B and 14.

⁽²⁾ the 2 groups of control clusters of infants were pooled

⁽³⁾ p-value<0.0001

⁽⁴⁾ p-value=0.0009

⁽⁵⁾ 93.0% (95% CI, 74.9;98.9; 2 versus 14 cases) regardless of the primary schedule

Catch-up cohorts

Among the 15,447 children in the catch-up cohorts, there were no culture-confirmed IPD cases in the *Synflorix* groups while 7 IPD cases were observed in the control groups (2 and 5 cases in the 7-11 months and 12-18 months cohorts respectively).

1.2. Pneumonia

Vaccine efficacy of *Synflorix* against likely bacterial Community Acquired Pneumonia (CAP), i.e. radiologically confirmed CAP cases with either alveolar consolidation/pleural effusion on the chest X-ray, or with non-alveolar infiltrates but with C reactive protein (CRP) ≥ 40 mg/L, was demonstrated in the according-to-protocol (ATP) cohort (immunized with at least the 3-dose primary series) as the primary objective of COMPAS during a follow up of 38 months from study start: 22.0% (95% CI: 7.7; 34.2); P value ≤ 0.002 ; 240 cases/10,295 subjects in the *Synflorix* group versus 304 cases/10,201 subjects in the control group.

Vaccine efficacy against CAP with alveolar consolidation or pleural effusion was 25.7% (95% CI: 8.4; 39.6) and against clinically suspected CAP referred for X-ray was 6.7% (95% CI: 0.7; 12.3).

During an observation period of 48 months from study start, vaccine efficacy against likely bacterial CAP was 18.2% (95% CI: 4.1; 30.3), against CAP with alveolar consolidation or pleural effusion 22.4% (95% CI: 5.7; 36.1) and against clinically suspected CAP referred for X-ray 7.3% (95% CI: 1.6; 12.6).

In the FinIP study, vaccine effectiveness in reducing hospital-diagnosed pneumonia cases was 26.7% (95% CI: 4.9; 43.5) in the 3+1 infant schedule and 29.3% (95% CI: 7.5; 46.3) in the 2+1 infant schedule. Vaccine effectiveness was 33.2% (95% CI: 3.0; 53.4) in the 7-11 month cohort and 22.4% (95% CI: -8.7; 44.8) in the 12-18 month cohort.

1.3. Acute Otitis Media (AOM)

Table 2: Vaccine efficacy against AOM⁽¹⁾ in COMPAS (ATP⁽²⁾: 5,989 subjects)

Type or cause of AOM	Vaccine efficacy	95% CI
Clinical AOM regardless of aetiology	16.1%	-1.1; 30.4 ⁽³⁾
Any pneumococcal serotype	56.1%	13.4; 77.8
10 pneumococcal vaccine serotypes	67.1%	17.0; 86.9
Vaccine-related pneumococcal serotypes	25.7%	-232.2; 83.4
Non-vaccine/non-vaccine related pneumococcal serotypes	25.7%	-231.9; 83.4
NTHi	15.0%	-83.8;60.7

CI Confidence Interval

⁽¹⁾ First episode

⁽²⁾ Follow up period for a maximum of 40 months from 2 weeks after third primary dose

(3) Not statistically significant by pre-defined criteria (One sided p=0.032). However, in TVC cohort, efficacy against clinical AOM episodes was 19% (95% CI: 4.4; 31.4)

Table 3: Vaccine efficacy against AOM⁽²⁾ in POET⁽¹⁾ (ATP⁽³⁾: 4,907 subjects)

Type or cause of AOM	Vaccine efficacy	95% CI
Clinical AOM regardless of aetiology	33.6%	20.8; 44.3
Any pneumococcal serotype	51.5%	36.8; 62.9
10 pneumococcal serotypes in common with Synflorix	67.9%	53.0; 78.1
Vaccine-related pneumococcal serotypes	65.5%	22.4; 84.7
Non-vaccine/non-vaccine related pneumococcal serotypes	8.5%	-64.2; 49.0
NTHi	35.3%	1.8; 57.4

CI Confidence Interval

(1) Large randomised double-blind trial in the Czech Republic and Slovakia where infants received either an 11-valent investigational vaccine containing the 10 serotypes of **Synflorix** (along with serotype 3 for which efficacy was not demonstrated), or a control vaccine, according to a 3-4-5 and 12-15 months schedule.

(2) All episodes

(3) Follow up period for a maximum of 24 months from 2 weeks after third primary dose

No increase in the incidence of AOM due to non-vaccine/non-vaccine related serotypes, or due to other bacterial pathogens was observed in either COMPAS (based on the few cases reported) or POET trial.

Impact on antimicrobial prescriptions

In the FinIP infant total vaccinated cohort, vaccination with **Synflorix** reduced outpatient prescriptions for amoxicillin, the most frequently prescribed antibiotic for AOM, by 7.9% (95% CI: 2.0; 13.4) in the 3+1 schedule and 7.5% (95% CI: 0.9; 13.6) in the 2+1 schedule. In the **Synflorix** groups, there was a trend for a reduction in any outpatient antimicrobial prescriptions and in antimicrobial prescriptions usually recommended for otitis media and respiratory infections.

1.4 Impact on nasopharyngeal carriage (NPC)

The effect of **Synflorix** on NPC was studied in the nested study of FinIP (5,092 subjects) and in COMPAS (1,921 subjects). In both studies, **Synflorix** significantly reduced vaccine type carriage (combined and 6B, 19F and 23F individually) with a trend for increase after booster vaccination in non-vaccine/non-vaccine related type NPC resulting in net decrease in overall pneumococcal carriage. In the nested study, a significant reduction was also observed for vaccine serotype 14 and for serotype 19A.

In a clinical study assessing NPC in HIV positive infants (HIV+/+, N=83) and HIV negative infants born from an HIV positive mother (HIV+/-, N=101), the HIV exposure or infection did not appear to alter the effect of **Synflorix** on pneumococcal carriage when compared to the effect in HIV negative infants born from an HIV negative mother (HIV-/, N=100).

2. Effectiveness in post-marketing surveillance

In Brazil, **Synflorix** was introduced into the national immunization program (NIP) in March 2010, using a 3+1 schedule in infants with a catch-up campaign in children up to 2 years of age. Based on almost 3 years of surveillance following **Synflorix** introduction, a matched case-control study reported a significant decrease in culture or PCR confirmed IPD due to any vaccine serotype (83.8% (95% CI: 65.9;92.3)).

In Finland, **Synflorix** was introduced into NIP in September 2010, with a 2+1 schedule in infants without catch-up campaign. The relative rate reduction of IPD incidence in children ≤5 years of age during the first 3 years after NIP introduction was evaluated. Before and after NIP comparison suggests a significant decrease in the incidence of any culture confirmed IPD (80% (95% CI: 72;85)), any vaccine serotype IPD (92% (95% CI: 86;95)).

In Quebec, Canada, **Synflorix** was introduced into the infant immunization programme (2 primary doses to infants <6 months of age and a booster dose at 12 months) following 4.5 years of use of 7-valent Pneumococcal Conjugate Vaccine (PCV). Based on 1.5 years of surveillance following **Synflorix** introduction, with over 90% coverage in the vaccine-eligible age group, a decrease in vaccine serotype IPD incidence (largely due to changes in serotype 7F disease) was observed with no concomitant increase in non-vaccine serotype IPD incidence, leading to an overall decrease in IPD incidence in the target age group compared to the incidence reported during the preceding period.

3. Immunogenicity data

3.1 Immunologic non-inferiority to 7-valent PCV

In a head-to-head comparative trial with 7-valent PCV, non-inferiority of the immune response to **Synflorix** measured by ELISA was demonstrated for all serotypes, except for 6B and 23F. The clinical relevance of these differences is unclear, as **Synflorix** was observed to be effective against IPD caused by serotype 6B in a clinical study (see Table 1). The percentage of vaccinees reaching the ELISA antibody threshold (i.e. 0.20 µg/mL) for serotypes 1, 5 and 7F in **Synflorix** was at least as good as the aggregate 7-valent PCV response against the 7 common serotypes. The proportion of functional antibody responders (OPA titre ≥ 8) to all serotypes contained in each vaccine were high (>87.7%) with the exception of serotype 1 for **Synflorix** post-primary (65.7%). Immunological memory was shown for all vaccine serotypes after a booster dose in the second year of life.

It has also been demonstrated that **Synflorix** induces an immune response to serotype 19A with 6.1 fold increases in both antibody GMC and OPA GMT observed 1 month after a booster dose compared to pre-booster concentrations.

3.2 Immunogenicity in infants

In clinical trials, the immunogenicity of **Synflorix** was evaluated after a 2- or 3-dose primary course according to different schedules (2-4, 3-5 months of age or 2-3-4, 3-4-5, 2-4-6 months of age) and after a booster dose given at least 6 months after the last primary dose.

A 3-dose primary schedule has shown higher response against protein D compared to a 2-dose primary schedule. However, the clinical relevance of this observation remains unknown.

Immune memory

After a single challenge dose of **Synflorix** in the 4th year of life, similar anamnestic immune response was observed for all vaccine serotypes. Anamnestic immune responses to protein D were shown with both schedules.

3.3 Immunogenicity in unvaccinated infants and children ≥ 7 months of age (catch-up)

In studies in previously unvaccinated 7-11 months children (2+1 schedule) and children 12 months up to 5 years of age (2 dose schedule), antibody GMCs and OPA GMTs for vaccine serotypes were similar or higher than those induced by 3-dose primary infant series. A similar immune response was observed for protein D in 2 to 5 years old children and infants after a 3-dose primary series.

3.4 Immunogenicity in preterm infants

Immunogenicity of **Synflorix** in very preterm and preterm (gestation period of 27-30 weeks and 31-36 weeks respectively) as well as full term infants was evaluated (3 primary doses at 2, 4, 6 months of age with a booster dose at 15-18 months of age).

After primary vaccination, for each vaccine serotype the proportion of subjects with ELISA antibody concentrations ≥ 0.20 µg/mL and OPA titres ≥ 8 was similar regardless of maturity. With respect to full term, similar immunogenicity was observed in preterm groups except lower antibody GMCs for vaccine serotypes 4, 5, 9V and lower OPA GMT for serotype 5. Immunological memory was shown for vaccine serotype one month after the booster dose.

3.5 Immunogenicity in special populations

In a clinical study in South Africa, **Synflorix** was given to HIV+/+ (asymptomatic or mild disease), HIV+/- and HIV-/ infants (3 primary doses at 6-10-14 weeks of age with a booster dose at 9-10 months of age). Group comparisons (HIV+/+ and HIV+/- versus HIV/-) suggest comparable immune responses for most vaccine serotypes and protein D, except a trend for lower post primary OPA response in HIV+/+ group for most vaccine serotypes with unknown clinical relevance. Immunological memory was shown for each vaccine serotype after the booster dose.

A clinical study in Burkina Faso in children with or without SCD receiving vaccination according to their age (<6 months, 7-11 months, 12-23 months of age) suggests that the immunogenicity of **Synflorix** is not influenced by SCD.

Immunogenicity and safety of **Synflorix** were assessed in a limited number of subjects with congenital or acquired asplenia, splenic dysfunction or complement deficiencies: 6 subjects 2-5 years of age and

40 subjects 6-17 years of age (**Synflorix** is indicated up to 5 years of age). **Synflorix** was shown to be immunogenic and no new safety concerns were observed in this study.

Non-clinical Information

A repeated dose toxicity study of pneumococcal conjugate vaccine in rabbit revealed no evidence of any significant local or systemic toxic effects.

PHARMACEUTICAL INFORMATION

List of Excipients

Sodium chloride, water for injections.

Shelf Life

When stored at the recommended temperature of +2°C to +8°C, the shelf life of **Synflorix** is 4 years.

Synflorix should be administered as soon as possible after being removed from the refrigerator. However, stability data generated indicate that **Synflorix** remains stable and can be administered when the vaccine has been stored outside the refrigerator for up to 72 hours at temperatures between 8°C and 25°C.

Storage

Store in a refrigerator (2°C – 8°C).

Do not freeze.

Store in the original packaging in order to protect from light.

Nature and Contents of Container

0.5 mL of suspension in a pre-filled syringe (type I glass) with a plunger stopper (butyl rubber) and with a rubber tip cap.

Pack sizes of 1 with needle.

The tip cap and rubber plunger stopper of the pre-filled syringe are not made with natural rubber latex.

Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

Use and Handling


A fine white deposit with a clear colourless supernatant may be observed upon storage of the syringe/vial. This does not constitute a sign of deterioration.

The content of the syringe/vial should be inspected visually both before and after shaking for any foreign particulate matter and/or abnormal physical appearance prior to administration.

In the event of either being observed, discard the vaccine.


The vaccine should be well shaken before use.

Instructions for the pre-filled syringe



Hold the syringe by the barrel, not by the plunger.

Unscrew the syringe cap by twisting it anticlockwise.



To attach the needle, connect the hub to the Luer Lock Adaptor and rotate a quarter turn clockwise until you feel it lock.

Do not pull the syringe plunger out of the barrel. If it happens, do not administer the vaccine.

Disposal

Any unused product or waste material should be disposed of in accordance with local requirements.

Presentations

Box, 1 pre-filled syringe @ 0.5 mL (1 dose) + needle

Reg. No.: DKI2276703643A1

HARUS DENGAN RESEP DOKTER

Imported by

PT Glaxo Wellcome Indonesia
Jakarta, Indonesia

Manufactured by

GlaxoSmithKline Biologicals s.a.
Rixensart, Belgium

Version number: GDS18/IPI18

Date of issue: 12 August 2022

Trademarks are owned by or licensed to the GSK group of companies.

©2022 GSK group of companies or its licensor.

Informasi Produk Untuk Pasien

**Synflorix**

Suspensi untuk Injeksi

Pneumococcal polysaccharide and Non-Typeable *Haemophilus influenzae* (NTHi) protein D conjugate vaccine, adsorbed**Baca brosur ini dengan saksama sebelum anak Anda mendapatkan vaksin ini.**

- Simpan brosur ini. Anda mungkin butuh untuk membacanya lagi.
- Jika Anda memiliki pertanyaan lebih lanjut, Anda dapat menghubungi dokter atau apoteker Anda.
- Vaksin ini diresepkan untuk anak Anda. Jangan diberikan untuk orang lain.
- Jika kejadian ikutan pasca imunisasi (KIP) menjadi serius atau jika Anda mendapatkan kejadian ikutan pasca imunisasi (KIP) yang tidak terdapat pada brosur ini, harap hubungi dokter atau apoteker Anda.

Pada brosur ini:

1. Apakah itu **Synflorix** dan digunakan untuk apa
2. Sebelum anak Anda mendapatkan **Synflorix**
3. Bagaimana cara pemberian **Synflorix**
4. Kemungkinan kejadian ikutan pasca imunisasi (KIP)
5. Isi kemasan dan informasi lain

1. Apakah itu *Synflorix* dan digunakan untuk apa

Synflorix adalah vaksin yang akan membantu melindungi anak Anda terhadap penyakit yang disebabkan oleh beberapa jenis bakteri yang disebut *Streptococcus pneumoniae* serotipe 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F. Bakteri ini dapat menyebabkan penyakit serius termasuk meningitis, infeksi darah, pneumonia dan infeksi telinga.

Vaksin bekerja dengan membantu tubuh untuk membuat antibodi sendiri, yang melindungi anak Anda dari penyakit-penyakit tersebut.

Seperti halnya semua vaksin, **Synflorix** mungkin tidak sepenuhnya melindungi semua anak yang divaksinasi.

Synflorix hanya akan melindungi terhadap infeksi yang disebabkan oleh kelompok *Streptococcus pneumoniae* yang menjadi alasan dikembangkannya vaksin ini.

Anak-anak dengan sistem kekebalan yang lemah, misalnya karena infeksi *Human Immunodeficiency Virus* (HIV), mungkin tidak mendapatkan manfaat penuh dari **Synflorix**.

2. Sebelum anak Anda mendapatkan *Synflorix****Synflorix* sebaiknya tidak diberikan jika anak Anda:**

- Sebelumnya pernah mengalami reaksi alergi terhadap **Synflorix** atau bahan apa pun yang terkandung dalam **Synflorix**. Bahan-bahan dalam **Synflorix** tercantum di bagian akhir brosur. Tanda-tanda reaksi alergi mungkin termasuk ruam kulit yang gatal, sesak napas dan pembengkakan wajah atau lidah.

Memerlukan perhatian khusus pada *Synflorix*:

Sebelum anak Anda divaksinasi, pastikan dokter Anda tahu jika hal-hal berikut berlaku untuk anak Anda:

- Memiliki infeksi berat dengan suhu tinggi. Mungkin perlu untuk menunda vaksinasi sampai pemulihannya. Infeksi ringan seperti pilek seharusnya tidak menjadi masalah, tetapi konsultasikan dengan dokter Anda terlebih dahulu.
- Memiliki masalah perdarahan atau mudah memar.
- Mengalami kesulitan bernapas, silakan hubungi dokter Anda. Ini mungkin lebih sering terjadi dalam tiga hari pertama setelah vaksinasi jika anak Anda lahir prematur (sebelum atau pada kehamilan 28 minggu).

Pingsan dapat terjadi setelah, atau bahkan sebelum penyuntikan apa pun, oleh karena itu beri tahu dokter atau perawat jika anak Anda pingsan saat penyuntikan sebelumnya.

Menggunakan obat atau vaksin lain

Konsultasikan dengan dokter Anda jika anak Anda sedang atau baru saja selesai menggunakan obat lain, termasuk obat-obatan yang diperoleh tanpa resep atau baru saja menerima vaksin lain.

Synflorix tidak berfungsi sebaik yang diinginkan jika anak Anda mengonsumsi obat yang mengurangi efektivitas sistem kekebalan tubuh dalam melawan infeksi.

Synflorix dapat diberikan bersamaan dengan vaksin anak lainnya. Tempat injeksi yang berbeda digunakan untuk setiap jenis vaksin.

3. Bagaimana cara pemberian *Synflorix*

Bayi dari usia 2 sampai 6 bulan:

Biasanya, anak Anda akan menerima tiga suntikan dengan selang waktu setidaknya satu bulan antar suntikan. Suntikan pertama biasanya diberikan pada usia 2 bulan dengan selang waktu satu bulan antar suntikan. Setidaknya enam bulan setelah suntikan terakhir anak Anda akan menerima suntikan tambahan (*booster*).

Atau, anak Anda mungkin menerima dua suntikan dengan selang waktu dua bulan antara suntikan. Suntikan pertama dapat diberikan mulai usia 2 bulan ke atas. Setidaknya enam bulan setelah suntikan terakhir, anak Anda akan menerima suntikan tambahan (*booster*).

Bayi prematur

Anak Anda akan menerima tiga suntikan dengan selang waktu setidaknya satu bulan antara setiap dosis. Setidaknya enam bulan setelah suntikan terakhir, anak Anda akan menerima suntikan tambahan (*booster*).

Bayi dan Anak yang belum divaksinasi sebelumnya

- Bayi usia 7-11 bulan:
anak Anda akan menerima dua suntikan dengan selang waktu setidaknya satu bulan antara suntikan. Setidaknya dua bulan setelah suntikan terakhir dan selama tahun kedua umurnya, anak Anda akan menerima suntikan ketiga (*booster*).
- Anak-anak berusia 12 bulan - 5 tahun:
anak Anda akan menerima total dua suntikan dengan selang waktu setidaknya dua bulan antara suntikan.

Populasi khusus

Keamanan dan imunogenisitas **Synflorix** pada anak-anak dengan kondisi *immunocompromising* yang mendasari penyakit invasif pneumokokus (seperti SCD, atau infeksi HIV) telah diteliti pada anak-anak di bawah usia 2 tahun. Tersedia data yang terbatas untuk anak-anak dengan asplenia atau disfungsi limpa usia 2 sampai 5 tahun. **Synflorix** boleh diberikan sesuai jadwal yang disebutkan di atas, kecuali jadwal 3 dosis harus diberikan sebagai vaksinasi primer pada bayi mulai vaksinasi dari usia 2 bulan hingga 6 bulan. Konsultasikan dengan dokter Anda untuk informasi tentang jumlah dan waktu suntikan untuk anak Anda.

Synflorix selalu disuntikkan ke otot, biasanya di paha atau lengan atas.

Anda akan diberitahu ketika anak Anda harus kembali untuk suntikan berikutnya. Jika anak Anda tidak mendapatkan suntikan yang dijadwalkan, penting untuk Anda menjadwalkan di hari lain.

Pastikan anak Anda menyelesaikan jadwal vaksinasi lengkap.

4. Kemungkinan kejadian ikutan pasca imunisasi (KIP)

Seperti semua obat-obatan, **Synflorix** dapat menyebabkan kejadian ikutan pasca imunisasi (KIP), meskipun tidak semua orang mendapatkannya.

Kejadian ikutan pasca imunisasi (KIP) yang terjadi selama uji klinis dengan **Synflorix** adalah sebagai berikut:

Sangat umum (terjadi lebih dari 1 dari 10 dosis vaksin):

- Nyeri, kemerahan dan Bengkak di tempat suntikan
- Demam 38°C atau lebih tinggi (untuk umur di bawah 2 tahun)

- Mengantuk
- Irritabilitas
- Kehilangan selera makan

Umum (terjadi hingga 1 dari 10 dosis vaksin):

- Nyeri dan Bengkak di tempat suntikan
- Demam lebih tinggi dari 39°C (untuk umur di bawah 2 tahun)

Tidak umum (terjadi hingga 1 dari 100 dosis vaksin):

- Gatal, pembekuan darah, perdarahan, dan benjolan kecil di tempat suntikan
- Mual, diare, muntah
- Menangis tanpa alasan
- Henti napas sementara (*apnoea*)
- Sakit kepala
- Ruam kulit

Jarang (terjadi hingga 1 dari 1.000 dosis vaksin):

- Reaksi alergi, seperti alergi kulit
- Kejang dengan atau tanpa demam

Sangat jarang (ini dapat terjadi hingga 1 dari 10.000 dosis vaksin)

- Reaksi alergi yang parah yang dapat dikenali:
 - biduran dan ruam gatal (gatal-gatal)
 - bengkak, kadang-kadang di wajah atau mulut (*angioedema*), menyebabkan kesulitan bernafas-kehilangan kesadaran.
- Reaksi-reaksi ini biasanya akan terjadi sebelum meninggalkan ruangan dokter. Namun, jika anak Anda mengalami gejala-gejala ini, Anda harus segera menghubungi dokter.
- Penyakit Kawasaki (tanda-tanda utama penyakit tersebut misalnya: demam yang berlangsung selama lebih dari lima hari, yang dikaitkan dengan ruam pada dada, perut dan punggung (*trunk*) diikuti dengan pengelupasan kulit pada tangan dan jari, bengkaknya kelenjar di leher, mata merah, bibir, tenggorokan dan lidah).

Jika anak Anda berusia lebih dari 12 bulan ketika dia menerima suntikan tambahan (*booster*), dia lebih mungkin mengalami reaksi di tempat suntikan.

Jika Anda mengalami kejadian ikutan pasca imunisasi (KIP)

Jika salah satu dari kejadian ikutan pasca imunisasi (KIP) ini menjadi serius atau Anda menyadari adanya kejadian ikutan pasca imunisasi (KIP) yang tidak terdapat dalam brosur ini, beritahukan dokter atau apoteker Anda.

5. Isi kemasan dan Informasi lain

Apa kandungan *Synflorix*

Setiap dosis 0.5 mL mengandung 1 mikrogram sakarida untuk serotype 1, 5, 6B, 7F, 9V, 14 dan 23F, dan 3 mikrogram sakarida untuk serotype 4, 18C dan 19F.

Bahan lainnya;

Sodium chloride, water for injections.

Bagaimana bentuk dari *Synflorix* dan Isi kemasan

Synflorix adalah suspensi injeksi dalam kemasan sebagai berikut:

- kemasan 1 atau 10 *pre-filled syringes* dengan atau tanpa jarum terpisah
- kemasan 1, 10 atau 100 vial

Tidak semua kemasan dipasarkan.

Diproduksi oleh:

GlaxoSmithKline Biologicals s.a.
89, rue de l'Institut - 1330 Rixensart
Belgia

Diimpor oleh:

PT Glaxo Wellcome Indonesia
Jakarta, Indonesia

HARUS DENGAN RESEP DOKTER

Dus, 1 pfs @ 0.5 mL (1 dosis) + 1 jarum suntik Reg. No. XXXXXXXXXXXXXXXXX

Merek dagang dimiliki oleh atau dilisensikan kepada grup perusahaan GSK.
©2021 grup perusahaan GSK atau pemberi lisensinya.

Version 4 – Sep 2021