



COMPOSITION

CoAprovel 150 mg /12.5 mg Film coated tablets

Each film-coated tablet contains 150 mg ibresartan and 12.5 mg hydrochlorothiazide.

CoAprovel 300 mg/12.5 mg Film coated tablets

Each film-coated tablet contains 300 mg ibresartan and 12.5 mg hydrochlorothiazide.

PHARMACEUTICAL FORM

150 mg/12.5 mg film coated tablet

Film-coated tablet: Peach, biconvex, oval-shaped, with a heart debossed on one side and the number <2875> engraved on the other side.

300 mg/12.5 mg film coated tablet

Film-coated tablet: Peach, biconvex, oval-shaped, with a heart debossed on one side and the number <2876> engraved on the other side.

CLINICAL PARTICULARS

Therapeutic indications

Treatment of essential hypertension.

This fixed dose combination is indicated in patients whose blood pressure is not adequately controlled on ibresartan or hydrochlorothiazide alone.

This fixed dose combination is also indicated as initial treatment when hypertension is severe and rapid control of blood pressure (within days to weeks) is of primary clinical importance (see "Pharmacodynamic Properties").

Posology and method of administration

CoAprovel can be taken once daily, with or without food.

Dose titration with the individual components (i.e. ibresartan and hydrochlorothiazide) may be recommended.

When clinically appropriate direct change from monotherapy to the fixed combinations may be considered:

- CoAprovel 150 mg/12.5 mg may be administered in patients whose blood pressure is not adequately controlled with hydrochlorothiazide or ibresartan 150 mg alone;
- CoAprovel 300 mg/12.5 mg may be administered in patients insufficiently controlled by ibresartan 300 mg or by CoAprovel 150 mg/12.5 mg;
- CoAprovel 300 mg/25 mg may also be administered in patients insufficiently controlled by CoAprovel 300 mg/12.5 mg.

Doses higher than 300 mg ibresartan/25 mg hydrochlorothiazide once daily are not recommended.

When necessary, CoAprovel may be administered with another antihypertensive medicinal product (see "Interaction with other medicinal products and other forms of interaction").

Renal impairment: due to the hydrochlorothiazide component, CoAprovel is not recommended for patients with severe renal impairment (creatinine clearance < 30 ml/min). Loop diuretics are preferred to thiazides in this population. No dosage adjustment is necessary in patients with renal impairment whose renal creatinine clearance is ≥ 30 ml/min (see "Contraindications" and "Special warnings and special precautions for use").

Hepatic impairment: CoAprovel is not indicated in patients with severe hepatic impairment. Thiazides should be used with caution in patients with impaired hepatic function. No dosage adjustment of CoAprovel is necessary in patients with mild to moderate hepatic impairment (see "Contraindications").

Intravascular volume depletion: volume and/or sodium depletion should be corrected prior to administration of CoAprovel (see "Special warnings and special precautions for use, Hypotension - Volume-depleted patients").

Elderly patients: no dosage adjustment of CoAprovel is necessary in elderly patients.

Paediatric patients: CoAprovel is not recommended for use in children and adolescents due to a lack of data on safety and efficacy.

Contraindications

- Hypersensitivity to the active substances, to any of the excipients (see "List of excipients"), or to other sulfonamide-derived substances (hydrochlorothiazide is a sulfonamide-derived substance)
- Second and third trimester of pregnancy (see "Special warnings and special precautions for use" and "Pregnancy and lactation")
- Severe renal impairment (creatinine clearance < 30 ml/min)
- Refractory hypokalaemia, hypercalcaemia
- Severe hepatic impairment, biliary cirrhosis and cholestasis
- The concomitant use of CoAprovel with aliskiren-containing products is contraindicated in patients with diabetes mellitus or renal impairment (glomerular filtration rate (GFR) <60 ml/min/1.73 m²) (see "Interaction with other medicinal products and other forms of interaction" and "Pharmacodynamic properties")

DISETUVUI OLEH BPOM : 21/06/2021

Special warnings and special precautions for use

Hypotension - Volume-depleted patients: CoAprovel has been rarely associated with symptomatic hypotension in hypertensive patients without other risk factors for hypotension. Symptomatic hypotension may be expected to occur in patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea or vomiting. Such conditions should be corrected before initiating therapy with CoAprovel.

Renal artery stenosis - Renovascular hypertension: there is an increased risk of severe hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with angiotensin converting enzyme inhibitors or angiotensin-II receptor antagonists. While this is not documented with CoAprovel, a similar effect should be anticipated.

Renal impairment and kidney transplantation: when CoAprovel is used in patients with impaired renal function, a periodic monitoring of potassium, creatinine and uric acid serum levels is recommended. There is no experience regarding the administration of CoAprovel in patients with a recent kidney transplantation. CoAprovel should not be used in patients with severe renal impairment (creatinine clearance < 30 ml/min) (see "Contraindications"). Thiazide diuretic-associated azotemia may occur in patients with impaired renal function. No dosage adjustment is necessary in patients with renal impairment whose creatinine clearance is ≥ 30 ml/min. However, in patients with mild to moderate renal impairment (creatinine clearance ≥ 30 ml/min but < 60 ml/min) this fixed dose combination should be administered with caution.

Dual blockade of the renin-angiotensin-aldosterone system (RAAS):

There is evidence that the concomitant use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren increases the risk of hypotension, hyperkalaemia and decreased renal function (including acute renal failure). Dual blockade of RAAS through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is therefore not recommended (see "Interactions with other medicinal products and other forms of interaction" and "Pharmacodynamic properties"). If dual blockade therapy is considered absolutely necessary, this should only occur under specialist supervision and subject to frequent close monitoring of renal function, electrolytes and blood pressure. ACE-inhibitors and angiotensin II receptor blockers should not be used concomitantly in patients with diabetic nephropathy.

Psoriasis: The use of in patients with psoriasis or a history of psoriasis should be carefully weighed as it may exacerbate psoriasis.

Hepatic impairment: thiazides should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma. There is no clinical experience with CoAprovel in patients with hepatic impairment.

Aortic and mitral valve stenosis, obstructive hypertrophic cardiomyopathy: as with other vasodilators, special caution is indicated in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy.

Primary aldosteronism: patients with primary aldosteronism generally will not respond to anti-hypertensive medicinal products acting through inhibition of the renin-angiotensin system. Therefore, the use of CoAprovel is not recommended.

Metabolic and endocrine effects: thiazide therapy may impair glucose tolerance. In diabetic patients dosage adjustments of insulin or oral hypoglycemic agents may be required.

Latent diabetes mellitus may become manifest during thiazide therapy.

Increases in cholesterol and triglyceride levels have been associated with thiazide diuretic therapy; however at the 12.5 mg dose contained in CoAprovel, minimal or no effects were reported.

Hyperuricaemia may occur or frank gout may be precipitated in certain patients receiving thiazide therapy.

Electrolyte imbalance: as for any patient receiving diuretic therapy, periodic determination of serum electrolytes should be performed at appropriate intervals.

Thiazides, including hydrochlorothiazide, can cause fluid or electrolyte imbalance (hypokalaemia, hyponatraemia, and hypochloraemic alkalosis). Warning signs of fluid or electrolyte imbalance are dryness of mouth, thirst, weakness, lethargy, drowsiness, restlessness, muscle pain or cramps, muscular fatigue, hypotension, oliguria, tachycardia, and gastrointestinal disturbances such as nausea or vomiting.

Although hypokalaemia may develop with the use of thiazide diuretics, concurrent therapy with ibresartan may reduce diuretic-induced hypokalaemia. The risk of hypokalaemia is greatest in patients with cirrhosis of the liver, in patients experiencing brisk diuresis, in patients who are receiving inadequate oral intake of electrolytes and in patients receiving concomitant therapy with corticosteroids or ACTH. Conversely, due to the ibresartan component of CoAprovel hyperkalaemia might occur, especially in the presence of renal impairment and/or heart failure, and diabetes mellitus. Adequate monitoring of serum potassium in patients at risk is recommended. Potassium-sparing diuretics, potassium supplements or potassium-containing salts substitutes should be co-administered cautiously with CoAprovel (see "Interaction with other medicinal products and other forms of interaction").

There is no evidence that ibresartan would reduce or prevent diuretic-induced hyponatraemia. Chloride deficit is generally mild and usually does not require treatment.

Thiazides may decrease urinary calcium excretion and cause an intermittent and slight elevation of serum calcium in the absence of known disorders of calcium metabolism.

Non-steroidal anti-inflammatory drugs: when angiotensin II antagonists are administered simultaneously with non-steroidal antiinflammatory drugs (i.e. selective COX-2 inhibitors, acetylsalicylic acid (> 3 g/day) and non-selective NSAIDs), attenuation of the antihypertensive effect may occur.

As with ACE inhibitors, concomitant use of angiotensin II antagonists and NSAIDs may lead to an increased risk of worsening of renal function, including possible acute renal failure, and an increase in serum potassium, especially in patients with poor pre-existing renal function. The combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring renal function after initiation of concomitant therapy, and periodically thereafter.

Additional information on ibresartan interactions: in clinical studies, the pharmacokinetic of ibresartan is not affected by hydrochlorothiazide. Ibresartan is mainly metabolized by CYP2C9 and to a lesser extent by glucuronidation. No significant pharmacokinetic or pharmacodynamic interactions were observed when ibresartan was coadministered with warfarin, a medicinal product metabolized by CYP2C9. The effects of CYP2C9 inducers such as rifampicin on the pharmacokinetic of ibresartan have not been evaluated. The pharmacokinetic of digoxin was not altered by co-administration of ibresartan.

Additional information on hydrochlorothiazide interactions: when administered concurrently, the following medicinal products may interact with thiazide diuretics:

Alcohol: potentiation of orthostatic hypotension may occur.

Antidiabetic medicinal products (oral agents and insulins): dosage adjustment of the antidiabetic medicinal product may be required (see "Special warnings and precautions for use").

Colestyramine and Colestipol resins: absorption of hydrochlorothiazide is impaired in the presence of anionic exchange resins. CoAprovel should be taken at least one hour before or four hours after these medications.

Corticosteroids, ACTH: electrolyte depletion, particularly hypokalaemia, may be increased; **Digitalis glycosides:** thiazide induced hypokalaemia or hypomagnesaemia favour the onset of hypersensitivity reactions to hydrochlorothiazide may occur in patients with or without a history of allergy or bronchial asthma, but are more likely in patients with such a history.

Exacerbation or activation of systemic lupus erythematosus has been reported with the use of

Thiazide diuretics.

Cases of photosensitivity reactions have been reported with thiazides diuretics (see "Undesirable effects"). If photosensitivity reaction occurs during treatment, it is recommended to stop the treatment. If a re-administration of the diuretic is deemed necessary, it is recommended to protect exposed areas to the sun or to artificial UVA.

Pregnancy: angiotensin II receptor antagonists including CoAprovel should not be initiated during pregnancy. Unless continued therapy with angiotensin II receptor antagonists is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with angiotensin II receptor antagonists should be stopped immediately, and, if appropriate, alternative therapy should be started (see "Contraindications" and "Pregnancy and lactation").

Acute Myopia and Secondary Acute Angle-Closure Glaucoma: sulfonamide drugs or sulfonamide derivative drugs can cause an idiosyncratic reaction, resulting in transient myopia and acute angle-closure glaucoma. While hydrochlorothiazide is a sulfonamide, only isolated cases of acute angle-closure glaucoma have been reported so far with hydrochlorothiazide. Symptoms include acute onset of decreased visual acuity or ocular pain and typically occur within hours to weeks of drug initiation. Untreated acute angle-closure glaucoma can lead to permanent vision loss.

The primary treatment is to discontinue drug intake as rapidly as possible. Prompt medical or surgical treatments may need to be considered if the intraocular pressure remains uncontrolled. Risk factors for developing acute angle-closure glaucoma may include a history of sulfonamide or penicillin allergy (see "Undesirable effects").

Lactose: Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Non-melanoma skin cancer:

An increased risk of non-melanoma skin cancer (NMSC) [basal cell carcinoma (BCC) and squamous cell carcinoma (SCC)] with increasing cumulative dose of hydrochlorothiazide (HCTZ) exposure has been observed in two epidemiological studies based on the Danish National Cancer Registry. Photosensitizing actions of HCTZ could act as a possible mechanism for NMSC. Patients taking HCTZ should be informed of the risk of NMSC and advised to regularly check their skin for any new lesions and promptly report any suspicious skin lesions. Possible preventive measures such as limited exposure to sunlight and UV rays and, in case of exposure, adequate protection should be advised to the patients in order to minimize the risk of skin cancer. Suspicious skin lesions should be promptly examined potentially including histological examinations of biopsies. The use of HCTZ may also need to be reconsidered in patients who have experienced previous NMSC (see also section Undesirable effects).

Interaction with other medicinal products and other forms of interaction

Other antihypertensive agents: the antihypertensive effect of CoAprovel may be increased with the concomitant use of other antihypertensive agents. Irbesartan and hydrochlorothiazide (at doses up to 300 mg ibresartan/25 mg hydrochlorothiazide) have been safely administered with other antihypertensive agents including calcium channel blockers and beta-adrenergic blockers.

Prior treatment with high dose diuretics: may result in volume depletion and a risk of hypotension when initiating therapy with ibresartan with or without thiazide diuretics unless the volume depletion is corrected first (see "Special warnings and precautions for use").

Aliskiren-containing products or ACE-inhibitors: Clinical trial data has shown that dual blockade of the renin-angiotensin-aldosterone system (RAAS) through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is associated with a higher frequency of adverse events such as hypotension, hyperkalaemia and decreased renal function (including acute renal failure) compared to the use of a single RAAS-acting agent (see "Contraindications", "Special warnings and precautions for use" and "Pharmacodynamic properties").

Lithium: reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with angiotensin converting enzyme inhibitors. Similar effects have been very rarely reported with ibresartan so far. Furthermore, renal clearance of lithium is reduced by thiazides so the risk of lithium toxicity could be increased with CoAprovel. Therefore, the combination of lithium and CoAprovel is not recommended (see "Special warnings and precautions for use"). If the combination proves necessary, careful monitoring of serum lithium levels is recommended.

Medicinal products affecting potassium: the potassium-depleting effect of hydrochlorothiazide is attenuated by the potassium-sparing effect of ibresartan. However, this effect of hydrochlorothiazide on serum potassium would be expected to be potentiated by other medicinal products associated with potassium loss and hypokalaemia (e.g. other kaliuretic diuretics, laxatives, amphotericin, carbenoxolone, penicillin G sodium). Conversely, based on the experience with the use of other medicinal products that blunt the renin-angiotensin system, concomitant use of potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium or other medicinal products that may increase serum potassium levels (e.g. heparin sodium) may lead to increases in serum potassium. Adequate monitoring of serum potassium in patients at risk is recommended (see "Special warnings and precautions for use").

Medicinal products affected by serum potassium disturbances: periodic monitoring of serum potassium is recommended when CoAprovel is administered with medicinal products affected by serum potassium disturbances (e.g. digitalis glycosides, antiarrhythmics).

Non-steroidal anti-inflammatory drugs: when angiotensin II antagonists are administered simultaneously with non-steroidal antiinflammatory drugs (i.e. selective COX-2 inhibitors, acetylsalicylic acid (> 3 g/day) and non-selective NSAIDs), attenuation of the antihypertensive effect may occur.

As with ACE inhibitors, concomitant use of angiotensin II antagonists and NSAIDs may lead to an increased risk of worsening of renal function, including possible acute renal failure, and an increase in serum potassium, especially in patients with poor pre-existing renal function. The combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring renal function after initiation of concomitant therapy, and periodically thereafter.

Additional information on ibresartan interactions: in clinical studies, the pharmacokinetic of ibresartan is not affected by hydrochlorothiazide. Ibresartan is mainly metabolized by CYP2C9 and to a lesser extent by glucuronidation. No significant pharmacokinetic or pharmacodynamic interactions were observed when ibresartan was coadministered with warfarin, a medicinal product metabolized by CYP2C9. The effects of CYP2C9 inducers such as rifampicin on the pharmacokinetic of ibresartan have not been evaluated. The pharmacokinetic of digoxin was not altered by co-administration of ibresartan.

Additional information on hydrochlorothiazide interactions: when administered concurrently, the following medicinal products may interact with thiazide diuretics:

Alcohol: potentiation of orthostatic hypotension may occur.

Antidiabetic medicinal products (oral agents and insulins): dosage adjustment of the antidiabetic medicinal product may be required (see "Special warnings and precautions for use").

Colestyramine and Colestipol resins: absorption of hydrochlorothiazide is impaired in the presence of anionic exchange resins. CoAprovel should be taken at least one hour before or four hours after these medications.

Corticosteroids, ACTH: electrolyte depletion, particularly hypokalaemia, may be increased; **Digitalis glycosides:** thiazide induced hypokalaemia or hypomagnesaemia favour the onset of hypersensitivity reactions to hydrochlorothiazide may occur in patients with or without a history of allergy or bronchial asthma, but are more likely in patients with such a history.

Exacerbation or activation of systemic lupus erythematosus has been reported with the use of

digitalis-induced cardiac arrhythmias (see "Special warnings and precautions for use"); **Non-steroidal anti-inflammatory drugs:** the administration of a non-steroidal anti-inflammatory drug may reduce the diuretic, natriuretic and antihypertensive effects of thiazide diuretics in some patients;

Pressor amines (e.g. noradrenaline): the effect of pressor amines may be decreased, but not sufficiently to preclude their use;

Nondepolarizing skeletal muscle relaxants (e.g. tubocurarine): the effect of nondepolarizing skeletal muscle relaxants may be potentiated by hydrochlorothiazide;

Antigout medicinal products: dosage adjustments of antigout medicinal products may be necessary as hydrochlorothiazide may raise the level of serum uric acid. Increase in dosage of probenecid or sulfinpyrazone may be necessary. Co-administration of thiazide diuretics may

increase the incidence of hypersensitivity reactions to allopurinol;

Calcium salts: thiazide diuretics may increase serum calcium levels due to decreased excretion. If calcium supplements or calcium sparing medicinal products (e.g. Vitamin D therapy) must be prescribed, serum calcium levels should be monitored and calcium dosage adjusted accordingly;

Carbamazepine: concomitant use of carbamazepine and hydrochlorothiazide has been associated with the risk of symptomatic hyponatraemia. Electrolytes should be monitored during concomitant use. If possible, another class of diuretics should be used;

Other interactions: the hyperglycaemic effect of beta-blockers and diazoxide may be enhanced by thiazides. Anticholinergic agents (e.g. atropine, beperiden) may increase the bioavailability of thiazide-type diuretics by decreasing gastrointestinal motility and stomach emptying rate. Thiazides may increase the risk of adverse effects caused by amantadine. Thiazides may reduce the renal excretion of cytotoxic medicinal products (e.g. cyclophosphamide, methotrexate) and potentiate their myelosuppressive effects.

Pregnancy and lactation

Pregnancy: CoAprovel is not recommended during the first trimester of pregnancy (see "Special warnings and precautions for use"). The use of angiotensin II receptor antagonists is contraindicated during the second and third trimester of pregnancy (see "Contraindications" and "Special warnings and precautions for use").

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Whilst there is no controlled epidemiological data on the risk with Angiotensin II Receptor Antagonists (AIIAs), similar risks may exist for this class of drugs. Unless continued AIIA therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIAs should be stopped immediately, and, if appropriate, alternative therapy should be started. Exposure to AIIA therapy during the second and third trimesters is known to induce human foetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia) (see "Preclinical safety data").

Should exposure to AIIAs have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended. Infants whose mothers have taken AIIAs should be closely observed for hypotension (see "Contraindications" and "Special warnings and precautions for use").

Thiazides cross the placental barrier and appear in cord blood. They may cause a decrease in placental perfusion, foetal electrolyte disturbances and possibly other reactions that have occurred in the adults. Cases of neonatal thrombocytopenia, or foetal or neonatal hypocalcaemia have been reported with maternal thiazide therapy. Since CoAprovel contains hydrochlorothiazide, it is not recommended during the first trimester of pregnancy. A switch to a suitable alternative treatment should be carried out in advance of a planned pregnancy.

Breast-feeding: Because no information is available regarding the use of CoAprovel during breast-feeding, CoAprovel is not recommended and alternative treatments with better established safety profiles during breastfeeding are preferable, especially while nursing a newborn or preterm infant. It is unknown whether ibresartan or its metabolites are excreted in human milk. Available pharmacodynamic/toxicological data in rats have shown excretion of ibresartan or its metabolites in milk (for details see "Preclinical safety data"). Hydrochlorothiazide is excreted in human milk in small amounts. Thiazides in high doses causing intense diuresis can inhibit the milk production. The use of CoAprovel during breast feeding is not recommended. If CoAprovel is used during breast feeding, doses should be kept as low as possible.

Ibresartan had no effect upon fertility of treated rats and their offspring up to the dose levels inducing the first signs of parental toxicity (see section "Preclinical safety data").

Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. Based on its pharmacodynamic properties, CoAprovel is unlikely to affect this ability. When driving vehicles or operating machines, it should be taken into account that occasionally dizziness or weariness may occur during treatment of hypertension.

Undesirable effects

Ibresartan

Table 1: Adverse Reactions in Placebo-Controlled Trials and Spontaneous Reports*

<i>Respiratory, thoracic and mediastinal disorders:</i>	Not known:	cough
<i>Gastrointestinal disorders:</i>	Common:	nausea/vomiting
	Uncommon:	diarrhoea
	Not known:	dyspepsia, dysgeusia
<i>Renal and urinary disorders:</i>	Common:	abnormal urination
	Uncommon:	impaired renal function including isolated cases of renal failure in patients at risk (see "Special warnings and special precautions for use")
<i>Musculoskeletal and connective tissue disorders:</i>	Common:	swelling extremity
	Not known:	arthralgia, myalgia
<i>Metabolism and nutrition disorders:</i>	Not known:	hyperkalaemia
<i>Vascular disorders:</i>	Uncommon:	flushing
<i>General disorders and administration site conditions:</i>	Common:	fatigue
<i>Immune system disorders:</i>	Not known:	cases of hypersensitivity reactions such as angioedema, rash, urticaria
<i>Hepatobiliary disorders:</i>	Not known:	hepatitis, abnormal liver function
<i>Reproductive system and breast disorders:</i>	Uncommon:	sexual dysfunction, libido changes

*Frequency for adverse reactions detected by spontaneous reports is described as "not known"

Additional information on individual components: in addition to the adverse reactions listed above for the combination product, other adverse reactions previously reported with one of the individual components may be potential adverse reactions with CoAprovel. Tables 2 and 3 below detail the adverse reactions reported with the individual components of CoAprovel.

Table 2: Adverse reactions reported with the use of irbesartan alone

<i>Blood and lymphatic system disorders:</i>	Not known:	thrombocytopenia
<i>General disorders and administration site conditions:</i>	Uncommon:	chest pain
<i>Immune system disorders:</i>	Not known:	Anaphylactic reaction including anaphylactic shock, thrombocytopenic purpura, psoriasis (and psoriasis exacerbation), and photosensitivity

Table 3: Adverse reactions (regardless of relationship to medicinal product) reported with the use of hydrochlorothiazide alone

<i>Investigations:</i>	Not known:	electrolyte imbalance (including hypokalaemia and hyponatraemia, see "Special warnings and special precautions for use"), hyperuricaemia, glycosuria, hyperglycaemia, increases in cholesterol and triglycerides
<i>Cardiac disorders:</i>	Not known:	cardiac arrhythmias
<i>Blood and lymphatic system disorders:</i>	Not known:	aplastic anaemia, bone marrow depression, neutropenia/agranulocytosis, haemolytic anaemia, leucopenia, thrombocytopenia
<i>Nervous system disorders:</i>	Not known:	vertigo, paraesthesia, light-headedness, restlessness
<i>Eye disorders:</i>	Not known:	transient blurred vision, xanthopsia, acute myopia and secondary acute angle-closure glaucoma
<i>Respiratory, thoracic and mediastinal disorders:</i>	Not known:	respiratory distress (including pneumonitis and pulmonary oedema)
<i>Gastrointestinal disorders:</i>	Not known:	pancreatitis, anorexia, diarrhoea, constipation, gastric irritation, sialadenitis, loss of appetite
<i>Renal and urinary disorders:</i>	Not known:	interstitial nephritis, renal dysfunction
<i>Skin and subcutaneous tissue disorders:</i>	Not known:	anaphylactic reactions, toxic epidermal necrolysis, necrotizing angitis (vasculitis, cutaneous vasculitis), cutaneous lupus erythematosus-like reactions, reactivation of cutaneous lupus erythematosus, photosensitivity reactions, rash, urticaria
<i>Musculoskeletal and connective tissue disorders:</i>	Not known:	weakness, muscle spasm
<i>Vascular disorders:</i>	Not known:	postural hypotension
<i>General disorders and administration site conditions:</i>	Not known:	fever
<i>Hepatobiliary disorders:</i>	Not known:	jaundice (intrahepatic cholestatic jaundice)
<i>Psychiatric disorders:</i>	Not known:	depression, sleep disturbances
<i>Neoplasms benign, malignant and unspecified (incl cysts and polyps)</i>	Not known:	non-melanoma skin cancer (basal cell carcinoma and squamous cell carcinoma)

Non-melanoma skin cancer: Based on available data from epidemiological studies, cumulative dose dependent association between HCTZ and NMSC has been observed (see also sections Special warnings and precautions for use, Pharmacodynamic).

The dose dependent adverse events of hydrochlorothiazide (Particularly electrolyte disturbances) may increase, when titrating the hydrochlorothiazide.

Reporting of suspected adverse reactions
Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

Overdose

No specific information is available on the treatment of overdosage with CoAprovel. The patient should be closely monitored, and the treatment should be symptomatic and supportive. Management depends on the time since ingestion and the severity of the symptoms. Suggested measures include induction of emesis and/or gastric lavage. Activated charcoal may be useful in the treatment of overdosage. Serum electrolytes and creatinine should be monitored frequently. If hypotension occurs, the patient should be placed in a supine position, with salt and volume

replacements given quickly.

The most likely manifestations of irbesartan overdosage are expected to be hypotension and tachycardia; bradycardia might also occur. Overdosage with hydrochlorothiazide is associated with electrolyte depletion (hypokalaemia, hypochloremia, hyponatraemia) and dehydration resulting from excessive diuresis. The most common signs and symptoms of overdosage are nausea and somnolence. Hypokalaemia may result in muscle spasms and/or accentuate cardiac arrhythmias associated with the concomitant use of digitalis glycosides or certain anti-arrhythmic medicinal products. Irbesartan is not removed by haemodialysis. The degree to which hydrochlorothiazide is removed by haemodialysis has not been established.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic properties

Pharmacotherapeutic group: angiotensin-II antagonists, combinations

ATC code: C09DA04.

Mechanism of action

CoAprovel is a combination of an angiotensin-II receptor antagonist, irbesartan, and a thiazide diuretic, hydrochlorothiazide. The combination of these ingredients has an additive antihypertensive effect, reducing blood pressure to a greater degree than either component alone.

Irbesartan is a potent, orally active, selective angiotensin-II receptor (AT, subtype) antagonist. It is expected to block all actions of angiotensin-II mediated by the AT₁ receptor, regardless of the source or route of synthesis of angiotensin-II. The selective antagonism of the angiotensin-II (AT₁) receptors results in increases in plasma renin levels and angiotensin-II levels, and a decrease in plasma aldosterone concentration. Serum potassium levels are not significantly affected by irbesartan alone at the recommended doses in patients without risk of electrolyte imbalance (see "Special warnings and special precautions for use" and "Interaction with other medicinal products and other forms of interactions"). Irbesartan does not inhibit ACE (kininase-II), an enzyme which generates angiotensin-II and also degrades bradykinin into inactive metabolites. Irbesartan does not require metabolic activation for its activity.

Hydrochlorothiazide is a thiazide diuretic. The mechanism of antihypertensive effect of thiazide diuretics is not fully known. Thiazides affect the renal tubular mechanisms of electrolyte reabsorption, directly increasing excretion of sodium and chloride in approximately equivalent amounts. The diuretic action of hydrochlorothiazide reduces plasma volume, increases plasma renin activity, increases aldosterone secretion, with consequent increases in urinary potassium and bicarbonate loss, and decreases in serum potassium. Presumably through blockade of the renin-angiotensin-aldosterone system, co-administration of irbesartan tends to reverse the potassium loss associated with these diuretics. With hydrochlorothiazide, onset of diuresis occurs in 2 hours, and peak effect occurs at about 4 hours, while the action persists for approximately 6-12 hours.

The combination of hydrochlorothiazide and irbesartan produces dose-related additive reductions in blood pressure across their therapeutic dose ranges. The addition of 12.5 mg hydrochlorothiazide to 300 mg irbesartan once daily in patients not adequately controlled on 300 mg irbesartan alone resulted in further placebo-corrected diastolic blood pressure reductions with 63.067 population controls, using a risk-set sampling strategy. A cumulative dose-response relationship was demonstrated with an adjusted OR 2.1 (95% CI: 1.7-2.6) increasing to OR 3.9 (3.0-4.9) for high use (~25,000 mg) and OR 7.7 (5.7-10.5) for the highest cumulative dose (~100,000 mg) (see also section Special warnings and precautions for use).

Pharmacokinetic properties

Concomitant administration of hydrochlorothiazide and irbesartan has no effect on the pharmacokinetics of either medicinal product.

Absorption

Irbesartan and hydrochlorothiazide are orally active agents and do not require biotransformation for their activity. Following oral administration of CoAprovel, the absolute oral bioavailability is 60-80% and 50-80% for irbesartan and hydrochlorothiazide, respectively. Food does not affect the bioavailability of CoAprovel. Peak plasma concentration occurs at 1.5-2 hours after oral administration for irbesartan and 1-2.5 hours for hydrochlorothiazide.

Distribution

Plasma protein binding of irbesartan is approximately 96%, with negligible binding to cellular blood components. The volume of distribution for irbesartan is 53-93 litres. Hydrochlorothiazide is 68% protein-bound in the plasma, and its apparent volume of distribution is 0.83-1.14 l/kg.

Linearity/non-linearity

Irbesartan exhibits linear and dose proportional pharmacokinetics over the dose range of 10 to 600 mg. A less than proportional increase in oral absorption at doses beyond 600 mg was observed; the mechanism for this is unknown. The total body and renal clearance are 157-176 and 3.0-3.5 ml/min, respectively. The terminal elimination half-life of irbesartan is 11-15 hours.

Steady-state plasma concentrations are attained within 3 days after initiation of a once-daily dosing regimen.

Limited accumulation of irbesartan (<20%) is observed in plasma upon repeated once-daily dosing. In a study, somewhat higher plasma concentrations of irbesartan were observed in female hypertensive patients. However, there was no difference in the half-life and accumulation of irbesartan. No dosage adjustment is necessary in female patients.

Irbesartan AUC and Cmax values were also somewhat greater in elderly subjects (≥ 65 years) than those of young subjects (18-40 years). However the terminal half-life was not significantly altered. No dosage adjustment is necessary in elderly patients. The mean plasma half-life of hydrochlorothiazide reportedly ranges from 5-15 hours.

Biotransformation

Following oral or intravenous administration of ¹⁴C irbesartan, 80-85% of the circulating plasma radioactivity is attributable to unchanged irbesartan. Irbesartan is metabolized by the liver via glucuronide conjugation and oxidation. The major circulating metabolite is irbesartan glucuronide (approximately 6%). *In vitro* studies indicate that irbesartan is primarily oxidized by the cytochrome P450 enzyme CYP2C9; isoenzyme CYP3A4 has negligible effect.

Elimination

Irbesartan and its metabolites are eliminated by both biliary and renal pathways. After either oral or intravenous administration of ¹⁴C irbesartan, about 20% of the radioactivity is recovered in the urine, and the remainder in the faeces. Less than 2% of the dose is excreted in the urine as unchanged irbesartan. Hydrochlorothiazide is not metabolized but is eliminated rapidly by the kidneys. At least 61% of the oral dose is eliminated unchanged within 24 hours. Hydrochlorothiazide crosses the placental but not the blood-brain barrier, and is excreted in breast milk.

Renal impairment: in patients with renal impairment or those undergoing haemodialysis, the pharmacokinetic parameters of irbesartan are not significantly altered. Irbesartan is not removed by haemodialysis. In patients with creatinine clearance < 20 ml/min, the elimination half-life of hydrochlorothiazide was reported to increase to 21 hours.

Hepatic impairment: in patients with mild to moderate cirrhosis, the pharmacokinetic parameters of irbesartan are not significantly altered. Studies have not been performed in patients with severe hepatic impairment.

Preclinical safety data

Irbesartan/hydrochlorothiazide:

The potential toxicity of the irbesartan/hydrochlorothiazide combination after oral administration was evaluated in rats and macaques in studies lasting up to 6 months. There were no toxicological findings observed of relevance to human therapeutic use. The following changes, observed in rats and macaques receiving the irbesartan/hydrochlorothiazide combination at 10/10 and 90/90 mg/kg/day, were also seen with one of the two medicinal products alone and/or were secondary to decreases in blood pressure (no significant toxicologic interactions were observed):

- kidney changes, characterized by slight increases in serum urea and creatinine, and

hyperplasia/hypertrophy of the juxtaglomerular apparatus, which are a direct consequence of the interaction of irbesartan with the renin-angiotensin system;

- slight decreases in erythrocyte parameters (erythrocytes, haemoglobin, haematocrit);
- stomach discolouration, ulcers and focal necrosis of gastric mucosa were observed in few rats in a 6 months toxicity study at irbesartan 90 mg/kg/day, hydrochlorothiazide 90 mg/kg/day, and irbesartan/hydrochlorothiazide 10/10 mg/kg/day. These lesions were not observed in macaques;
- decreases in serum potassium due to hydrochlorothiazide and partly prevented when hydrochlorothiazide was given in combination with irbesartan.

Most of the above mentioned effects appear to be due to the pharmacological activity of irbesartan (blockade of angiotensin-II-induced inhibition of renin release, with stimulation of the renin-producing cells) and occur also with angiotensin converting enzyme inhibitors, hydrochlorothiazide in humans.

No teratogenic effects were seen in rats given irbesartan and hydrochlorothiazide in combination at doses that produced maternal toxicity. The effects of the irbesartan/hydrochlorothiazide combination on fertility have not been evaluated in animal studies, as there is no evidence of adverse effect on fertility in animals or humans with either irbesartan or hydrochlorothiazide when administered alone. However, another angiotensin-II antagonist affected fertility parameters in animal studies when given alone. These findings were also observed with lower doses of this other angiotensin-II antagonist when given in combination with hydrochlorothiazide.

There was no evidence of mutagenicity or clastogenicity with the irbesartan/hydrochlorothiazide combination. The carcinogenic potential of irbesartan and hydrochlorothiazide in combination has not been evaluated in animal studies.

Irbesartan: there was no evidence of abnormal systemic or target organ toxicity at clinically relevant doses. In preclinical safety studies, high doses of irbesartan (≥ 250 mg/kg/day in rats and ≥ 100 mg/kg/day in macaques) caused a reduction of red blood cell parameters (erythrocytes, haemoglobin, haematocrit). At very high doses (≥ 500 mg/kg/day) degenerative changes in the kidneys (such as interstitial nephritis, tubular distension, basophilic tubules, increased plasma concentrations of urea and creatinine) were induced by irbesartan in the rat and the macaque and are considered secondary to the hypotensive effects of the medicinal product which led to decreased renal perfusion. Furthermore, irbesartan-induced hyperplasia/hypertrophy of the juxtaglomerular cells (in rats at ≥ 90 mg/kg/day, in macaques at ≥ 10 mg/kg/day). All of these changes were considered to be caused by the pharmacological action of irbesartan. For therapeutic doses of irbesartan in humans, the hyperplasia/hypertrophy of the renal juxtaglomerular cells does not appear to have any relevance.

There was no evidence of mutagenicity, clastogenicity or carcinogenicity.

Fertility and reproductive performance were not affected in studies of male and female rats even at oral doses of irbesartan causing some parental toxicity (from 50 to 650 mg/kg/day), including mortality at the highest dose. No significant effects on the number of corpora lutea, implants, or live foetuses were observed. Irbesartan did not affect survival, development, or reproduction of offspring. Studies in animals indicate that the radiolabelled irbesartan is detected in rat and rabbit foetuses. Irbesartan is excreted in the milk of lactating rats.

Animal studies with irbesartan showed transient toxic effects (increased renal pelvic cavitation, hydroureter or subcutaneous oedema) in rat foetuses, which were resolved after birth. In rabbits, abortion or early resorption was noted at doses causing significant maternal toxicity, including mortality. No teratogenic effects were observed in the rat or rabbit.

Hydrochlorothiazide: although equivocal evidence for a genotoxic or carcinogenic effect was found in some experimental models, the extensive human experience with hydrochlorothiazide has failed to show an association between its use and an increase in neoplasms.

PHARMACEUTICAL PARTICULARS

List of Excipients

Tablet core: lactose monohydrate, microcrystalline cellulose, croscarmellose sodium, hypromellose, silicon dioxide, magnesium stearate

Film-coating:

- Opadry Pink 32 F 25403 consist of : lactose monohydrate, hypromellose, titanium dioxide (E171), macrogol, red and yellow ferric oxides (E172)

• Carnauba wax

Incompatibilities

Not applicable.

Special precautions for storage

Do not store above 30°C.

Store in the original package.

Shelf life

Do not use later than the date of Expiry.

Nature and contents of container

CoAprovel film coated tablets are packaged in cartons containing 2 blisters of 14 film coated tablets in PVC/PVDC/Aluminium blisters.

Instructions for use and handling, and disposal

No special requirements.

HARUS DENGAN RESEP DOKTER

COAPROVEL 150 mg/12.5 mg Reg. No. DKI 1077402817A1

COAPROVEL 300 mg/12.5 mg Reg. No. DKI 1077402817B1

Manufactured by:

SANOFI WITHROP INDUSTRIE

1, rue de la Vierge - Ambarès et Lagrave

33565 Carbon Blanc Cedex - France

Imported by:

PT. Aventis Pharma

Jakarta, Indonesia

GENERAL

Code : COACoMltID0321v1
Update : 23.03.2021 - ver 1
Local Code SCM : 000000
Current item code : 549150
Product/Item type : CoAprovel range leaflet

Pack Size
Box of : 28s
Blister : 2 x 14s

Country : ID
Languages : English & Bahasa
Market Barcode : 000000000000
Artwork by : Analyn NOVILLAS
Revised by : XXXXXX
Plant : REC SG

Format : 400 x 220 mm
Plant barcode :
Colour(s):
 PAN REFLEX BLUE CVC

Font(s) : Ocean Sans light, bold
Size : 7 pts (mini)
Technical Plans: xxxx