



Product name:

STELARA®

Ustekinumab

DOSAGE FORMS AND STRENGTHS

Ustekinumab is a fully human IgG1κ monoclonal antibody with an approximate molecular weight of 148600 daltons. Ustekinumab is produced by a recombinant cell line cultured by continuous perfusion and is purified by a series of steps that includes measures to inactivate and remove viruses.

STELARA is available in the following presentations:

Solution for injection for subcutaneous administration

Pre-filled Syringe:

45 mg Ustekinumab in 0.5 mL

90 mg/ 1.0 mL

Single-use Vial:

45 mg / 0.5 mL

Solution for intravenous infusion

Single-use Vial:

130 mg / 26 mL

For excipients, see List of Excipients.

CLINICAL INFORMATIONS

Indications

Stelara 45 mg / 0.5 mL & 90 mg / 1.0 mL in Pre-filled Syringe and 45 mg / 0.5 mL in Single-use Vial:

Plaque Psoriasis

Adults

STELARA is indicated for the treatment of moderate to severe plaque psoriasis in adults who failed to respond to, or who have a contraindication to, or are intolerant to other systemic therapies including cyclosporine, methotrexate or PUVA.

Pediatric plaque psoriasis

Stelara, as monotherapy, is indicated for the treatment of moderate to severe plaque psoriasis in children and adolescent patients from the age of 6 years and older, who are candidates for phototherapy or systemic treatment of psoriasis (either naïve or history of previous treatment) or have psoriasis considered inadequately controlled with topical therapy after an adequate dose and duration of therapy (see posology and clinical trial properties).

Psoriatic Arthritis (PsA)

STELARA alone or in combination with MTX, is indicated for the treatment of moderate to severe active psoriatic arthritis in adult patients when the response to previous nonbiological disease modifying anti rheumatic drug (DMARD) therapy has been inadequate,

- Reducing signs and symptoms
- Improving physical function
- Inhibiting the progression of structural damage
- Improving enthesitis
- Improving psoriasis
- Improving health-related QOL in adults with active psoriatic arthritis.

Crohn's Disease

Stelara is indicated for the treatment of adult patients with moderately to severely active Crohn's disease who have had an inadequate response with, lost response to, or were intolerant to either conventional therapy or a TNFα antagonist or have medical contraindications to such therapies.

Ulcerative Colitis

STELARA is indicated for the treatment of adult patients with moderately to severely active ulcerative colitis who have had an inadequate response with, lost response to, or were intolerant to either conventional therapy or a biologic or have medical contraindications to such therapies (see section 5.1).

Stelara 130 mg / 26 mL in Single-use Vial:

Crohn's Disease

Stelara is indicated for the treatment of adult patients with moderately to severely active Crohn's disease who have had an inadequate response with, lost response to, or were intolerant to either conventional therapy or a TNF α antagonist or have medical contraindications to such therapies.

Ulcerative Colitis

STELARA is indicated for the treatment of adult patients with moderately to severely active ulcerative colitis who have had an inadequate response with, lost response to, or were intolerant to either conventional therapy or a biologic or have medical contraindications to such therapies (see section 5.1).

DOSAGE AND ADMINISTRATION

Dosage – (Adults)

Plaque Psoriasis

For the treatment of plaque psoriasis, STELARA should be administered by subcutaneous injection. The recommended dose of STELARA is 45 mg administered at Weeks 0 and 4, then every 12 weeks thereafter. Alternatively, 90 mg may be used in patients with a body weight greater than 100 kg.

Dose Adjustment

For patients who inadequately respond to 45 mg every 12 weeks, consideration may be given to treating with 90 mg every 12 weeks. For patients who inadequately respond to dosing every 12 weeks, a 90 mg dose every 8 weeks may be considered.

Re-treatment

Re-treatment with a dosing regimen of Weeks 0 and 4 after interruption of therapy has been shown to be safe and effective.

Psoriatic Arthritis

For the treatment of psoriatic arthritis, STELARA is administered by subcutaneous injection.

The recommended dose of STELARA is 45 mg administered at Weeks 0 and 4, then every 12 weeks thereafter. Alternatively, 90 mg may be used in patients with a body weight greater than 100 kg.

Crohn's Disease and Ulcerative Colitis

In patients with Crohn's disease and ulcerative colitis, the recommended treatment regimen is a single intravenous (IV) tiered dose of STELARA based on body weight (refer to STELARA concentrate for solution for infusion), followed by 90 mg subcutaneous dosing 8 weeks later, then every 8 weeks thereafter (see Instructions for Use, Handling and Disposal).

| Table 1: Initial IV dosing of STELARA^a | | |
|--|-------------|---------------------------------------|
| Body Weight of Patient at the time of dosing | Dose | Number of 130 mg STELARA Vials |
| ≤ 55 kg | 260 mg | 2 |
| > 55 kg to ≤ 85 kg | 390 mg | 3 |
| > 85 kg | 520 mg | 4 |

^a approximately 6mg/kg

For some patients, a single IV dose based on body weight (Table 1) (refer to STELARA concentrate for solution for infusion) followed by 90 mg subcutaneous dosing 8 weeks later, then every 12 weeks thereafter may be acceptable. Patients who inadequately respond to 90 mg subcutaneous dosing every 12 weeks may benefit from an increase in dosing frequency to every 8 weeks (refer to STELARA solution for injection) (see Clinical Studies).

Immunomodulators and/or corticosteroids may be continued during treatment with STELARA. In patients who have responded to treatment with STELARA corticosteroids may be reduced or discontinued in accordance with standard of care.

If therapy in Crohn's disease or Ulcerative Colitis is interrupted, resumption of treatment with subcutaneous dosing every 8 weeks is safe and effective.

Dosage – (Pediatric population, 6 years and older)

Plaque Psoriasis

For the treatment of plaque psoriasis, STELARA should be administered by subcutaneous injection. The recommended dose of STELARA based on body weight is shown below (Table 2). STELARA should be administered at Weeks 0 and 4, then every 12 weeks thereafter.

Table 2: Recommended dose of STELARA for pediatric psoriasis

| Weight | Recommended Dose | Dosage Form |
|------------------|------------------|--------------------------|
| < 60 kg | 0.75 mg/kg* | Vial |
| ≥ 60 to ≤ 100 kg | 45 mg | Pre-filled syringe, vial |
| > 100 kg | 90 mg | Pre-filled syringe |

* To calculate the volume of injection (mL) for patients < 60 kg, use the following formula: $body\ weight\ (kg) \times 0.0083\ (mL/kg)$. The calculated volume should be rounded to the nearest 0.01 mL and administered using a 1 mL graduated syringe. A 45 mg vial is available for pediatric patients who need to receive less than the full 45 mg dose.

General Consideration for Administration

Subcutaneous administration

STELARA is intended for use under the guidance and supervision of a physician. Patients or their caregivers may inject STELARA if a physician determines that it is appropriate and with medical follow-up as necessary, after proper training in subcutaneous injection technique and disposal (see *Instructions for Use, Handling and Disposal*).

In pediatric patients, it is recommended that STELARA be administered by a healthcare provider.

Comprehensive instructions for the subcutaneous administration of STELARA are given in the "Core Patient Package Insert (CPPI)". Patients should be instructed to inject the prescribed amount of STELARA subcutaneously according to the directions provided in the patient information leaflet. The needle cover on the pre-filled syringe contains dry natural rubber (a derivative of latex), which may cause allergic reactions in individuals sensitive to latex. Each pre-filled syringe is for single dose only.

Intravenous infusion (Crohn's Disease and Ulcerative Colitis)

STELARA 130 mg vial is for IV infusion only. Intravenous infusion of STELARA should be administered by qualified health care professionals (For preparation, see *Instructions for Use, Handling and Disposal*).

Special populations

Pediatrics

Studies of STELARA in pediatric patients with psoriasis below 6 years of age have not been conducted. No studies have been conducted in pediatric patients with psoriatic arthritis, Crohn's disease or ulcerative colitis.

Elderly

Of the 6710 patients exposed to STELARA, a total of 353 were 65 years or older (183 patients with psoriasis, 69 patients with psoriatic arthritis, 58 with Crohn's disease, and 43 patients with ulcerative colitis). No major age-related differences in clearance or volume of distribution were observed in clinical studies. Although no overall differences in safety or efficacy were observed between older and younger patients in clinical studies in approved indications, the number of patients aged 65 and over is not sufficient to determine whether they respond differently from younger patients.

Renal impairment

Specific studies have not been conducted in patients with renal insufficiency.

Hepatic impairment

Specific studies have not been conducted in patients with hepatic insufficiency.

CONTRAINDICATIONS

Hypersensitivity to the active substance or to any of the excipients (See section List of excipients).

Clinically important, active infection (e.g. active tuberculosis).

WARNINGS AND PRECAUTIONS

Infections

STELARA is a selective immunosuppressant and may have the potential to increase the risk of infections and reactivate latent infections.

In clinical studies, serious bacterial, fungal, and viral infections have been observed in patients receiving STELARA.

STELARA should not be given to patients with a clinically important, active infection. Caution should be exercised when considering the use of STELARA in patients with a chronic infection or a history of recurrent infection.

Prior to initiating treatment with STELARA, patients should be evaluated for tuberculosis infection. STELARA should not be given to patients with active tuberculosis. Treatment of latent tuberculosis infection should be initiated prior to administering STELARA. Anti-tuberculosis therapy should also be considered prior to initiation of STELARA in patients with a past history of latent or active tuberculosis in whom an adequate course of treatment cannot be confirmed. Patients receiving STELARA should be monitored closely for signs and symptoms of active tuberculosis during and after treatment.

Patients should be instructed to seek medical advice if signs or symptoms suggestive of an infection occur. If a patient develops a serious infection they should be closely monitored and STELARA should not be administered until the infection resolves (see *Adverse Reactions*).

Malignancies

STELARA is a selective immunosuppressant. Immunosuppressive agents have the potential to increase the risk of malignancy. Some patients who received STELARA in clinical studies developed cutaneous and noncutaneous malignancies (see *Adverse Reactions*).

STELARA has not been studied in patients with a history of malignancy. Caution should be exercised when considering the use of STELARA in patients with a history of malignancy or when considering continuing treatment in patients who develop a malignancy.

All patients, in particular those greater than 60 years of age, patients with a medical history of prolonged immunosuppressant therapy or those with a history of PUVA treatment, should be monitored for the appearance of non-melanoma skin cancer (see *Adverse Reactions*).

Hypersensitivity reactions

In post-marketing experience, serious hypersensitivity reactions, including anaphylaxis and angioedema, have been reported. If an anaphylactic or other serious hypersensitivity reaction occurs, institute appropriate therapy and administration of STELARA should be discontinued (see *Adverse Reactions*).

Immunizations

It is recommended that live viral or live bacterial vaccines not be given concurrently with STELARA.

No data are available on the secondary transmission of infection by live vaccines in patients receiving STELARA. Caution is advised when administering some live vaccines to household contacts of patients receiving STELARA because of the potential risk for shedding from the household contact and transmission to the patient.

Patients receiving STELARA may receive concurrent inactivated or non-live vaccinations.

Long term treatment with STELARA does not suppress the humoral immune response to pneumococcal polysaccharide or tetanus vaccines (see *Pharmacodynamic Properties*).

Immunosuppression

In psoriasis studies, the safety and efficacy of STELARA in combination with immunosuppressive agents or phototherapy have not been evaluated. In psoriatic arthritis studies, concomitant MTX use did not appear to influence the safety or efficacy of STELARA. In Crohn's disease and ulcerative colitis studies, concomitant use of immunomodulators (6-mercaptopurine (6-MP), azathioprine (AZA), MTX) or corticosteroids did not appear to influence the safety or efficacy of STELARA. Caution should be exercised when considering concomitant use of immunosuppressive agents and STELARA or when transitioning from other biologic agents.

Immunotherapy

STELARA has not been evaluated in patients who have undergone allergy immunotherapy.

STELARA may affect allergy immunotherapy. Caution should be exercised in patients receiving or who have received allergy immunotherapy particularly for anaphylaxis.

General

The needle cover on the pre-filled syringe contains dry natural rubber (a derivative of latex), which may cause allergic reactions in individuals sensitive to latex.

INTERACTIONS

- The effects of IL-12 or IL-23 on the regulation of CYP450 enzymes were evaluated in an *in vitro* study using human hepatocytes, which showed that IL-12 and/or IL-23 at levels of 10 ng/mL did not alter human CYP450 enzyme activities (CYP1A2, 2B6, 2C9, 2C19, 2D6, or 3A4). Results from a Phase 1 study in subjects with active Crohn's disease suggest no clinically relevant drug interactions are likely. These results do not suggest the need for dose adjustments in patients who are receiving concomitant CYP450 substrates (see *Pharmacokinetic Properties*).
- Live vaccines should not be given concurrently with STELARA (see *Warnings & Precautions*).

Pregnancy, Breast-feeding and Fertility

Pregnancy

There is no evidence from animal studies of teratogenicity, birth defects or developmental delays at exposures up to approximately 150-fold higher compared to C_{max} following 4 weekly 90 mg subcutaneous injections or up to 21-fold higher compared to serum concentrations 1h following 6 mg/kg IV administration (See *Non-Clinical Information*). However, animal reproductive and developmental studies are not always predictive of human response.

Data from prospectively collected pregnancies following exposure to STELARA resulting in live birth with known outcomes, including more than 450 pregnancies exposed during the first trimester, do not indicate an increased risk of malformations in the newborn.

Overall, data from observational studies, pharmacovigilance, and published case reports and cohort studies do not indicate an increase in the risk of major birth defects, pattern of major or minor anomalies, miscarriage, or adverse infant outcomes.

STELARA should be given to a pregnant woman if the benefit clearly outweighs the risk.

Breast-feeding

STELARA is excreted in the milk of lactating monkeys administered STELARA. It is not known if STELARA is absorbed systemically after ingestion. Because many drugs and immunoglobulins are excreted in human milk, and because of the potential for adverse reactions in nursing infants from STELARA, a decision should be made whether to discontinue nursing or to discontinue the drug.

Fertility

The effect of STELARA on human fertility has not been evaluated. No adverse effects on female fertility parameters were identified in a female fertility toxicity study conducted in mice (see *Non-Clinical Information*).

It is not known whether STELARA can affect reproductive potential.

Effects on Ability to Drive and Use Machines

No studies on the effects on the ability to drive and use machines have been performed.

ADVERSE REACTIONS

Throughout this section, adverse reactions are presented. Adverse reactions are adverse events that were considered to be reasonably associated with the use of ustekinumab based on the comprehensive assessment of the available adverse event information. A causal relationship with ustekinumab cannot be reliably established in individual cases. Further, because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Clinical Studies Experience in Adult Patients with Psoriasis, Psoriatic Arthritis, Crohn's Disease, and Ulcerative Colitis

The safety data described below reflect exposure to STELARA in 14 Phase 2 and Phase 3 studies in 6710 patients (4135 with psoriasis and/or psoriatic arthritis, 1749 for Crohn's disease, and 826 with ulcerative colitis in UC-1 and UC-2 clinical trials), with duration of exposure to STELARA presented in Table 3.

Table 3: Long term exposure to STELARA in Phase 2 and Phase 3 clinical studies

| Exposure | Number of patients |
|-----------|--------------------|
| 6 months | 4577 ^a |
| 1 year | 3648 ^a |
| ≥ 4 years | 2194 ^b |
| ≥ 5 years | 1148 ^b |

^a Total number of patients in the psoriasis, psoriatic arthritis, Crohn's disease and ulcerative colitis studies

^b Number of patients with psoriasis, psoriatic arthritis, and Crohn's disease

The most common adverse reactions (>5%) in controlled periods of the clinical studies with STELARA among all indications were nasopharyngitis and headache. Most were considered to be mild and did not necessitate drug discontinuation. The overall safety profile of STELARA was similar for patients among all indications.

Table 4 provides a summary of Adverse Drug Reactions from the clinical studies. The frequency of these adverse reactions was based on those that occurred during the initial controlled periods of the clinical studies. The adverse drug reactions are ranked by frequency, using the following convention:

Very common (≥1/10)

Common (frequent) (≥1/100, <1/10)

Uncommon (infrequent) (≥1/1,000, <1/100)

Rare (≥1/10,000, <1/1,000)

Table 4: SUMMARY OF ADRs IN PSORIASIS CLINICAL STUDIES

| | |
|--|--|
| Infections and infestations | Common: Upper respiratory tract infection, nasopharyngitis, sinusitis Uncommon: Cellulitis, dental infections, herpes zoster, viral upper respiratory tract infection, vulvovaginal mycotic infection |
| Psychiatric disorders | Uncommon: Depression |
| Nervous system disorders | Common: Dizziness, headache |
| Respiratory, thoracic and mediastinal disorders | Common: Oropharyngeal pain Uncommon: Nasal congestion |
| Gastrointestinal disorders | Common: Diarrhea, nausea, vomiting |
| Skin and subcutaneous tissue disorders | Common: Pruritus Uncommon: Acne |
| Musculoskeletal and connective tissue disorders | Common: Back pain, myalgia, arthralgia |
| General disorders and administration site conditions | Common: Fatigue, injection site erythema, injection site pain Uncommon: Injection site reactions (including hemorrhage, hematoma, induration, swelling and pruritus), asthenia |

Infections

In the placebo-controlled studies of patients with psoriasis, psoriatic arthritis, Crohn's disease, and ulcerative colitis, the rates of infection or serious infection were similar between STELARA-treated patients and those treated with placebo. In the placebo-controlled period of the clinical studies of patients with psoriasis, patients with psoriatic arthritis, patients with Crohn's disease, and patients with ulcerative colitis, the rate of infection was 1.36 and 1.34 per patient-year of follow-up in STELARA and placebo-treated patients, respectively. Serious infections occurred at a rate of 0.03 per patient-year of follow-up in STELARA-treated patients (30 serious infections in 930 patient-years of follow-up) and 0.03 per patient-year of follow-up in placebo-treated patients (15 serious infections in 434 patient-years of follow-up).

In the controlled and non-controlled periods of psoriasis, psoriatic arthritis, Crohn's disease, and ulcerative colitis clinical studies representing 15227 patient-years of STELARA exposure in 6710 patients, the median follow-up was 1.2 years; 1.7 years for psoriatic disease studies, 0.6 year for Crohn's disease studies, and 2.3 years for ulcerative colitis studies. The rate of infections was 0.85 per patient-year of follow-up in STELARA-treated patients. The rate of serious infections was 0.02 per patient-year of follow-up in STELARA-treated patients (289 serious infections in 15227 patient-years of follow-up) and included pneumonia, anal abscess, cellulitis, diverticulitis, gastroenteritis and viral infections.

In clinical studies, patients with latent tuberculosis who were concurrently treated with isoniazid did not develop tuberculosis.

Malignancy

In the placebo-controlled period of the psoriasis, psoriatic arthritis, Crohn's disease, and ulcerative colitis clinical studies, the incidence of malignancies excluding non-melanoma skin cancer was 0.11 per 100 patient-years of follow-up for STELARA-treated

patients (1 patient in 929 patient-years of follow-up) and 0.23 per 100 patient-years of follow-up for placebo-treated patients (1 patient in 434 patient-years of follow-up). The incidence of non-melanoma skin cancer was 0.43 per 100 patient-years of follow-up for STELARA-treated patients (4 patients in 929 patient-years of follow-up) compared with 0.46 per 100 patient-years of follow-up for placebo-treated patients (2 patients in 433 patient-years of follow-up).

In the controlled and non-controlled periods of the psoriasis, psoriatic arthritis, Crohn's disease, and ulcerative colitis clinical studies, representing 15205 patient-years of STELARA exposure in 6710 patients, the median follow-up was 1.2 years; 1.7 years for psoriasis studies, 0.6 year for Crohn's disease studies, and 2.3 years for ulcerative colitis studies. Malignancies, excluding non-melanoma skin cancers, were reported in 76 patients in 15205 patient-years of follow-up (with an incidence of 0.50 per 100 patient-years of follow-up for STELARA-treated patients). The incidence of non-melanoma skin cancer was 0.46 per 100 patient-years of follow-up for STELARA-treated patients. The incidence of malignancies, reported in STELARA-treated patients was comparable to the incidence expected in the general population (standardized incidence ratio = 0.94 [95% confidence interval: 0.73, 1.18], adjusted for age, gender and race). The most frequently observed malignancies, other than non-melanoma skin cancer, were prostate, melanoma, colorectal and breast. The ratio of patients with basal versus squamous cell skin cancers (3:1) is comparable with the ratio expected in the general population (see *Warnings and Precautions*).

Hypersensitivity and Infusion Reactions

Subcutaneous administration

During the controlled periods of the psoriasis and psoriatic arthritis clinical studies of STELARA, rash and urticaria have each been observed in <1% of patients.

IV administration

In Crohn's disease and ulcerative colitis intravenous induction studies, no events of anaphylaxis or other serious infusion reactions were reported. In these studies, 2.2% of 785 placebo treated patients and 1.9% of 790 patients treated with the recommended dose of STELARA reported adverse events occurring during or within an hour of the infusion.

Immunogenicity

In psoriasis and psoriatic arthritis clinical studies, up to 12.4% of patients treated with STELARA developed antibodies to ustekinumab. Patients positive for antibodies to ustekinumab tended to have lower efficacy, however, antibody positivity did not preclude a clinical response. The majority of patients who were positive for antibodies to ustekinumab had neutralizing antibodies. In Crohn's disease and ulcerative colitis clinical studies, 2.9 and 4.6% of patients, respectively, developed antibodies to ustekinumab when treated with ustekinumab for approximately one year. No apparent association between the development of antibodies to ustekinumab and the development of injection site reactions was observed.

Overdose

Single doses up to 6 mg/kg intravenously have been administered in clinical studies without dose-limiting toxicity. In case of overdosage, it is recommended that the patient be monitored for any signs or symptoms of adverse reactions or effects and appropriate symptomatic treatment be instituted immediately.

Clinical Studies Experience in pediatric patients with Psoriasis

The safety of STELARA has been studied in two phase 3 studies of pediatric patients with moderate to severe plaque psoriasis. The first study was in 110 patients from 12 to 17 years of age treated for up to 60 weeks (CADMUS) and the second study was in 44 patients from 6 to 11 years of age treated for up to 56 weeks (CADMUS Jr.). In general, the adverse events reported in these two studies were similar to those seen in previous studies in adults with plaque psoriasis. (see *Clinical Studies Experience in Adult Patients with Psoriasis and/or Psoriatic Arthritis section above*).

Post Marketing Experience

The adverse reactions in Table 5 are ranked by frequency* using the following convention:

Very common: $\geq 1/10$

Common: $\geq 1/100$ and $<1/10$

Uncommon: $\geq 1/1,000$ and $<1/100$

Rare: $\geq 1/10,000$ and $<1/1000$

Very rare: $<1/10,000$, including isolated reports

Table 5: Post-Marketing Reports

| | |
|-------------------------|--|
| Immune system disorders | Uncommon: Hypersensitivity reactions (including rash, urticaria) |
|-------------------------|--|

| | |
|---|--|
| | Rare: Serious hypersensitivity reactions (including anaphylaxis and angioedema) |
| Infections and infestations | Uncommon: Lower respiratory tract infection |
| Respiratory, thoracic and mediastinal disorders | Rare: Allergic alveolitis, eosinophilic pneumonia |
| Skin and subcutaneous tissue disorders | Uncommon: Pustular psoriasis Rare: Erythrodermic psoriasis, hypersensitivity vasculitis |

*Post-marketing adverse reaction frequency is derived from the placebo-controlled portion of the 11 clinical trials if the adverse reaction was observed in those trials. Otherwise, it is estimated to be lower than a certain frequency given the exposure in the 11 clinical trials where the adverse reaction was not observed.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic Properties

Pharmacotherapeutic group: Immunosuppressants, interleukin inhibitors, ATC code: [L04AC05](#).

Mechanism of action

STELARA is a fully human IgG1κ monoclonal antibody that binds with specificity to the shared p40 protein subunit of human cytokines interleukin (IL)-12 and IL-23. STELARA inhibits the bioactivity of human IL-12 and IL-23 by preventing p40 from binding to the IL-12R β 1 receptor protein expressed on the surface of immune cells. STELARA cannot bind to IL-12 or IL-23 that is already bound to IL-12R β 1 cell surface receptors. Thus, STELARA is not likely to contribute to complement or antibody mediated cytotoxicity of cells expressing IL-12 and/or IL-23 receptors.

IL-12 and IL-23 are heterodimeric cytokines secreted by activated antigen presenting cells, such as macrophages and dendritic cells. IL-12 stimulates natural killer (NK) cells and drives the differentiation of CD4+ T cells toward the T helper 1 (Th1) phenotype and stimulates interferon gamma (IFN γ) production. IL-23 induces the T helper 17 (Th17) pathway and promotes secretion of IL-17A, IL-21, and IL-22. Levels of IL-12 and IL-23 are elevated in the skin and blood of patients with psoriasis and serum IL 12/23p40 distinguishes patients with psoriatic arthritis from healthy individuals, implicating IL-12 and IL-23 in the pathophysiology of psoriatic inflammatory disease. Genetic polymorphisms in IL23A, IL23R, and IL12B genes confer susceptibility to these disorders. Additionally, IL-12 and IL-23 are highly expressed in lesional psoriatic skin, and IL-12-mediated induction of IFN γ correlates with psoriasis disease activity. IL-23 responsive T-cells have been found in the enthuses in a mouse model of inflammatory arthritis, where IL-23 drives enthesal inflammation. In addition, there is pre-clinical evidence implicating IL-23 and downstream pathways in bone erosion and destruction through up-regulation of receptor activator of nuclear factor- κ B ligand (RANKL), which activates osteoclasts.

In patients with Crohn's disease, IL-12 and IL-23 are elevated in the intestines and lymph nodes. This is accompanied by increases in serum IFN γ and IL-17A levels, suggesting that IL-12 and IL-23 promote Th1 and Th17 activation in Crohn's disease. Both IL-12 and IL-23 can also stimulate TNF α production by T cells, resulting in chronic intestinal inflammation and epithelial cell injury. Significant associations have been found between Crohn's disease and genetic polymorphisms in the IL23R and IL12B genes, suggesting a potential causal role for IL-12/23 signaling in the disease. This is supported by pre-clinical data demonstrating that IL-12/23 signaling is required for intestinal injury in mouse models of inflammatory bowel disease.

By binding the shared p40 subunit of IL-12 and IL-23, STELARA may exert its clinical effects in psoriasis, psoriatic arthritis, Crohn's disease, and ulcerative colitis through interruption of the Th1 and Th17 cytokine pathways, which are central to the pathology of these diseases.

Pharmacodynamic effects

Treatment with STELARA resulted in significant improvement in histological measures of psoriasis including epidermal hyperplasia and cell proliferation. These results are consistent with the clinical efficacy observed.

In patients with psoriasis and/or psoriatic arthritis, STELARA had no apparent effect on the percentages of circulating immune cell populations including memory and naive T cell subsets or circulating cytokine levels. Systemic markers of inflammation were measurable in the serum at baseline and 4 markers (MDC, VEGF, MCSF-1 and YKL-40) showed modest differences in concentration post-treatment in STELARA-treated patients as compared to placebo.

Treatment with STELARA resulted in a decrease in the gene expression of its molecular targets IL-12 and IL-23 as shown by analyses of mRNA obtained from lesional skin biopsies of psoriatic patients at baseline and up to 2 weeks post-treatment. In addition, STELARA down regulated the gene expression of inflammatory cytokines and chemokines such as MCP-1, TNF-alpha, IP-10, and IL-8 in lesional skin biopsies. These results are consistent with the significant clinical benefit observed with STELARA treatment in psoriasis.

In psoriasis and psoriatic arthritis studies, clinical response (improvement in PASI or ACR measurements, respectively) appeared to be related to serum ustekinumab levels. Patients with psoriasis with higher PASI response had higher median serum concentrations of ustekinumab than those with lower clinical responses. In psoriasis studies, the proportion of patients who achieved PASI 75 response increased with increasing serum levels of ustekinumab. The proportion of patients who achieved PASI 75 response at Week 28 increased with increasing serum ustekinumab trough levels at Week 28. In psoriatic arthritis studies, patients achieving an ACR 20 response had higher median serum concentrations of ustekinumab than ACR 20 non-responders. The proportion of patients who achieved ACR 20 and ACR 50 response increased with increasing serum levels of ustekinumab.

In patients with Crohn's disease, treatment with STELARA resulted in a significant decrease in inflammatory markers including C-Reactive Protein (CRP) and fecal calprotectin. CRP was assessed during the study extension and the reductions observed during maintenance were generally sustained through week 252. Reductions in serum IFN γ and IL-17A, which are IL-12 and IL-23 regulated pro-inflammatory cytokines, were achieved and maintained in STELARA treated patients through Week 44 compared to placebo. Expression of genes such as IL-12R β 1 and IL-23 was reduced in inflamed colon tissue from Crohn's disease patients, responders to STELARA treatment while no significant changes were observed in placebo treated patients at Week 6.

In patients with ulcerative colitis, treatment with STELARA resulted in a decrease in inflammatory markers including CRP and fecal calprotectin during the induction phase, which were maintained throughout the maintenance phase and study extension through week 200.

Immunization

During the long term extension of a Phase 3 psoriasis study (PHOENIX 2), patients treated with STELARA for at least 3.5 years mounted similar antibody responses to both pneumococcal polysaccharide and tetanus vaccines as a non-systemically treated psoriasis control group. Similar proportions of patients developed protective levels of anti-pneumococcal and anti-tetanus antibodies and antibody titers were similar among STELARA-treated and control patients.

Clinical studies

Clinical Efficacy - Plaque Psoriasis (Adults)

The safety and efficacy of STELARA was assessed in 2 Phase 3, multicenter, randomized, double-blind, placebo-controlled studies in patients with moderate to severe plaque psoriasis (PHOENIX 1 and PHOENIX 2). A total of 1996 patients were enrolled in these studies.

The studies enrolled adults (\geq 18 years) with chronic ($>$ 6 months) plaque psoriasis who had a minimum body surface area (BSA) involvement of 10%, and PASI score \geq 12 and who were candidates for systemic therapy or phototherapy. Patients with guttate, erythrodermic, or pustular psoriasis were excluded from the studies. No concomitant antipsoriatic therapies were allowed during the study with the exception of low-potency topical corticosteroids on the face and groin after week 12.

The PASI is a composite score that assesses the fraction of body surface area involved with psoriasis and the severity of psoriatic changes within the affected regions (plaque thickness/induration, erythema, and scaling). PASI numeric scores range from 0 to 72, with higher scores representing more severe disease.

Patients achieving \geq 75% improvement in PASI from baseline (PASI 75) were considered PASI 75 responders. Patients originally randomized to STELARA who were PASI 75 responders at both Weeks 28 and 40 were considered long-term PASI 75 responders. Patients achieving \geq 90% improvement in PASI from baseline (PASI 90) were considered PASI 90 responders and patients with \geq 50% improvement in PASI from baseline (PASI 50) were considered PASI 50 responders. Patients who achieved \geq 50% but less than 75% improvement in PASI from baseline were considered partial responders. Patients with $<$ 50% improvement in PASI from baseline were considered nonresponders.

Other key efficacy assessments included:

- The Physician's Global Assessment (PGA), a 6-category scale: 0 = cleared, 1 = minimal, 2 = mild, 3 = moderate, 4 = marked and 5 = severe, that indicates the physician's overall assessment of psoriasis focusing on plaque thickness/induration, erythema, and scaling. The PGA was assessed in PHOENIX 1 and 2.
- The Dermatology Life Quality Index (DLQI), a dermatology-specific quality of life instrument designed to assess the impact of the disease on a patient's quality of life. DLQI scores range from 0 to 30, with a lower score representing a better quality of life. A decrease of 5 in the DLQI score from baseline is considered a clinically meaningful improvement. The DLQI was assessed in PHOENIX 1 and 2.

- The SF-36, a health survey questionnaire consisting of multi-item scales measuring 8 health concepts. The SF-36 yields composite scores that provide a measure of disease impact on physical and mental health status. Higher SF-36 scores indicate a better quality of life. The SF-36 was assessed in PHOENIX 1.
- The Nail Psoriasis Severity Index (NAPSI), a physician-assessed score that measures the severity of nail involvement. The scale consists of 4 components of nail matrix disease and 4 components of nail bed disease with scores from 0 to 8, with a lower scores representing milder disease. The NAPSI was assessed in PHOENIX 1.
- The Hospital Anxiety and Depression Scale (HADS), a self-rating tool developed to evaluate psychological measures in patients with physical ailments. It consists of 2 subscales, one measuring anxiety (A-scale) and one measuring Depression (D-scale), which are scored separately. Lower HADS scores correspond to lesser psychological impairment. The HADS was assessed in PHOENIX 2.
- The Work Limitations Questionnaire (WLQ), a 25-item, self-administered questionnaire that was used to measure the impact of chronic health conditions on job performance and work productivity among employed populations. The WLQ assesses four aspects of work and productivity: Physical Demands, Time Management, Mental-Interpersonal Demand, and Output Demand. The four subscales range from 0-100 with the lower score indicating fewer work limitations. The WLQ was assessed in PHOENIX 2.
- The Itch Visual Analog Scale, used to assess the severity of itch at the time of the assessment. Itch is assessed using a 10 cm horizontal line, or a Visual Analog Scale (VAS), representing the range of itch severity, from 0 (no itch at all) to 10 (severe itch). The Itch VAS was assessed in PHOENIX 1.

PHOENIX 1

PHOENIX 1 evaluated the safety and efficacy of STELARA versus placebo in 766 patients with plaque psoriasis and the efficacy of every 12 week dosing for patients who were PASI 75 responders.

Patients randomized to STELARA received 45 mg or 90 mg doses at Weeks 0 and 4 followed by the same doses every 12 weeks. Patients randomized to receive placebo at Weeks 0 and 4 crossed over to receive STELARA (either 45 mg or 90 mg) at Weeks 12 and 16 followed by the same dose every 12 weeks.

Maintenance dosing (every 12 weeks)

To evaluate the therapeutic benefit of maintenance dosing with STELARA, patients originally randomized to STELARA who were PASI 75 responders at both Weeks 28 and 40 were re-randomized to either maintenance dosing of STELARA every 12 weeks or to placebo (i.e., withdrawal of therapy). Patients who were re-randomized to placebo at Week 40 reinitiated STELARA at their original dosing regimen when they experienced at least a 50% loss of their PASI improvement obtained at Week 40.

Dose Adjustment (every 8 weeks)

At Week 28, patients who were nonresponders discontinued treatment and patients who were partial responders were adjusted to every-8-week dosing.

PASI 75 responders at week 28 who became partial responders or nonresponders at Week 40 were adjusted to every-8-week dosing.

All patients were followed for up to 76 weeks following first administration of study treatment.

PHOENIX 2

PHOENIX 2 evaluated the safety and efficacy of STELARA versus placebo in 1230 patients with plaque psoriasis. Patients randomized to STELARA received 45 mg or 90 mg doses at Weeks 0 and 4 followed by an additional dose at Week 16. Patients randomized to receive placebo at Weeks 0 and 4 crossed over to receive STELARA (either 45 mg or 90 mg) at Weeks 12 and 16 followed by the same dose every 12 weeks.

Dose Adjustment (every 8 weeks)

At Week 28, patients who were nonresponders discontinued treatment and patients who were partial responders were re-randomized to continue every-12-week dosing or switch to every-8-week dosing.

PASI 75 responders at week 28 who became partial responders or nonresponders at Week 40 were adjusted to every-8-week dosing.

All patients were followed for up to 52 weeks following first administration of study agent.

Baseline disease characteristics: PHOENIX 1 and 2

Baseline disease characteristics across PHOENIX 1 and 2 were similar (Table 6).

Table 6: Baseline Disease Characteristics

| | PHOENIX 1 | | PHOENIX 2 | |
|---|------------------|------------------|------------------|------------------|
| | Placebo N=255 | STELARA N=511 | Placebo N=410 | STELARA N=820 |
| Patients randomized at Week 0 | | | | |
| Median BSA | 22.0 | 21.0 | 20.0 | 21.0 |
| BSA ≥ 20% | 145 (57%) | 276 (54%) | 217 (53%) | 445 (54%) |
| Median PASI | 17.80 | 17.40 | 16.90 | 17.60 |
| PASI ≥ 20 | 91 (36%) | 169 (33%) | 133 (32%) | 300 (37%) |
| PGA of marked or severe | 112 (44%) | 223 (44%) | 160 (39%) | 328 (40%) |
| History of psoriatic arthritis | 90 (35%) | 168 (33%) | 105 (26%) | 200 (24%) |
| Prior phototherapy | 150 (59%) | 342 (67%) | 276 (67%) | 553 (67%) |
| Prior conventional systemic therapy excluding biologics | 142 (56%) | 282 (55%) | 241 (59%) | 447 (55%) |
| Prior conventional systemic or biologic therapy | 189 (74%) | 364 (71%) | 287 (70%) | 536 (65%) |
| Failed to respond to, had contraindication for, or intolerant to ≥ 1 conventional therapy | 139 (55%) | 270 (53%) | 254 (62%) | 490 (60%) |
| Failed to respond to, had contraindication for, or intolerant to ≥ 3 conventional therapies | 30 (12%) | 54 (11%) | 66 (16%) | 134 (16%) |

Efficacy at the Primary Endpoint, PHOENIX 1 and 2

In both the PHOENIX 1 and PHOENIX 2 studies, a significantly greater proportion of patients randomized to treatment with STELARA were PASI 75 responders compared with placebo at Week 12 (Table 7). In the PHOENIX 1 study, 67% and 66% of patients receiving STELARA 45 mg and 90 mg, respectively, achieved a PASI 75 response at Week 12 compared with 3% of patients receiving placebo. In the PHOENIX 2 study, 67% and 76% of patients receiving STELARA 45 mg and 90 mg respectively achieved a PASI 75 response at Week 12 compared with 4% of patients receiving placebo.

All 3 components of the PASI (plaque thickness/induration, erythema, and scaling) contributed comparably to the improvement in PASI.

The efficacy of STELARA was significantly superior ($p<0.001$) to placebo across all subgroups defined by baseline demographics, clinical disease characteristics (including patients with a history of psoriatic arthritis) and prior medication usage. While pharmacokinetic modeling suggested a trend towards higher CL/F in patients with diabetes, a consistent effect on efficacy was not observed.

Other efficacy measures at Week 12

In both PHOENIX 1 and PHOENIX 2, compared with placebo, significantly greater proportions of patients randomized to 45 mg or 90 mg STELARA achieved a cleared or minimal PGA score, and significantly greater proportions of patients randomized to 45 mg or 90 mg STELARA were PASI 90 and PASI 50 responders at Week 12 (Table 2). In the PHOENIX 1 study, 59% and 61% of the patients treated with 45 mg and 90 mg STELARA, respectively, achieved PGA scores of cleared or minimal compared with 4% of placebo-treated patients. In PHOENIX 2, 68% and 73 % of patients receiving 45 mg or 90 mg STELARA, respectively, had cleared or minimal PGA scores compared with 4% of the placebo patients. In PHOENIX 1, PASI 90 was achieved by 42% and 37% of the patients treated with 45 mg and 90 mg STELARA, respectively, compared with 2% of placebo-treated patients. In PHOENIX 2, the percentage of patients achieving PASI 90 was 42% in the 45 mg STELARA group, 51% in the 90 mg STELARA group and 1% in the placebo group. The percentage of patients achieving PASI 50 in PHOENIX 1 was 84% and 86% in the 45 mg and 90 mg STELARA groups, respectively, compared with 10% in the placebo group. Similarly, 84% of patients treated with 45 mg STELARA, 89% of patients treated with 90 mg STELARA and 10% of patients treated with placebo reached PASI 50 in PHOENIX 2 (Table 7).

Table 7: Key psoriasis endpoints – PHOENIX 1 and PHOENIX 2

| Week 12 | PHOENIX 1 | | | PHOENIX 2 | | |
|-------------------------------|-----------|-----------|-----------|-----------|-----------|-----------|
| | | STELARA | | | STELARA | |
| | | Placebo | 45 mg | 90 mg | Placebo | 45 mg |
| Patients randomized at Week 0 | 255 | 255 | 256 | 410 | 409 | 411 |
| PASI response | | | | | | |
| PASI 50 response ^a | 26 (10%) | 213 (84%) | 220 (86%) | 41 (10%) | 342 (84%) | 367 (89%) |
| PASI 75 response ^a | 8 (3%) | 171 (67%) | 170 (66%) | 15 (4%) | 273 (67%) | 311 (76%) |
| PASI 90 response ^a | 5 (2%) | 106 (42%) | 94 (37%) | 3 (1%) | 173 (42%) | 209 (51%) |

| | | | | | | |
|--|---------|------------------|--------------|---------|------------------|--------------|
| PGA of Cleared or Minimal ^{a,b} | 10 (4%) | 151 (59%) | 156 (61%) | 18 (4%) | 277 (68%) | 300 (73%) |
| PASI 75 response by weight | | | | | | |
| ≤ 100 kg | | | | | | |
| N | 166 | 168 | 164 | 290 | 297 | 289 |
| PASI 75 response | 6 (4%) | 124 (74%) | 107 (65%) | 12 (4%) | 218 (73%) | 225 (78%) |
| > 100 kg | | | | | | |
| N | 89 | 87 | 92 | 120 | 112 | 121 |
| PASI 75 response | 2 (2%) | 47 (54%) | 63 (68%) | 3 (3%) | 55 (49%) | 86 (71%) |
| PGA of Cleared or Minimal by weight | | | | | | |
| ≤ 100 kg | | | | | | |
| N | 166 | 168 | 164 | 290 | 297 | 289 |
| PGA response ^b | 7 (4%) | 108 (64%) | 103 (63%) | 14 (5%) | 220 (74%) | 216 (75%) |
| > 100 kg | | | | | | |
| N | 89 | 87 | 92 | 120 | 112 | 121 |
| PGA response ^b | 3 (3%) | 43 (49%) | 53 (58%) | 4 (3%) | 57 (51%) | 84 (69%) |
| Week 28 | | | | | | |
| | | PHOENIX 1 | | | PHOENIX 2 | |
| | | STELARA | | | STELARA | |
| | | <u>45 mg</u> | <u>90 mg</u> | | <u>45 mg</u> | <u>90 mg</u> |
| N | | 250 | 243 | | 397 | 400 |
| PASI response | | | | | | |
| PASI 50 response | | 228 (91%) | 234 (96%) | | 369 (93%) | 380 (95%) |
| PASI 75 response | | 178 (71%) | 191 (79%) | | 276 (70%) | 314 (79%) |
| PASI 90 response | | 123 (49%) | 135 (56%) | | 178 (45%) | 217 (54%) |
| PGA of Cleared or Minimal ^b | | 146 (58%) | 160 (66%) | | 241 (61%) | 279 (70%) |
| PASI 75 response by weight | | | | | | |
| ≤ 100 kg | | | | | | |
| N | | 164 | 153 | | 287 | 280 |
| PASI 75 response | | 130 (79%) | 124 (81%) | | 217 (76%) | 226 (81%) |
| > 100 kg | | | | | | |
| N | | 86 | 90 | | 110 | 119 |
| PASI 75 response | | 48 (56%) | 67 (74%) | | 59 (54%) | 88 (74%) |
| PGA of Cleared or Minimal by weight | | | | | | |
| ≤ 100 kg | | | | | | |
| N | | 164 | 153 | | 287 | 280 |
| PGA response ^b | | 106 (65%) | 106 (69%) | | 192 (67%) | 207 (74%) |
| > 100 kg | | | | | | |
| N | | 86 | 90 | | 110 | 119 |
| PGA response | | 40 (47%) | 54 (60%) | | 49 (45%) | 71 (60%) |

^a p < 0.001 for 45 mg or 90 mg comparison with placebo

^b data corrected post EMEA inspection


Response over time

In PHOENIX 1, significantly greater proportions of STELARA-treated patients had PASI 50 responses (9% and 10% for the 45 mg and 90 mg groups, respectively) compared with placebo (2%) by Week 2 (p< 0.001). Significantly greater proportions of patients treated with STELARA achieved PASI 75 responses (9% and 12% for the 45 mg and 90 mg STELARA groups, respectively) compared with placebo (0.4%) by Week 4 (p< 0.001). Maximum response was generally achieved by Week 24 in the 45 mg and 90 mg STELARA treatment groups, and response rates were generally sustained through Week 36 (Figure 1). In PHOENIX 1, PASI 75 rates at Week 24 were 76% for the 45 mg group, and 85% for the 90 mg group. Higher response rates were observed in patients

receiving STELARA 90 mg than in those receiving STELARA 45 mg by Week 16 and these higher response rates were sustained through Week 36 (Figure 1). Similar results were observed in the PHOENIX 2 study through Week 28.

In pre-specified analyses of efficacy by body weight in PHOENIX 1 and PHOENIX 2, no consistent pattern of dose response was seen in patients ≤ 100 kg. In patients who weighed >100 kg, higher PASI 75 response rates were seen with 90 mg dosing compared with 45 mg dosing, and a higher proportion of patients receiving 90 mg dosing had PGA scores of cleared or minimal compared with patients receiving 45 mg dosing (Table 6).

Figure 1: shows PASI 75 response over time in PHOENIX 1 and 2:



Therapeutic benefit of Long-term continuous use

At Week 40 in PHOENIX 1, 162 patients were randomized to receive STELARA (maintenance) and 160 were randomized to receive placebo (treatment withdrawal). Maintenance of PASI 75 was significantly superior with continuous treatment compared with treatment withdrawal ($p<0.001$). Similar results were seen with each dose of STELARA (Figure 2). At 1 year (Week 52), 89% of patients re-randomized to maintenance treatment were PASI 75 responders compared with 63% of patients re-randomized to placebo (treatment withdrawal) ($p<0.001$). At 18 months (Week 76), 84% of patients re-randomized to maintenance treatment were PASI 75 responders compared with 19% of patients re-randomized to placebo (treatment withdrawal). At 3 years (Week 148), 82% of patients re-randomized to maintenance treatment were PASI 75 responders. At 5 years (Week 244), 80% of patients re-randomized to maintenance treatment were PASI 75 responders.




Figure 2: Life-table estimate of percent of patients maintaining PASI 75 response; patients randomized at Week 40 (PHOENIX 1)

Efficacy of retreatment

In PHOENIX 1, after withdrawal from therapy, patients reinitiated their original STELARA treatment regimen after loss of $\geq 50\%$ of PASI improvement. Retreatment with STELARA resulted in 71% of evaluated patients regaining PASI 75 response within 8 weeks after reinitiating therapy and 85% of evaluated patients regaining PASI 75 response within 12 weeks after reinitiating therapy.

Dosing Interval Adjustment

In PHOENIX 1, Week 28 and Week 40 Partial Responders and Week 40 Nonresponders were adjusted from every 12 week to every 8 week dosing. Approximately 40%-50% of Week 28 Partial Responders to every 12 week dosing achieved PASI 75 response after adjustment to every 8 week dosing and this proportion of PASI 75 responders was maintained through Week 52. A similar proportion of patients who were PASI 75 responders at Week 28 and subsequently became partial responders or nonresponders at Week 40 achieved PASI 75 response following a dosing interval adjustment to every 8 weeks.

Quality of Life

In PHOENIX 1 and 2, the mean baseline DLQI scores ranged from 11 to 12. In PHOENIX 1, the mean baseline SF-36 Physical Component ranged from 47-49 and the mean baseline SF-36 Mental Component was approximately 50. Quality of life improved significantly in patients randomized to 45 mg or 90 mg STELARA compared with patients randomized to placebo as evaluated by DLQI in PHOENIX 1 and 2 and SF-36 in PHOENIX 1 (Tables 8 and 9). Quality of life improvements were significant as early as 2 weeks in patients treated with STELARA and these improvements were maintained over time with continued dosing.

Table 8: Quality of Life endpoints through Week 40 – PHOENIX 1

| | STELARA | | |
|-------------------------------|-----------------|-----------------|-----------------|
| | Placebo | 45 mg | 90 mg |
| Patients randomized at Week 0 | 255 | 255 | 256 |
| DLQI | | | |
| Baseline | | | |
| N | 254 | 255 | 255 |
| Mean \pm SD | 11.8 \pm 7.41 | 11.1 \pm 7.09 | 11.6 \pm 6.92 |
| Median | 10.0 | 10.0 | 11.0 |
| Change from baseline | | | |
| Week 2 ^a | | | |
| N | 253 | 255 | 254 |
| Mean \pm SD | -0.9 \pm 4.88 | -3.6 \pm 4.51 | -4.5 \pm 5.31 |
| Median | -1.0 | -3.0 | -4.0 |

| | | | |
|----------------------------|--------------------|--------------------|--------------------|
| Week 12 ^a | | | |
| N | 252 | 254 | 249 |
| Mean \pm SD | -0.6 \pm 5.97 | -8.0 \pm 6.87 | -8.7 \pm 6.47 |
| Median | 0.0 | -6.0 | -7.0 |
| Week 28 | | | |
| N | NA | 249 | 241 |
| Mean \pm SD | NA | -8.1 \pm 7.23 | -9.6 \pm 7.17 |
| Median | NA | -7.0 | -8.0 |
| Week 40 | | | |
| N | NA | 246 | 236 |
| Mean \pm SD | NA | -8.2 \pm 7.23 | -9.5 \pm 6.96 |
| Median | NA | -7.0 | -9.0 |
| SF-36 | | | |
| Physical component summary | | | |
| Baseline | | | |
| N | 254 | 255 | 255 |
| Mean \pm SD | 47.22 \pm 10.240 | 48.90 \pm 9.555 | 47.51 \pm 9.224 |
| Median | 50.70 | 51.60 | 49.60 |
| Change from Baseline | | | |
| Week 12 ^a | | | |
| N | 250 | 255 | 249 |
| Mean \pm SD | -0.53 \pm 7.457 | 1.97 \pm 7.422 | 3.23 \pm 7.590 |
| Median | -0.25 | 1.30 | 1.50 |
| Week 28 | | | |
| N | NA | 250 | 239 |
| Mean \pm SD | NA | 1.86 \pm 8.301 | 3.17 \pm 7.855 |
| Median | NA | 1.00 | 1.90 |
| Week 40 | | | |
| N | NA | 246 | 236 |
| Mean \pm SD | NA | 1.77 \pm 8.402 | 2.96 \pm 8.027 |
| Median | NA | 0.80 | 2.10 |
| Mental component summary | | | |
| Baseline | | | |
| N | 254 | 255 | 255 |
| Mean \pm SD | 49.62 \pm 10.582 | 50.02 \pm 10.425 | 49.86 \pm 10.175 |
| Median | 53.35 | 52.90 | 53.10 |
| Change from Baseline | | | |
| Week 12 ^a | | | |
| N | 250 | 255 | 249 |
| Mean \pm SD | -1.33 \pm 7.473 | 2.12 \pm 9.308 | 2.54 \pm 9.506 |
| Median | -0.60 | 0.80 | 1.50 |
| Week 28 | | | |
| N | NA | 250 | 239 |
| Mean \pm SD | NA | 1.80 \pm 9.578 | 3.47 \pm 9.587 |
| Median | NA | 0.40 | 1.50 |
| Week 40 | | | |
| N | NA | 246 | 236 |
| Mean \pm SD | NA | 2.17 \pm 9.137 | 2.91 \pm 9.418 |
| Median | NA | 0.95 | 1.10 |

^a p < 0.001 for 45 mg or 90 mg comparison with placebo.

NA = not applicable

Table 9: Quality of Life endpoints through Week 24 – PHOENIX 2

| | Placebo | STELARA | |
|-------------------------------|---------|--------------|--------------|
| | | <u>45 mg</u> | <u>90 mg</u> |
| Patients randomized at Week 0 | 410 | 409 | 411 |
| DLQI | | | |

| | | | |
|----------------------|-----------------|-----------------|------------------|
| Baseline | | | |
| N | 408 | 406 | 408 |
| Mean \pm SD | 12.3 \pm 6.86 | 12.2 \pm 7.07 | 12.6 \pm 7.29 |
| Median | 11.0 | 12.0 | 12.0 |
| Change from baseline | | | |
| Week 4 ^a | | | |
| N | 405 | 404 | 404 |
| Mean \pm SD | -1.4 \pm 4.68 | -6.9 \pm 6.07 | -7.0 \pm 5.86 |
| Median | -1.0 | -6.0 | -6.0 |
| Week 12 ^a | | | |
| N | 400 | 401 | 402 |
| Mean \pm SD | -0.5 \pm 5.66 | -9.3 \pm 7.12 | -10.0 \pm 6.67 |
| Median | -0.5 | -8.0 | -9.0 |
| Week 24 | | | |
| N | NA | 394 | 399 |
| Mean \pm SD | NA | -9.5 \pm 7.26 | -10.3 \pm 6.96 |
| Median | NA | -8.0 | -9.0 |

^a p < 0.001 for 45 mg or 90 mg comparison with placebo.

NA = not applicable

Nail Psoriasis

In PHOENIX 1, the median baseline NAPSI score for nail psoriasis was 4.0 and the median number of fingernails involved with psoriasis was 8.0. Nail psoriasis improved significantly in patients randomized to 45 mg or 90 mg STELARA compared with patients randomized to placebo when measured by the NAPSI score (Tables 10 and 11). Nail psoriasis continued to improve over time through Week 52 in patients treated with STELARA.

Table 10: Summary of percent improvement from baseline in NAPSI at Week 12; patients randomized at Week 0 with nail psoriasis present at Week 0 – PHOENIX 1

| | STELARA | | |
|--|------------------|------------------|------------------|
| | Placebo | 45 mg | 90 mg |
| Placebo randomized at Week 0 with nail psoriasis present at Week 0 | 176 | 182 | 187 |
| Week 12 ^a | | | |
| N | 174 | 182 | 184 |
| Mean \pm SD | 11.8 \pm 51.09 | 26.7 \pm 56.80 | 24.9 \pm 48.90 |
| Median | 0.0 | 25.0 | 25.0 |

^ap ≤ 0.001 for 45 mg or 90 mg comparison with placebo

Table 11: Summary of percent improvement from baseline in NAPSI at Week 24; patients randomized at Week 0 with nail psoriasis present at Week 0 – PHOENIX 1

| | STELARA | | | |
|---|-----------------------------|-----------------------------|-----------------|-----------------|
| | Placebo \rightarrow 45 mg | Placebo \rightarrow 90 mg | 45 mg | 90 mg |
| Patients randomized at Week 0 with nail psoriasis present at Week 0 | 93 | 83 | 182 | 187 |
| Week 24 | | | | |
| N | 89 | 77 | 179 | 181 |
| Mean \pm SD | 29.1 \pm 60.83 | 40.5 \pm 43.37 | 46.5 \pm 47.4 | 48.7 \pm 45.5 |
| Median | 33.3 | 42.9 | 50.0 | 50.0 |

Hospital Anxiety and Depression Scale

At baseline in PHOENIX 2, the mean HADS anxiety and depression scores were 6.9 and 5.1, respectively. Both anxiety and depression scores were reduced significantly in patients randomized to 45 mg or 90 mg STELARA at Week 12 compared with patients randomized to placebo (Table 12). HADS improvements were maintained through Week 24 (Table 13).

Table 12: Summary of change from baseline in Hospital Anxiety and Depression at Week 12; patients randomized at Week 0 – PHOENIX 2

| | STELARA | | |
|--|---------|-------|-------|
| | Placebo | 45 mg | 90 mg |

| | | | |
|-------------------------------|-------------|-------------|-------------|
| Patients randomized at Week 0 | 410 | 409 | 411 |
| Anxiety score ^a | | | |
| N | 395 | 399 | 399 |
| Mean ± SD | -0.11±2.689 | -1.59±3.570 | -1.60±3.351 |
| Median | 0.00 | -1.00 | -1.00 |
| Depression score ^a | | | |
| N | 398 | 399 | 401 |
| Mean ± SD | 0.21±2.757 | -1.71±3.124 | -2.06±3.420 |
| Median | 0.00 | -1.00 | -1.00 |

^ap ≤ 0.001 for 45 mg or 90 mg comparison with placebo.

Table 13: Summary of change from baseline in Hospital Anxiety and Depression at Week 24; patients randomized at Week 0 – PHOENIX 2

| | <u>STELARA</u> | | | |
|-------------------------------|------------------------|------------------------|--------------|--------------|
| | <u>Placebo → 45 mg</u> | <u>Placebo → 90 mg</u> | <u>45 mg</u> | <u>90 mg</u> |
| Patients randomized at Week 0 | 205 | 205 | 409 | 411 |
| Anxiety score ^a | | | | |
| N | 183 | 191 | 393 | 395 |
| Mean ± SD | -1.52±3.148 | -1.76±3.245 | -1.80±3.725 | -1.99±3.463 |
| Median | -1.00 | -1.00 | -1.00 | -1.00 |
| Depression score ^a | | | | |
| N | 184 | 190 | 391 | 398 |
| Mean ± SD | -1.65±3.207 | -1.42±3.013 | -1.77±3.449 | -2.26±3.490 |
| Median | -1.00 | -1.00 | -1.00 | -2.00 |

Work Limitations Questionnaire

The Work Limitations Questionnaire obtained at baseline showed impaired work productivity among patients with psoriasis evaluated in PHOENIX 2 for the Physical Demands, Time Management, Mental-Interpersonal and Output Demands component scores. Work productivity improved significantly more in patients randomized to STELARA at Week 12 compared with patients randomized to placebo as measured by the four WLQ subscales (Physical Demands, Time Management, Mental-Interpersonal, and Output Demands; Table 14).

Table 14: Summary of change from baseline in Work Limitations Questionnaire at Week 12; patients randomized at Week 0 – PHOENIX 2

| | <u>STELARA</u> | | |
|---|----------------|--------------|--------------|
| | <u>Placebo</u> | <u>45 mg</u> | <u>90 mg</u> |
| Patients randomized at Week 0 | 410 | 409 | 411 |
| Physical Demands score ^a | | | |
| N | 277 | 277 | 281 |
| Mean ± SD | -0.20±30.991 | -7.61±30.917 | -5.05±34.050 |
| Median | 0.00 | 0.00 | 0.00 |
| Time Management score ^b | | | |
| N | 259 | 255 | 265 |
| Mean ± SD | 0.74±18.962 | -6.58±21.634 | -9.06±24.239 |
| Median | 0.00 | -5.00 | -3.30 |
| Mental - Interpersonal score ^b | | | |
| N | 272 | 275 | 276 |
| Mean ± SD | 1.11±18.881 | -7.82±22.684 | -7.51±19.366 |
| Median | 0.00 | -2.80 | -1.35 |
| Output Demands score ^b | | | |
| N | 276 | 274 | 279 |
| Mean ± SD | 1.08±16.062 | -6.82±22.367 | -6.98±20.866 |
| Median | 0.00 | 0.00 | 0.00 |

^ap = 0.001 and 0.060 for the 45 mg and 90 mg comparisons, respectively, with placebo

^bp < 0.001 for 45 mg or 90 mg comparison with placebo

Itch VAS

Itch associated with psoriasis improved significantly ($p<0.001$) at Week 12 in patients randomized to 45 mg or 90 mg STELARA compared with patients randomized to placebo as evaluated by Itch VAS in PHOENIX 1 (Table 15).

Table 15: Summary of change from baseline in itch VAS at Week 12; patients randomized at Week 0 – PHOENIX 1

| | STELARA | | |
|-------------------------------|----------------|--------------|--------------|
| | <u>Placebo</u> | <u>45 mg</u> | <u>90 mg</u> |
| Patients randomized at Week 0 | 255 | 255 | 256 |
| Week 12 ^a | | | |
| N | 252 | 253 | 249 |
| Mean ± SD | -0.78±2.538 | -4.91±3.142 | -5.14±3.020 |
| Median | -0.30 | -5.50 | -5.50 |

^a $p < 0.001$ for 45 mg or 90 mg comparison with placebo

ACCEPT

In addition, a multicenter, randomized, single-blind, active-controlled study (ACCEPT) compared the safety and efficacy of ustekinumab and etanercept in patients 18 years of age and older with chronic (>6 months) plaque psoriasis who had a minimum BSA involvement of 10%, PASI score ≥ 12 , Physician Global Assessment (PGA) score ≥ 3 , who were candidates for phototherapy or systemic therapy, and who had had an inadequate response to, intolerance to, or contraindication to cyclosporine, MTX, or PUVA therapy. A total of 903 patients were enrolled in the study.

The ACCEPT trial compared the efficacy of ustekinumab to etanercept and evaluated the safety of ustekinumab and etanercept in patients with moderate to severe psoriasis. The active-controlled portion of the study was from Week 0 to Week 12, during which patients were randomized to receive etanercept (50mg twice a week) ustekinumab 45 mg at Weeks 0 and 4, or ustekinumab 90 mg at Weeks 0 and 4. This trial was powered to test the superiority of each ustekinumab dose to etanercept on the primary endpoint of the proportion of patients who achieved a PASI 75 at week 12.

Significantly greater proportions of subjects treated with ustekinumab 45 mg (67%; $p = 0.012$) or 90 mg (74%; $p < 0.001$) were PASI 75 responders at Week 12 compared with the etanercept group (57%). PASI 90 response was observed in 36% and 45 % of patients in the ustekinumab 45 mg and 90 mg groups, respectively, compared with 23% of patients receiving etanercept ($p<0.001$ for each comparison versus etanercept). PASI 100 response was observed in 12% and 21% of patients in the ustekinumab 45 mg and 90 mg groups, respectively, compared to 6% of patients receiving etanercept (Table 11). In addition, a greater proportion of patients in the ustekinumab 45 mg and 90 mg treatment groups achieved a PGA score of “cleared” or “minimal” (65 % and 71 %, respectively) compared with patients in the etanercept treatment group (49 %) ($p<0.001$ for each comparison versus etanercept).

In pre-specified analyses of efficacy by body weight in ACCEPT, minimal dose response to ustekinumab was evident in patients ≤ 100 kg. In patients who weighed >100 kg, higher PASI 75 response rates were seen with 90 mg dosing compared with 45 mg dosing, and a higher proportion of patients receiving 90 mg dosing had PGA scores of cleared or minimal compared with patients receiving 45 mg dosing (Table 16).

Table 16: Key psoriasis endpoints at Week 12: ACCEPT

| | ACCEPT | | |
|-----------------------------------|--------------------------------|---------------------------------|------------------------|
| | Etanercept (50mg twice a week) | Ustekinumab (week 0 and week 4) | |
| | | 45 mg | 90 mg |
| Patients randomized | 347 | 209 | 347 |
| PASI RESPONSE | | | |
| PASI 50 response | 286 (82%) | 181 (87%) | 320 (92%) ^a |
| PASI 75 response | 197 (57%) | 141 (67%) ^b | 256 (74%) ^a |
| PASI 90 response | 80 (23%) | 76 (36%) ^a | 155 (45%) ^a |
| PASI 100 response | 22 (6%) | 25 (12%) ^c | 74 (21%) ^a |
| PGA of Cleared or Minimal | 170 (49%) | 136 (65%) ^a | 245 (71%) ^a |
| PASI 75 RESPONSE BY WEIGHT | | | |
| ≤ 100 kg | | | |
| N | 251 | 151 | 244 |
| PASI 75 response | 154 (61%) | 109 (72%) | 189 (77%) |

| | | | |
|--|-----------|------------------------|------------------------|
| > 100 kg | | | |
| N | 96 | 58 | 103 |
| PASI 75 response | 43 (45%) | 32 (55%) | 67 (65%) |
| PGA OF CLEARED OR MINIMAL BY WEIGHT | | | |
| ≤ 100 kg | | | |
| N | 251 | 151 | 244 |
| PGA response | 131 (52%) | 110 (73%) | 185 (76%) |
| > 100 kg | | | |
| N | 96 | 58 | 103 |
| PGA response | 39 (41%) | 26 (45%) | 60 (58%) |
| PASI 75 RESPONSE BY NUMBER OF UNSUITABLE CONVENTIONAL SYSTEMIC AGENTS^b | | | |
| -at least one therapy | | | |
| N | 347 | 209 | 346 |
| PASI 75 Response | 197 (57%) | 141 (67%) ^b | 256 (74%) ^a |
| -at least two therapies | | | |
| N | 186 | 118 | 185 |
| PASI 75 Response | 94 (51%) | 79 (67%) ^d | 137 (74%) ^a |
| -at least three therapies | | | |
| N | 52 | 31 | 47 |
| PASI 75 Response | 20 (38%) | 17 (55%) ^e | 34 (72%) ^f |

^ap < 0.001 for ustekinumab 45 mg or 90 mg comparison with etanercept

^bp = 0.012 for ustekinumab 45 mg comparison with etanercept

^cp = 0.020 for ustekinumab 45 mg comparison with etanercept

^dp = 0.004 for ustekinumab 45 mg comparison with etanercept

^ep = 0.303 for ustekinumab 45 mg comparison with etanercept

^fp = 0.001 for ustekinumab 90 mg comparison with etanercept

^b Conventional systemic agents include psoralen plus ultraviolet A, MTX, and cyclosporine. Unsuitable conventional systemic agents are defined as those to which patients had had an inadequate response, were intolerant, or had a contraindication.

Clinical Efficacy – Pediatric plaque psoriasis

Adolescent patients (12 to 17 years of age)

The efficacy of STELARA was studied in 110 pediatric patients 12 to 17 years of age, in a multicenter, Phase 3, randomized, double blind, placebo controlled study (CADMUS). Patients were randomized to receive either placebo (n=37), or the recommended dose of STELARA (n=36) (see Dosage and Administration) or half the recommended dose of STELARA (n=37) by subcutaneous injection at Weeks 0 and 4 followed by every 12 week (q12w) dosing. At Week 12, placebo treated patients crossed over to receive STELARA. Efficacy observed in patients treated with the recommended dose of STELARA is presented below.

The baseline disease characteristics of randomized subjects are summarized in Table 16. Patients with PASI ≥ 12, PGA ≥ 3 and BSA involvement of at least 10%, who were candidates for systemic or phototherapy, were eligible for the study. Approximately half of the patients had prior exposure to conventional systemic or biologic therapy.

Table 17: Baseline Disease Characteristics in pediatric patients 12 to 17 years of age: CADMUS

| | <u>Placebo</u> | <u>STELARA*</u> |
|-------------------------------|----------------|-----------------|
| Patients randomized at Week 0 | N= 37 | N= 36 |
| Median Age (years) | 16.0 | 15.0 |
| Males | 20 (54.1%) | 16 (44.4%) |

| | | |
|---|---------------------|---------------------|
| Mean Weight (range; kg) | 64.74 (43.8; 107.0) | 62.00 (33.8; 109.5) |
| Median BMI (kg/m ²) | 22.70 | 22.15 |
| Median BSA | 21.0 | 21.5 |
| BSA ≥ 20% | 20 (54.1%) | 20 (55.6%) |
| Median PASI | 19.6 | 16.8 |
| Median CDLQI** (0-30) | 10.0 | 9.0 |
| Median PedsQL*** (0-100) | 77.17 | 79.35 |
| PGA of marked or severe | 15 (40.5%) | 12 (33.3%) |
| Psoriasis Disease Duration (years) | 5.11 | 5.54 |
| Prior topical therapy | 34 (91.9%) | 33 (91.7%) |
| Prior phototherapy | 11 (29.7%) | 14 (38.9%) |
| Prior conventional systemic therapy | 16 (43.2%) | 17 (47.2%) |
| Prior conventional systemic therapy or phototherapy | 22 (59.5%) | 22 (61.1%) |
| Prior biologic therapy | 5 (13.5%) | 3 (8.3%) |
| Prior conventional systemic or biologic therapy | 18 (48.6%) | 17 (47.2%) |

* Data presented for the recommended dose of STELARA

** CDLQI: The CDLQI is a dermatology instrument to assess the effect of a skin problem on the health-related quality of life in the pediatric population, with higher scores indicating greater negative effect on health-related quality of life.

*** PedsQL: The PedsQL is a general health-related quality of life measure developed for use in children and adolescent populations.

The primary endpoint was the proportion of patients who achieved a PGA score of cleared (0) or minimal (1) at Week 12. Secondary endpoints included PASI 75, PASI 90, change from baseline in Children's Dermatology Life Quality Index (CDLQI), change from baseline in the total score of PedsQL (Pediatric Quality of Life Inventory) at Week 12. At Week 12, subjects treated with STELARA showed significantly greater improvement in their psoriasis and health related quality of life compared with placebo (Table 18).

Table 18: Summary of Primary and Secondary End-points at Week 12: CADMUS (Age 12-17)

| | Placebo | STELARA* |
|---|--------------|---------------------------|
| | N (%) | N (%) |
| Patients randomized at Week 0 | 37 | 36 |
| Number of patients who achieved a PGA score of cleared (0) or minimal (1) | 2 (5.4%) | 25 (69.4%) ^a |
| PGA of Cleared (0) | 1 (2.7%) | 17 (47.2%) ^a |
| | | |
| PASI 75 responders | 4 (10.8%) | 29 (80.6%) ^a |
| PASI 90 responders | 2 (5.4%) | 22 (61.1%) ^a |
| PASI 100 responders | 1 (2.7%) | 14 (38.9%) ^a |
| | | |
| Change from baseline in CDLQI score | | |
| N | 32 | 32 |
| Mean (SD) | -1.5 (3.18) | -6.7 (5.63) |
| Median | 0.0 | -5.5 |
| CDLQI of 0 or 1** | 4 (13.3%) | 17 (56.7%) ^a |
| | | |
| Change from baseline in PedsQL Total Scale Score | | |
| N | 36 | 36 |
| Mean (SD) | 3.35 (10.04) | 8.03 (10.44) ^b |

* Data presented for the recommended dose of STELARA


^ap<0.001

^bp=0.028

** CDLQI of 0 or 1 indicates no effect on child's quality of life.

All patients were followed for efficacy for up to 52 weeks following first administration of study agent. The proportion of patients with a PGA score of cleared (0) or minimal (1) and the proportion achieving PASI 75 showed separation between the STELARA treated group and placebo at the first post-baseline visit at Week 4 reaching a maximum at Week 8. Improvements in PGA, PASI, CDLQI and PedsQL were maintained through Week 52. The PGA scores of cleared (0) or minimal (1) over time through Week 52 are summarized in Figure 3 below.

Figure 3: Percent of patients achieving PGA score of cleared (0) or minimal (1) through Week 52 by visit*.



*Data presented for the recommended dose of STELARA

Pediatric patients (Children 6 to 11 years of age)

The efficacy of STELARA was studied in 44 pediatric patients 6 to 11 years of age with moderate to severe plaque psoriasis in an open label, single-arm, multicenter, Phase 3 study (CADMUS Jr.). Patients were treated with the recommended dose of STELARA (n=44) (see *Dosage and Administration*) by subcutaneous injection at Weeks 0 and 4 followed by every 12 week (q12w) dosing. The baseline disease characteristics of enrolled patients are summarized in Table 19. Patients with PASI \geq 12, PGA \geq 3 and BSA involvement of at least 10%, who were candidates for systemic therapy or phototherapy, were eligible for the study. Approximately 23% of the patients had prior exposure to conventional systemic therapy or biologic therapy.

Table 19: Baseline Disease Characteristics in pediatric patients 6 to 11 years of age; CADMUS Jr.

| | STELARA* |
|---|---------------|
| Patients enrolled at Week 0 | N= 44 |
| Median Age (years) | 9.5 |
| Males | 17 (38.6%) |
| Mean Weight (range; kg) | 38.4 (19; 99) |
| Median BMI (kg/m ²) | 18.0 |
| Median BSA | 18.0 |
| BSA \geq 20% | 19 (43.2%) |
| Median PASI | 16.1 |
| Median CDLQI** (0-30) | 7.0 |
| PGA of marked or severe | 15 (34.1%) |
| Median Psoriasis Disease Duration (years) | 2.9 |
| Prior topical therapy | 43 (97.7%) |
| Prior phototherapy | 15 (34.1%) |
| Prior conventional systemic therapy | 8 (18.2%) |
| Prior conventional systemic therapy or phototherapy | 19 (43.2%) |
| Prior biologic therapy | 2 (4.5%) |
| Prior conventional systemic or biologic therapy | 10 (22.7%) |

* Data presented for the recommended dose of STELARA

** CDLQI: The CDLQI is a dermatology instrument to assess the effect of a skin problem on the health-related quality of life in the pediatric population, with higher scores indicating greater negative effect on health-related quality of life.

The primary endpoint was the proportion of patients who achieved a PGA score of cleared (0) or minimal (1) at Week 12. Secondary endpoints included PASI 75, PASI 90, and change from baseline in Children's Dermatology Life Quality Index (CDLQI) at Week 12. At Week 12, patients treated with STELARA showed clinically meaningful improvements in their psoriasis and health related quality of life (Table 20).

Table 20: Summary of Primary and Secondary End-points at Week 12 and 52: CADMUS Jr. (Age 6-11).

| | STELARA Week 12 | STELARA Week 52 |
|---|--------------------|--------------------|
| | N (%) | N (%) |
| Patients enrolled at Week 0 | 44 | 41 |
| Number of patients who achieved a PGA score of cleared (0) or minimal (1) | 34 (77.3%) | 31 (75.6%) |
| PGA of cleared (0) | 17 (38.6%) | 23 (56.1%) |
| | | |
| PASI 75 responders | 37 (84.1%) | 36 (87.8%) |
| PASI 90 responders | 28 (63.6%) | 29 (70.7%) |
| PASI 100 responders | 15 (34.1%) | 22 (53.7%) |
| | | |
| Patients with a CDLQI >1 at baseline | N = 39 | N = 36 |
| CDLQI of 0 or 1* | 24 (61.5%) | 21 (58.3%) |

*The CDLQI is a dermatology instrument to assess the effect of a skin problem on the health-related quality of life in the pediatric population. CDLQI of 0 or 1 indicates no effect on child's quality of life.

All patients were followed for efficacy for up to 52 weeks following first administration of study agent. Efficacy measured by PGA score of 0 or 1 was observed as early as the first post-baseline visit at Week 4 and increased through Week 16 and then remained relatively stable through Week 52. Improvements in PGA, PASI, and CDLQI were maintained through Week 52.

Clinical efficacy – Psoriatic arthritis (PsA)

The safety and efficacy of STELARA was assessed in two multicenter, randomized, double-blind, placebo-controlled, Phase 3 studies, PSUMMIT I and PSUMMIT II, in patients with active psoriatic arthritis. Patients were randomized to receive treatment with either STELARA 45 mg, 90 mg, or placebo subcutaneous injections at Weeks 0 and 4 followed by every 12 week (q12w) dosing. The primary endpoint in these studies was the reduction of ACR 20 responders at Week 24. Secondary endpoints included change from baseline in Disability Index of the Health Assessment Questionnaire (HAQ-DI), PASI 75, ACR 50, ACR 70 and change from baseline in total radiographic scores of the hands and feet, at Week 24. Efficacy data were collected and analyzed through Week 52 for both studies and through Week 100 for PSUMMIT I. These studies included 927 (PSUMMIT I n=615; PSUMMIT II, n=312) adult patients (\geq 18 years) who had active psoriatic arthritis (\geq 5 swollen joints and \geq 5 tender joints, despite modifying antirheumatic (DMARD) and/or nonsteroidal anti-inflammatory (NSAID) therapy). Methotrexate use was allowed during the studies but was not mandatory. Approximately 50% of patients continued on stable doses of MTX (\leq 25 mg/week). In PSUMMIT I and PSUMMIT II, 80% and 86% of the patients, respectively, had been previously treated with DMARDs.

In PSUMMIT I patients, who had been previously treated with anti-TNF α therapy, prior to the first study dose, were excluded. In PSUMMIT II, the majority of patients (58%, n=180) had been previously treated with one or more anti-TNF α agent(s) for at least 8 weeks (14 weeks with infliximab) or had discontinued anti-TNF α for intolerance at any time. Among the patients who had been previously treated with an anti-TNF α agent, over 70% had discontinued their anti-TNF α treatment for lack of efficacy or intolerance.

Patients with each subtype of psoriatic arthritis were enrolled, including polyarticular arthritis with no evidence of rheumatoid nodules (39%, N=362), spondylitis with peripheral arthritis (28%, N=255), asymmetric peripheral arthritis (21%, N=193), distal interphalangeal (DIP) arthritis (12%, N=112) and arthritis mutilans (0.5%, N=5). Over 70% and 40% of the patients in both studies had enthesitis and dactylitis at baseline, respectively.

In both studies, a significantly greater proportion of patients achieved ACR 20 and ACR 50 responses at Week 24 in the STELARA 45 mg and 90 mg groups compared to placebo (see Table 12). In the PSUMMIT I, a significantly greater proportion of patients

and in PSUMMIT II a numerically greater proportion of patients (p=NS) achieved ACR 70 responses in the STELARA 45 mg and 90 mg groups compared to placebo (see Table 21).

In both studies, the proportion of patients achieving a modified PsA response criteria (PsARC) or a Disease Activity Index Score 28 using C-reactive protein (DAS28-CRP) response was significantly greater in the STELARA 45 mg and 90 mg groups compared to placebo. In PSUMMIT I the proportion of patients achieving DAS28-CRP remission was significantly greater in the STELARA 45 mg and 90 mg groups compared to placebo. In PSUMMIT II, the proportion of patients who achieved DAS28-CRP remission was significantly greater in the STELARA 90 mg group compared to placebo (see Table 21). DAS28-CRP and PsARC responses were maintained through Week 52 in both studies and through Week 100 in PSUMMIT I.

| Table 21: Number of patients who achieved ACR 20, ACR 50, ACR 70, PsARC, DAS28-CRP response and DAS28-CRP remission at Week 24 | | | | | | |
|--|--------------------|------------------------|------------------------|--------------------|-----------------------|-----------------------|
| | PSUMMIT I | | | PSUMMIT II | | |
| | STELARA | | STELARA | | | |
| | Placebo (N=206) | 45 mg (N=205) | 90 mg (N=204) | Placebo (N=104) | 45 mg (N=103) | 90 mg (N=105) |
| ACR 20 | 47 (23%) | 87 (42%) ^a | 101 (50%) ^a | 21 (20%) | 45 (44%) ^a | 46 (44%) ^a |
| ACR 50 | 18 (9%) | 51 (25%) ^a | 57 (28%) ^a | 7 (7%) | 18 (17%) ^b | 24 (23%) ^a |
| ACR 70 | 5 (2%) | 25 (12%) ^a | 29 (14%) ^a | 3 (3%) | 7 (7%) ^c | 9 (9%) ^c |
| PsARC | 77 (37%) | 115 (56%) ^a | 132 (65%) ^a | 32 (31%) | 57 (55%) ^a | 54 (51%) ^b |
| DAS28-CRP * | 71 (34%) | 135 (66%) ^a | 138 (68%) ^a | 31 (30%) | 56 (54%) ^a | 56 (53%) ^a |
| DAS28 Remission ** | 17 (8%) | 42 (20%) ^a | 40 (20%) ^a | 4 (4%) | 11 (11%) ^c | 16 (15%) ^b |

^a p<0.001

^b p<0.05

^c p=NS

* Combining tender joints (28 joints), swollen joints (28 joints), CRP, and the Patient Global Assessment of disease activity using CRP.

DAS28 responders include patients with moderate or good response.

**DAS28 remitters include patients with a DAS28 value of < 2.6 at a visit.

An ACR 20 response (Felson et al, 1995) was defined as:

1.≥ 20% improvement in swollen joint count (66 joints) and tender joint count (68 joints); and

2.≥ 20% improvement in 3 of the following 5 assessments:

- Patient's assessment of pain [Visual Analog Scale (VAS)]
- Patient's global assessment of disease activity (VAS)
- Physician's global assessment of disease activity (VAS)
- Patient's assessment of physical function as measured by the HAQ-DI
- CRP

ACR 50 or ACR 70 are similarly defined.

The time course for ACR 20 response rates during the first 24 weeks in both studies for patients receiving STELARA or placebo are summarized in Figure 3. ACR 20 responses showed improvement at the first assessment (Week 4). ACR 20, 50 and 70 responses continued to improve or were maintained through Week 52 (see Table 21). In PSUMMIT I, ACR responses were maintained through Week 100.

Figure 4: Percent of patients achieving ACR 20 response through Week 24

PSUMMIT I

PSUMMIT II




Table 22: Proportion of patients who achieved ACR 20, ACR 50, ACR 70 response at Week 52.

| | PSUMMIT I | | PSUMMIT II | |
|--------------|-----------|-------|------------|-------|
| | STELARA | | STELARA | |
| | 45 mg | 90 mg | 45 mg | 90 mg |
| N | 194 | 189 | 94 | 95 |
| ACR response | | | | |
| ACR 20 | 55.7% | 60.3% | 46.8% | 48.4% |
| ACR 50 | 31.4% | 37.0% | 27.7% | 26.3% |
| ACR 70 | 18.0% | 21.2% | 12.8% | 17.9% |

In PSUMMIT I, of 205 subjects randomized to STELARA 45 mg, 153 continued the same dose and were available for evaluation at Week 52. Among those, ACR 20, 50 and 70 responses were achieved by 99 (64.7%), 57 (37.3%) and 34 (22.2%) subjects respectively. Of 204 subjects randomized to STELARA 90 mg, 185 were available for evaluation at Week 52. Among those, ACR 20, 50 and 70 responses were achieved by 120 (64.9%), 74 (40%) and 41 (22.2%) subjects respectively.

In PSUMMIT I, of 205 subjects randomized to STELARA 45 mg, 138 continued the same dose and were available for evaluation at Week 100. Among those, ACR 20, 50 and 70 responses were achieved by 89 (64.5%), 63 (45.7%) and 41 (29.7%) subjects respectively. Of 204 subjects randomized to STELARA 90 mg, 166 were available for evaluation at Week 100. Among those, ACR 20, 50 and 70 responses were achieved by 116 (69.9%), 84 (50.6%) and 41 (24.7%) subjects respectively.

In PSUMMIT II, of 103 subjects randomized to STELARA 45 mg, 68 continued the same dose and were available for evaluation at Week 52. Among those, ACR 20, 50, and 70 responses were achieved by 41 (60.3%), 23 (33.8%) and 11 (16.2%) subjects respectively. Of 105 subjects randomized to STELARA 90 mg, 83 were available for evaluation at Week 52. Among those, ACR 20, 50 and 70 responses were achieved by 49 (59%), 26 (31.3%) and 17 (20.5%) subjects respectively.

Additionally, within each weight group (≤ 100 kg and > 100 kg), ACR 20, ACR 50 and ACR 70 responses were consistently higher in the STELARA 45 and 90 mg groups than in the placebo group (see Table 23).

Table 23: Number of patients who achieved ACR 20, ACR 50 and ACR 70 responses by weight through Week 24

| | PSUMMIT I | | | PSUMMIT II | | |
|---|-----------------|----------------|----------------|------------------|----------------|----------------|
| | Placebo (N=206) | STELARA | | Placebo (N= 104) | STELARA | |
| | | 45 mg (N= 205) | 90 mg (N= 204) | | 45 mg (N= 103) | 90 mg (N= 105) |
| Patients randomized with weight ≤ 100 kg at baseline | 154 | 153 | 154 | 74 | 74 | 73 |

| | | | | | | |
|---|----------|----------|----------|----------|----------|----------|
| ACR 20 | 39 (25%) | 67 (44%) | 78 (51%) | 17 (23%) | 32 (43%) | 34 (47%) |
| ACR 50 | 14 (9%) | 38 (25%) | 48 (31%) | 6 (8%) | 15 (20%) | 21 (29%) |
| ACR 70 | 5 (3%) | 20 (13%) | 26 (17%) | 3 (4%) | 6 (8%) | 8 (11%) |
| Patients randomized with weight >100 kg at baseline | 52 | 52 | 50 | 30 | 29 | 31 |
| ACR 20 | 8 (15%) | 20 (38%) | 23 (46%) | 4 (13%) | 13 (45%) | 12 (39%) |
| ACR 50 | 4 (8%) | 13 (25%) | 9 (18%) | 1 (3%) | 3 (10%) | 3 (10%) |
| ACR 70 | 0 | 5 (10%) | 3 (6%) | 0 | 1 (3%) | 1 (3%) |

STELARA treatment resulted in significantly greater improvement compared with placebo for each ACR component (see Table 24).

Table 24: Summary of percent improvement from baseline in ACR components at Week 24

| | PSUMMIT I | | | PSUMMIT II | | |
|---|-----------|--------------------|--------------------|-------------------|--------------------|--------------------|
| | | STELARA | | | STELARA | |
| | | Placebo (N=206) | 45 mg (N= 205) | 90 mg (N= 204) | Placebo (N=104) | |
| Number of swollen joints ^d | | | | | | |
| Median | 21.54 | 58.82 ^a | 60.00 ^a | 0.00 | 52.94 ^b | 50.00 ^c |
| Number of tender joints ^e | | | | | | |
| Median | 13.61 | 45.45 ^a | 51.51 ^a | 0.00 | 33.33 ^a | 35.00 ^c |
| Patient's assessment of pain ^f | | | | | | |
| Median | 0.00 | 31.33 ^a | 42.58 ^a | 0.00 | 24.19 ^a | 24.29 ^a |
| Patient global assessment ^f | | | | | | |
| Median | 4.11 | 32.84 ^a | 42.44 ^a | 0.00 | 21.25 ^a | 22.54 ^a |
| Physician global assessment ^f | | | | | | |
| Median | 17.64 | 48.39 ^a | 55.91 ^a | 0.83 | 36.67 ^a | 36.11 ^a |
| Disability index (HAQ-DI) ^g | | | | | | |
| Median | 0.00 | 22.22 ^a | 32.46 ^a | 0.00 | 12.50 ^a | 14.29 ^a |
| CRP (mg/dL) ^h | | | | | | |
| Median | 0.00 | 38.56 ^a | 48.30 ^a | 0.00 | 25.61 ^c | 33.69 ^a |

^a p<0.001

^b p<0.05

^c p<0.01

^d Number of swollen joints counted (0-66)

^e Number of tender joints counted (0-68)

^f Visual analogue scale; 0= best, 10=worst.

^g Disability Index of the Health Assessment Questionnaire; 0 = best, 3 = worst, measures the patient's ability to perform the following: dress/groom, arise, eat, walk, reach, grip, maintain hygiene, and maintain daily activity.

^h CRP: (Normal Range 0.0-1.0 mg/dL)

Methotrexate Use

The proportion of patients achieving ACR responses were consistently greater in patients treated with STELARA than those treated with placebo regardless of concomitant MTX use (see Table 25). Responses observed in the STELARA groups were similar in patients receiving or not receiving concomitant MTX. ACR responses were maintained through Week 52 in PSUMMIT I and II and through Week 100 in PSUMMIT I.

Table 25: Summary of patients achieving ACR 20, ACR 50 and ACR 70 responses through Week 24 by methotrexate usage

| | PSUMMIT I | | | | | | |
|---------------------|--------------------|---------------------------|-------------------|--------------------|-------------------------------|-------------------|--|
| | | Receiving MTX at baseline | | | Not receiving MTX at baseline | | |
| | | STELARA | | | STELARA | | |
| | Placebo (N=206) | 45 mg (N= 205) | 90 mg (N= 204) | Placebo (N=206) | 45 mg (N= 205) | 90 mg (N= 204) | |
| Patients randomized | 96 | 99 | 101 | 110 | 106 | 103 | |
| ACR 20 | 25 (26%) | 43 (43%) | 46 (46%) | 22 (20%) | 44 (42%) | 55 (53%) | |
| ACR 50 | 8 (8%) | 23 (23%) | 27 (27%) | 10 (9%) | 28 (26%) | 30 (29%) | |

| | | | | | | |
|---------------------|----------------------------------|---------------------------|---------------------------|--------------------------------------|---------------------------|---------------------------|
| ACR 70 | 2 (2%) | 11 (11%) | 13 (13%) | 3 (3%) | 14 (13%) | 16 (16%) |
| PSUMMIT II | | | | | | |
| | <i>Receiving MTX at baseline</i> | | | <i>Not receiving MTX at baseline</i> | | |
| | STELARA | | | | STELARA | |
| | Placebo (N=104) | 45 mg (N= 103) | 90 mg (N= 105) | Placebo (N=104) | 45 mg (N= 103) | 90 mg (N= 105) |
| Patients randomized | 49 | 54 | 52 | 55 | 49 | 53 |
| ACR 20 | 14 (29%) | 27 (50%) | 21 (40%) | 7 (13%) | 18 (37%) | 25 (47%) |
| ACR 50 | 4 (8%) | 10 (19%) | 12 (23%) | 3 (5%) | 8 (16%) | 12 (23%) |
| ACR 70 | 2 (4%) | 4 (7%) | 3 (6%) | 1 (2%) | 3 (6%) | 6 (11%) |

Prior Anti-TNF α therapy

PSUMMIT II evaluated 180 patients who were previously treated with one or more anti-TNF α agents for at least 8 weeks (14 weeks with infliximab), or had documented intolerance of anti-TNF α therapy at any time in the past.

Among patients previously treated with anti-TNF α agents, a significantly greater proportion of STELARA-treated patients achieved an ACR 20 response at Week 24 compared to placebo (see Table 26). ACR 20, 50 and 70 responses were generally maintained through Week 52.

Table 26: Number of patients previously treated with anti-TNF α agent(s) who achieved ACR 20, ACR 50 and ACR 70 responses through Week 24

| PSUMMIT II | | STELARA | | |
|---------------------|---------|-----------------------|-----------------------|-------------------|
| | | Placebo (N= 104) | 45 mg (N= 103) | 90 mg (N= 105) |
| Patients randomized | 62 | 60 | 58 | |
| ACR 20 | 9 (15%) | 22 (37%) ^a | 20 (34%) ^b | |
| ACR 50 | 4 (6%) | 9 (15%) ^c | 9 (16%) ^c | |
| ACR 70 | 1 (2%) | 3 (5%) ^c | 3 (5%) ^c | |

^a p<0.01

^b p<0.05

^c p=NS

Enthesitis and Dactylitis

For patients with enthesitis and/or dactylitis at baseline, in PSUMMIT I, a significant improvement in enthesitis and dactylitis score was observed in the STELARA 45 mg and 90 mg groups compared to placebo. In PSUMMIT II, a significant improvement in enthesitis score and numerical improvement in dactylitis score were observed in the 90 mg group (p=NS) compared with the placebo group (see Table 27). In both studies, improvement in enthesitis score and dactylitis score were maintained at Week 52. In PSUMMIT I, the improvement in enthesitis score and dactylitis score was maintained through Week 100.

Table 27: Summary of percent change in enthesitis and dactylitis scores at Week 24

| | | PSUMMIT I | | | PSUMMIT II | | |
|---|------|---------------------|---------------------|------------------|---------------------|---------------------|-------------------|
| | | STELARA | | | STELARA | | |
| | | Placebo (N=206) | 45 mg (N=205) | 90 mg (N=204) | Placebo (N= 104) | 45 mg (N= 103) | 90 mg (N= 105) |
| Enthesitis score^d | | | | | | | |
| Patients randomized with enthesitis at baseline | 145 | 142 | 154 | 73 | 72 | 76 | |
| N | 137 | 140 | 148 | 68 | 70 | 70 | |
| Median | 0.00 | -42.86 ^a | -50.00 ^b | 0.00 | -33.33 ^c | -48.33 ^a | |
| Dactylitis score^e | | | | | | | |
| Patients randomized with dactylitis at baseline | 96 | 101 | 99 | 38 | 48 | 41 | |
| N | 92 | 99 | 95 | 33 | 46 | 38 | |
| Median | 0.00 | -75.00 ^b | -70.83 ^b | 0.00 | 0.00 ^c | -64.58 ^c | |

^a p<0.01

^b p<0.001

^c p=NS

^d Enthesitis was assessed based on the Maastricht Ankylosing Spondylitis Enthesis Score (MASES) index modified for PSA (an instrument that counts 15 body sites).

^e Dactylitis was assessed in both hands and feet using a scoring system from 0 to 60.

A higher proportion of patients treated with STELARA, that have spondylitis with peripheral arthritis as their primary presentation, demonstrated Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) 50 and 70 percent improvement in BASDAI scores at Week 24 compared with placebo (see Table 28).

| Table 28: Number of patients who achieved improvement from baseline in BASDAI at Week 24 | | | | | | |
|--|---------------------|-----------------------|-----------------------|---------------------|-----------------------|-----------------------|
| | PSUMMIT I | | | PSUMMIT II | | |
| | | STELARA | | | STELARA | |
| | Placebo (N= 206) | 45 mg (N=205) | 90 mg (N=204) | Placebo (N= 104) | 45 mg (N=103) | 90 mg (N=105) |
| Patients randomized with spondylitis and peripheral joint involvement at baseline | 70 | 52 | 64 | 22 | 26 | 22 |
| N | 61 | 51 | 60 | 18 | 25 | 21 |
| BASDAI 20 | 16 (26%) | 25 (49%) ^a | 35 (58%) ^b | 10 (56%) | 15 (60%) ^c | 11 (52%) ^c |
| BASDAI 50 | 8 (13%) | 12 (24%) ^c | 19 (32%) ^a | 1 (6%) | 7 (28%) ^c | 8 (38%) ^a |
| BASDAI 70 | 0 | 7 (14%) ^d | 9 (15%) ^d | 0 | 3 (12%) [*] | 5 (24%) [*] |

^a p≤0.05

^b p<0.001

^c p=NS

^d p≤0.01

*p value not calculated

PASI Response

In PSUMMIT I and PSUMMIT II, the proportion of patients with psoriasis involvement of ≥3% BSA at baseline who achieved a ≥75% improvement in the PASI assessment at Week 24 was significantly greater in the STELARA 45 mg and 90 mg groups compared with the placebo group (see Table 20). In both studies the proportion of patients achieving the PASI 75 response was maintained through Week 52 (PSUMMIT I, STELARA 45mg-70.1% and 90mg- 68.1%; PSUMMIT II, STELARA 45mg-56.5% and 90mg- 64.4%). In PSUMMIT I, the PASI 75 response was maintained through Week 100.

The proportion of patients who achieved both a PASI 75 response and an ACR 20 response was evaluated for those patients with ≥3% BSA psoriasis skin involvement at baseline. A significantly higher proportion of patients achieved the combined response in the STELARA 45 mg and 90 mg groups compared with the placebo group at Week 24 (see Table 29). In both studies the proportion of patients achieving both a PASI 75 response and an ACR20 response was maintained through Week 52 (PSUMMIT I, STELARA 45mg-44.8% and 90mg-44.3%; PSUMMIT II, STELARA 45mg-36.8% and 90mg- 43.1%). In PSUMMIT I, the proportion of patients achieving the combined PASI 75 and ACR20 response was maintained through Week 100.

Table 29: Number of patients who achieved PASI 75, PASI 90 and PASI 100 responses as well as a combination of skin and joint responses at Week 24

| | PSUMMIT I | | | PSUMMIT II | | |
|--|---------------------|----------------------|------------------|---------------------|----------------------|------------------|
| | | STELARA ^a | | | STELARA ^a | |
| | Placebo (N= 206) | 45 mg (N=205) | 90 mg (N=204) | Placebo (N= 104) | 45 mg (N=103) | 90 mg (N=105) |
| Patients with ≥3% BSA psoriasis skin involvement at baseline | 146 | 145 | 149 | 80 | 80 | 81 |
| PASI 75 | 16 (11%) | 83 (57%) | 93 (62%) | 4 (5%) | 41 (51%) | 45 (56%) |
| PASI 90 | 4 (3%) | 60 (41%) | 65 (44%) | 3 (4%) | 24 (30%) | 36 (44%) |
| PASI 100 | 2 (1%) | 29 (20%) | 41 (28%) | 1 (1%) | 13 (16%) | 17 (21%) |
| Combination of skin and joint responses | | | | | | |
| PASI 75 and ACR 20 | 8 (5%) | 40 (28%) | 62 (42%) | 2 (3%) | 24 (30%) | 31 (38%) |

^a p<0.001 for 45 mg or 90 mg comparison with placebo.

Additionally, within each weight group (≤100 kg and >100 kg), PASI 75, 90 and 100 responses were consistently higher in the STELARA 45 and 90 mg groups than in the placebo group (see Table 30).

| Table 30: Summary of patients who achieved PASI 75, PASI 90 and PASI 100 responses by weight through Week 24 | | | | | | |
|---|--------------------|-------------------|-------------------|---------------------|-------------------|-------------------|
| | PSUMMIT I | | | PSUMMIT II | | |
| | STELARA | | STELARA | | | |
| | Placebo (N=206) | 45 mg (N= 205) | 90 mg (N= 204) | Placebo (N= 104) | 45 mg (N= 103) | 90 mg (N= 105) |
| Patients randomized with weight \leq 100 kg at baseline* | 105 | 105 | 111 | 54 | 58 | 57 |
| PASI 75 | 14 (13%) | 64 (61%) | 73 (66%) | 4 (7%) | 31 (53%) | 32 (56%) |
| PASI 90 | 4 (4%) | 46 (44%) | 48 (43%) | 3 (6%) | 20 (34%) | 27 (47%) |
| PASI 100 | 2 (2%) | 21 (20%) | 30 (27%) | 1 (2%) | 11 (19%) | 13 (23%) |
| Patients randomized with weight $>$ 100 kg at baseline* | 41 | 40 | 38 | 26 | 22 | 24 |
| PASI 75 | 2 (5%) | 19 (48%) | 20 (53%) | 0 | 10 (45%) | 13 (54%) |
| PASI 90 | 0 | 14 (35%) | 17 (45%) | 0 | 4 (18%) | 9 (38%) |
| PASI 100 | 0 | 8 (20%) | 11 (29%) | 0 | 2 (9%) | 4 (17%) |

* Patients randomized with \geq 3% BSA psoriasis skin involvement at baseline

Methotrexate Use

In both studies, the proportion of patients who achieved a PASI 75 response at Week 24 was consistently higher in STELARA 45 mg and 90 mg groups compared with placebo regardless of concomitant MTX use. PASI 75 responses were maintained through Week 52 in both PSUMMIT I and II. In PSUMMIT I, PASI 75 response was maintained at Week 100.

Prior Anti-TNF α Therapy

In PSUMMIT II, the proportion of patients who achieved a PASI 75 response at Week 24 was significantly greater in STELARA 45 mg and 90 mg groups compared with placebo in patients previously treated with an anti-TNF α agent.

Radiographic Response

Structural damage in both hands and feet was assessed by readers unaware of treatment group and order of visits, and expressed as change in total van der Heijde-Sharp score (vdH-S score), modified for PsA by addition of hand distal interphalangeal (DIP) joints, compared to baseline. A pre-specified integrated analysis combining data from 927 subjects in both PSUMMIT I & II was performed. At Week 24, based on this integrated analysis, the STELARA 45 mg or 90 mg treatment significantly inhibited progression of structural damage, when compared to placebo (see Table 22). Beyond Week 24, STELARA treatment continued to inhibit the progression of structural damage through Week 52. The mean change from Week 24 to 52 in total modified vdH-S score (0.18 and 0.26 in the STELARA 45mg and 90 mg groups respectively) was less than the mean change from Week 0 to 24 (see Table 31). In PSUMMIT I, the effect of STELARA on inhibition of structural damage progression was maintained through Week 100. Among subjects treated with STELARA 45 mg and 90 mg with no radiographic progression from baseline to Week 52 (n=103, and 113, respectively), 81.5% and 88.8% continued to show no radiographic progression at Week 100.

Table 31: Summary of change from baseline in total modified vdH-S score at Week 24 (Integrated analysis of PSUMMIT I and PSUMMIT II)

| | STELARA® | | |
|--|--------------------|-------------------------------|-------------------------------|
| | Placebo | 45mg | 90mg |
| Total Modified vdH-S score at Baseline | | | |
| N | 306 | 303 | 300 |
| Mean \pm SD | 28.01 \pm 55.771 | 30.40 \pm 50.688 | 27.97 \pm 42.137 |
| Change from Baseline | | | |
| N | 310 | 308 | 309 |
| Mean \pm SD | 0.97 \pm 3.852 | 0.40 \pm 2.110 ^b | 0.39 \pm 2.403 ^a |

^a p value < 0.001 for the difference between STELARA® and Placebo, Week 24 (integrated analysis)

^b p value < 0.05

At Week 24, patients treated with STELARA demonstrated less progression of structural damage compared to placebo, irrespective of concomitant MTX use.

The effect of STELARA on progression of structural damage in patients with prior anti-TNF α experience has not been established although it has not been adequately studied.

Physical Function and Health-Related Quality of Life

In PSUMMIT I and PSUMMIT II, physical function and health-related quality of life were assessed using the Disability Index of the Health Assessment Questionnaire (HAQ-DI), Dermatology Life Quality Index (DLQI) and the SF-36 health survey.

Patients treated with STELARA showed significant improvement in physical function as assessed by the HAQ-DI at Week 24. The proportion of patients achieving a clinically meaningful ≥ 0.3 improvement in HAQ-DI score from baseline at Week 24 was also significantly greater in the STELARA groups when compared with placebo (see Table 32). Improvement was observed at the first assessment (Week 4), reached maximum at Week 12 and was maintained through Week 24. Improvement in HAQ-DI score from baseline was maintained in both studies at Week 52 and through Week 100 in PSUMMIT I.

In both studies, the improvement in HAQ-DI at Week 24 was consistently greater in the STELARA 45 mg and 90 mg groups compared with placebo regardless of concomitant MTX use.

In PSUMMIT II, the improvement in HAQ-DI at Week 24 was significantly greater in the STELARA 45 mg and 90 mg groups compared with placebo in patients previously treated with anti-TNF α agents.

| Table 32: Improvement in physical function as measured by HAQ-DI at Week 24 | | | | | | |
|--|---------------------|-----------------------|-----------------------|---------------------|-----------------------|-----------------------|
| | PSUMMIT I | | | PSUMMIT II | | |
| | Placebo (N= 206) | 45 mg (N=205) | 90 mg (N=204) | Placebo (N= 104) | 45 mg (N=103) | 90 mg (N=105) |
| HAQ-DI Baseline Score | | | | | | |
| N | 204 | 205 | 204 | 104 | 103 | 104 |
| Mean (SD) | 1.24 (0.647) | 1.22 (0.610) | 1.22 (0.634) | 1.25 (0.723) | 1.34 (0.704) | 1.29 (0.666) |
| Median | 1.25 | 1.25 | 1.25 | 1.25 | 1.38 | 1.25 |
| Improvement in HAQ-DI | | | | | | |
| N | 206 | 205 | 204 | 104 | 103 | 105 |
| Mean (SD) | 0.10 (0.390) | 0.31 (0.521) | 0.40 (0.514) | 0.03 (0.380) | 0.21 (0.461) | 0.22 (0.436) |
| Median | 0.00 | 0.25 ^a | 0.25 ^a | 0.00 | 0.13 ^b | 0.25 ^a |
| HAQ-DI Responders* | 58 (28%) | 98 (48%) ^a | 97 (48%) ^a | 17 (16%) | 35 (34%) ^b | 40 (38%) ^a |

^a p<0.001

^b p<0.01

*achieving a ≥ 0.3 improvement from baseline

In PSUMMIT I, of 205 subjects randomized to STELARA 45 mg, 153 continued the same dose and were available for evaluation at Week 52. Among those, the HAQ-DI response was achieved by 83 (54.2%) subjects. Of 204 subjects randomized to STELARA 90 mg, 185 were available for evaluation at Week 52. Among those, HAQ-DI response was achieved by 102 (55.1%) subjects.

In PSUMMIT II, of 103 subjects randomized to STELARA 45 mg, 68 continued the same dose and were available for evaluation at Week 52. Among those, the HAQ-DI response was achieved by 29 (42.6%) subjects. Of 105 subjects randomized to STELARA 90 mg, 83 were available for evaluation at Week 52. Among those, HAQ-DI response was achieved by 44 (53%) subjects.

The DLQI was assessed by comparing the change in DLQI scores from baseline for those patients with $\geq 3\%$ BSA at baseline. In both studies at Week 24, there was a significant improvement from baseline in DLQI scores in both the STELARA 45 mg and 90 mg groups as compared with placebo (see Table 33) and the improvement was maintained at Week 52. In PSUMMIT I, the improvement from baseline in DLQI scores was maintained through Week 100.

In both PSUMMIT I and PSUMMIT II, at Week 24, the change from baseline in the SF-36 physical component summary (PCS) scores was significantly greater in the STELARA 45 mg and 90 mg groups compared with the placebo group. In both studies, the change from baseline in the SF-36 mental component summary (MCS) scores at Week 24 was greater in both STELARA groups compared with the placebo group (p<0.001 for PSUMMIT I - 90mg group, p=NS for other groups) (see Table 33). The change from baseline in the SF-36 PCS and MCS scores was maintained at Week 52 in both studies, and at Week 100 in PSUMMIT I.

In PSUMMIT II, a significant change from baseline in Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F) scores was observed at Week 24 in the STELARA 45 mg and 90 mg groups compared with the placebo group (median improvement, all 3.0 vs 0.0; $p<0.007$). Similarly, the percentage of patients with clinically significant improvement in fatigue from baseline (4 points in FACIT-F) was significantly greater in the STELARA 45 mg (49% [$p<0.001$]) and 90 mg groups (49% [$p<0.001$]) compared with the placebo group (25.8%). The change from baseline in the FACIT-F scores was maintained at Week 52.

Table 33: Summary of change from baseline in DLQI and SF-36 and scores at Week 24

| | PSUMMIT I | | | PSUMMIT II | | |
|--|---------------------|--------------------|--------------------|---------------------|--------------------|--------------------|
| | STELARA | | STELARA | | STELARA | |
| | Placebo (N= 206) | 45 mg (N=205) | 90 mg (N=204) | Placebo (N= 104) | 45 mg (N=103) | 90 mg (N=105) |
| DLQI | | | | | | |
| Patients randomized with $\geq 3\%$ BSA psoriasis skin involvement at baseline | 146 | 145 | 149 | 80 | 80 | 81 |
| Baseline | | | | | | |
| N | 145 | 145 | 149 | 80 | 80 | 81 |
| Mean (SD) | 11.68 (7.705) | 11.02 (7.308) | 10.54 (7.179) | 11.93 (7.622) | 12.09 (7.667) | 11.98 (7.754) |
| Median | 11.00 | 10.00 | 9.00 | 11.00 | 11.00 | 10.00 |
| Change from baseline | | | | | | |
| N | 140 | 142 | 146 | 73 | 77 | 75 |
| Mean (SD) | -1.40 (6.177) | -6.63 (6.776) | -7.54 (6.524) | -0.75 (5.666) | -6.95 (7.719) | -7.16 (6.748) |
| Median | -1.00 | -6.00 ^a | -6.00 ^a | 0.00 | -6.00 ^a | -6.00 ^a |
| SF-36 | | | | | | |
| Physical component summary | | | | | | |
| Baseline | | | | | | |
| N | 203 | 203 | 204 | 104 | 102 | 104 |
| Mean (SD) | 31.39 (8.785) | 31.16 (8.511) | 31.45 (8.152) | 30.28 (9.361) | 28.69 (8.501) | 28.93 (8.480) |
| Median | 30.40 | 29.80 | 29.70 | 29.35 | 27.95 | 28.15 |
| Change from baseline | | | | | | |
| N | 196 | 200 | 197 | 97 | 99 | 97 |
| Mean (SD) | 1.4(7.094) | 4.89 (9.333) | 6.22 (8.747) | 1.09 (5.892) | 4.29 (8.594) | 4.67 (8.758) |
| Median | 1.15 | 3.90 ^a | 5.80 ^a | 0.00 | 2.70 ^c | 3.50 ^a |
| Mental component summary | | | | | | |
| Baseline | | | | | | |
| N | 203 | 203 | 204 | 104 | 102 | 104 |
| Mean (SD) | 43.51 (10.848) | 42.77 (10.908) | 43.48 (11.608) | 42.11 (12.507) | 43.27 (12.911) | 42.81 (11.953) |
| Median | 43.90 | 42.00 | 41.65 | 41.80 | 43.70 | 41.40 |
| Change from baseline | | | | | | |
| N | 196 | 200 | 197 | 97 | 99 | 97 |
| Mean (SD) | 1.53 (9.582) | 3.35 (10.016) | 4.79 (10.054) | 0.63 (8.238) | 3.01 (11.144) | 3.52 (11.274) |
| Median | 0.25 | 2.65 ^b | 4.40 ^a | 0.00 | 0.70 ^b | 2.20 ^b |

^a $p\leq 0.001$

^b $p=NS$

^c $p<0.05$

Health Economics

Health economics data on time lost from work, employability, and daily productivity at work, school, or home were collected through questionnaires at baseline and Week 24. To assess productivity, patients were asked to indicate how much their disease affected their productivity at work, school or at home in the past 4 weeks, using a 10 cm Visual Analogue Scale (VAS) (not at all affected [0] to affected very much [10]).

The improvement in self-reported productivity was significantly greater in the STELARA 45 mg and 90 mg groups compared to placebo at Week 24. The improvement in self-reported productivity was maintained in both studies at Week 52 and through Week 100 in PSUMMIT I.

Clinical Efficacy – Crohn's Disease

The safety and efficacy of STELARA were evaluated in three randomized, double-blind, placebo-controlled clinical trials in adult patients with moderately to severely active Crohn's disease (Crohn's Disease Activity Index [CDAI] score of 220 to 450). The clinical development program consisted of two 8-week IV induction studies (UNITI-1 and UNITI-2) followed by a 44-week subcutaneous randomized withdrawal maintenance study (IM-UNITI) representing 52 weeks of therapy.

Induction of Clinical Response and Remission

UNITI-1 and UNITI-2 studies included 1409 (UNITI-1, n=769; UNITI-2 n=640) patients. In both studies, patients were permitted to concomitantly receive oral 5-ASA compounds, immunomodulators, corticosteroids, and/or antibiotics. Patients were randomized to receive a single IV administration of either 130 mg STELARA, or approximately 6 mg/kg STELARA designed as a tiered dose based on patient body weight (Table 1) or placebo at Week 0. The primary endpoint was clinical response (defined as a reduction in CDAI score of ≥ 100 points or CDAI score < 150) at Week 6. Secondary endpoints included clinical remission at Week 8, clinical response at Week 8, 70-point response at Week 3, and 70-point response at Week 6. Efficacy data were collected and analyzed through Week 8 for both studies.

In UNITI-1, patients had failed or were intolerant to prior anti-TNF α therapy. At baseline, approximately 46% (n=340) patients were receiving corticosteroids (including budesonide) and 31.4% of patients were receiving immunomodulators. Approximately 48% had failed 1 prior anti-TNF α therapy and 52% had failed 2 or 3 prior anti-TNF α therapies (40.8% and 10.4%, respectively). In this study, 29.1% patients had an inadequate initial response (primary non-responders), 69.4% responded but subsequently lost response (secondary non-responders), and 36.4% were intolerant to anti-TNF α therapies.

Patients in UNITI-2 had failed at least one conventional therapy (corticosteroids or immunomodulators) and were either anti-TNF α naïve (68.6%) or had previously received but not failed anti-TNF α therapy (31.4%). At baseline, approximately 40% patients were receiving corticosteroids (including budesonide) and 35% patients were receiving immunomodulators.

In these induction studies, efficacy was higher and better sustained in the tiered dose group compared to the 130 mg dose group, and tiered dosing is therefore the recommended IV induction dose. In both UNITI-1 and UNITI-2, a significantly greater proportion of patients were in clinical response and remission in the group treated with STELARA, compared to placebo (Table 34, Figure 5). Clinical response and remission were significant as early as Week 3 in STELARA treated patients and continued to improve through Week 8 (Figure 5).

| Table 34: Induction of Clinical Response and Remission in UNITI-1* and UNITI 2** | | | | |
|---|--------------------------|--------------------------|--------------------------|--------------------------|
| | UNITI-1 | | UNITI-2 | |
| | Placebo N=247 | STELARA N=249 | Placebo N=209 | STELARA N=209 |
| Clinical Remission, Week 8 | 18 (7.3%) | 52 (20.9%) ^a | 41 (19.6%) | 84 (40.2%) ^a |
| Clinical Response (100 point), Week 6 | 53 (21.5%) | 84 (33.7%) ^b | 60 (28.7%) | 116 (55.5%) ^a |
| Clinical Response (100 point), Week 8 | 50 (20.2%) | 94 (37.8%) ^a | 67 (32.1%) | 121 (57.9%) ^a |
| 70 Point Response, Week 3 | 67 (27.1%) | 101 (40.6%) ^b | 66 (31.6%) | 106 (50.7%) ^a |
| 70 Point Response, Week 6 | 75 (30.4%) | 109 (43.8%) ^b | 81 (38.8%) | 135 (64.6%) ^a |

Clinical remission is defined as CDAI score < 150 ; Clinical response is defined as reduction in CDAI score by at least 100 points or being in clinical remission
70 point response is defined as reduction in CDAI score by at least 70 points
* Anti-TNF α failures
** Conventional therapy failures
^a p < 0.001
^b p < 0.01




Figure 5: Proportion of STELARA treated patients in clinical response (A, B) and remission (C, D) through Week 8 in UNITI-1 and UNITI-2 studies

Maintenance of Response and Remission

The maintenance study (IM-UNITI) evaluated 388 patients who achieved clinical response (≥ 100 point reduction in CDAI score) at Week 8 of induction with STELARA in UNITI-1 or UNITI-2. Of those, approximately 60% of the patients entered the maintenance study in remission. Patients were randomized to receive a subcutaneous maintenance regimen of either 90 mg STELARA every 8 weeks, 90 mg STELARA every 12 weeks or placebo for 44 weeks.

Concomitant doses of oral 5-ASA compounds, immunomodulators, corticosteroids and antibiotics were permitted. Corticosteroids were tapered at the start of the maintenance trial. The primary endpoint was clinical remission (CDAI < 150) at Week 44. Secondary endpoints assessed at Week 44 included clinical response, clinical remission among STELARA treated patients in clinical remission after induction, corticosteroid-free remission, and clinical remission in the subset of patients who were refractory or intolerant to anti-TNF α treatment.

Significantly higher proportions of patients maintained clinical remission and response in the STELARA treated groups as compared to placebo at Week 44 (Table 35, Figure 6). A higher proportion of STELARA treated patients compared to placebo achieved sustained clinical remission (clinical remission at Week 36, 40 and 44).

Table 35: Maintenance of Clinical Response and Remission in IM-UNITI (Week 44; 52 weeks from initiation of the induction dose)

| | Placebo* | 90 mg STELARA every 8 weeks | 90 mg STELARA every 12 weeks |
|--|----------|-----------------------------|------------------------------|
| | | | |

| | N=131 [†] | N=128 [†] | N=129 [†] |
|---|--------------------|--------------------------|--------------------|
| Clinical Remission | 36% | 53% ^a | 49% ^b |
| Clinical Response | 44% | 59% ^b | 58% ^b |
| Corticosteroid-Free Clinical Remission | 30% | 47% ^a | 43% ^c |
| Sustained Clinical Remission [‡] | 26% | 46% ^c | 40% ^c |
| Clinical Remission in patients: | | | |
| in remission at the start of maintenance therapy | 46% (36/79) | 67% (52/78) ^a | 56% (44/78) |
| who are Anti-TNF α refractory/intolerant | 26% (16/61) | 41% (23/56) | 39% (22/57) |
| who failed conventional therapy but not anti-TNF α therapy | 44% (31/70) | 63% (45/72) ^c | 57% (41/72) |
| who are Anti-TNF α naïve | 49% (25/51) | 65% (34/52) ^c | 57% (30/53) |

Clinical remission is defined as CDAI score < 150; Clinical response is defined as reduction in CDAI of at least 100 points or being in clinical remission

* The placebo group consisted of patients who were in response to STELARA and were randomized to receive placebo at the start of maintenance therapy.

† Patients who achieved a clinical response to STELARA at start of maintenance therapy

‡ Defined as clinical remission at Week 36, 40 and 44.

^a p < 0.01

^b p < 0.05

^c nominally significant (p<0.05)




Figure 6: Proportion of patients in clinical remission at each visit through Week 44.

Delayed response

Patients who were not in clinical response to STELARA induction received a 90 mg subcutaneous injection of STELARA upon entry into the maintenance study. Eight weeks later, 50.5% of the patients achieved clinical response and continued to receive maintenance dosing every 8 weeks; among these patients with continued maintenance dosing, a majority achieved levels of response (68.1%) and remission (50.2%) similar to the patients who initially responded to STELARA induction.

Dosing in patients with a lower inflammatory burden

In patients with a lower inflammatory burden as reflected by CRP ≤ 10 mg/L at initiation of induction or initiation of maintenance therapy, the efficacy of the every 12 week dosing regimen was similar to that of the every 8 week dosing regimen.

Dosing frequency adjustment

In IM-UNITI, patients who did not maintain response to STELARA when treated every 12 weeks were allowed to increase the frequency of dosing and receive STELARA every 8 weeks. In these patients, clinical remission was achieved in 41.4% of patients 16 weeks after dosing frequency adjustment.

Resumption of treatment

Patients that responded to STELARA induction and who were randomized to the placebo group at the start of the maintenance study received 90 mg STELARA subcutaneously every 8 weeks at time of loss of response. Of these patients, 70.6% achieved clinical response and 39.2% achieved clinical remission 16 weeks after receiving the first subcutaneous dose of STELARA.

Long-Term Maintenance

In IM-UNITI, patients who completed the study through week 44 were eligible to continue treatment in a study extension. Among patients who entered the study extension, clinical remission and response were generally maintained through week 252. Results were consistent between patients who failed TNF-therapies versus those who did not.

No new safety concerns were identified in this study extension with up to 5 years of treatment in patients with Crohn's Disease.

Corticosteroid Use in maintenance

In patients that were in clinical response to STELARA induction therapy, a greater proportion of patients in the STELARA treated group were in remission and corticosteroid-free compared to the placebo group after 44 weeks of maintenance treatment (Table 35). In addition, a higher proportion of patients were in clinical response and not receiving corticosteroids in the STELARA treated group compared to placebo.

Endoscopic Healing of the Mucosa

Endoscopic healing of the mucosa was evaluated in 252 patients with baseline endoscopic disease activity in a substudy. At Week 8, after a single IV induction dose, reduction in mucosal inflammation, as measured by the Simplified Endoscopic Activity Score for Crohn's Disease (SES-CD), was greater in patients treated with STELARA (n=83) compared with patients treated with placebo (n=97) (-2.8 vs -0.7, p=0.009). Similar reductions in histologic inflammation were also observed.

Reduction in endoscopic and histologic inflammation was observed in patients treated with STELARA in maintenance. However, due to the small number of patients, the efficacy of STELARA in the maintenance of endoscopic healing could not be definitively established.

Fistula Response

In patients with draining fistulas at baseline (8.8%), a numerically greater proportion of STELARA treated patients achieved a fistula response (defined as $\geq 50\%$ reduction from baseline of the induction study in the number of draining fistulas) compared with placebo over 44 weeks (p=NS). The proportion of patients in fistula response at Week 44 was 45.5% (5/11) for placebo group, 71.4% (5/7) for STELARA 90 mg every 12 week dosing group, and 87.5% (7/8) for STELARA 90 mg every 8 week dosing group.

Health-Related Quality of Life Measures

Improvement in general and disease specific health-related quality of life was assessed using the SF-36 and Inflammatory Bowel Disease Questionnaire (IBDQ) respectively.

SF-36

A higher proportion of patients treated with STELARA showed clinically meaningful improvements in SF-36 Physical Component Summary (PCS) and Mental Component Summary (MCS) scores, and these improvements were significantly greater at week 8 compared with the placebo group in UNITI-1 (MCS) and UNITI-2 (PCS, MCS and all subscores). These improvements in the PCS and MCS scores were maintained in STELARA treated patients in the IM-UNITI maintenance study through Week 44.

IBDQ

At Week 8 in UNITI-1 and UNITI-2, significant improvement from baseline in the inflammatory bowel disease questionnaire (IBDQ) total score and all subscales, was observed in the patients treated with STELARA compared to placebo. In both studies, a higher proportion of patients with clinically meaningful improvement in IBDQ total scores were observed in patients treated with STELARA compared to placebo. These improvements in the IBDQ total scores were maintained in STELARA treated patients in the IM-UNITI maintenance study through Week 44.

Long-term maintenance of health-related quality of life measures

Improvement in health-related quality of life as measured by IBDQ and SF-36 was generally maintained during the extension through week 252.

Clinical Efficacy – Ulcerative Colitis

The safety and efficacy of ustekinumab was assessed in two randomized, double-blind, placebo-controlled, multicenter studies in adult patients with moderately to severely active ulcerative colitis (Mayo score 6 to 12; Endoscopy subscore ≥ 2 based on central review of the endoscopy). The clinical development program consisted of one intravenous induction study (referred to as UNIFI-induction) with treatment of up to 16 weeks followed by a 44-week subcutaneous randomized withdrawal maintenance study (referred to as UNIFI-maintenance) representing at least 52 weeks of therapy.

Efficacy results presented for UNIFI-induction and UNIFI-maintenance were based on central review of endoscopies.

UNIFI-induction included 961 patients. The primary endpoint for the induction study was the proportion of patients in clinical remission (defined as a Mayo score ≤ 2 points, with no individual subscore > 1) at Week 8. Patients were randomized to receive a single intravenous administration of either the recommended tiered dose of approximately 6 mg/kg (see Table 1; Initial IV dosing of STELARA^a), a fixed dose of 130 mg ustekinumab, or placebo at Week 0.

Concomitant use of oral corticosteroids, immunomodulators, and aminosalicylates were permitted and 90% of patients continued to receive at least one of these medications. Enrolled patients had to have failed conventional therapy (corticosteroids or immunomodulators) or at least one biologic (a TNF α antagonist and/or vedolizumab). 49% of patients had failed conventional therapy, but not a biologic (of which 94% were biological-naïve). 51% of patients had failed or were intolerant to a biologic. Approximately 50% of the patients had failed at least 1 prior anti-TNF α therapy (of which 48% were primary non-responders) and 17% had failed at least 1 anti-TNF α therapy and vedolizumab.

In UNIFI-induction a significantly greater proportion of patients were in clinical response and remission in the ustekinumab treated group compared to placebo (Table 36). As early as Week 2, the earliest scheduled study visit, and at each visit thereafter, a higher proportion of ustekinumab patients had no rectal bleeding or achieved normal stool frequency (defined as a stool frequency subscore of 0 or 1) as compared with placebo patients. Significant differences in partial Mayo score and symptomatic remission were observed between ustekinumab and placebo as early as Week 2. Efficacy was higher in the tiered dose group (6 mg/kg) compared to the 130 mg dose group in select endpoints, and tiered dosing is therefore the recommended intravenous induction dose.

Table 36: Summary of Key Efficacy Measures in UNIFI-Induction (Week 8)

| Endpoint | Placebo N = 319 | | Ustekinumab [†] N = 322 | |
|--|--------------------|-----|-------------------------------------|------------------|
| | N | % | N | % |
| Clinical Remission* | 17 | 5% | 50 | 16% ^a |
| <i>Biologic-naïve[‡]</i> | 15/151 | 10% | 27/147 | 18% |
| <i>Not biologic failure</i> | 15/158 | 9% | 29/156 | 19% |
| <i>Prior biological failure</i> | 2/161 | 1% | 21/166 | 13% |
| Clinical Response[§] | 100 | 31% | 199 | 62% ^a |
| <i>Biologic-naïve[‡]</i> | 54/151 | 36% | 98/147 | 67% |
| <i>Not biologic failure</i> | 56/158 | 35% | 104/156 | 67% |
| <i>Prior biological failure</i> | 44/161 | 27% | 95/166 | 57% |
| Endoscopic Healing[¶] | 44 | 14% | 87 | 27% ^a |
| <i>Biologic-naïve[‡]</i> | 32/151 | 21% | 49/147 | 33% |
| <i>Not biologic failure</i> | 33/158 | 21% | 52/156 | 33% |
| <i>Prior biologic failure</i> | 11/161 | 7% | 35/166 | 21% |
| Histo-Endoscopic Mucosal Healing[‡] | 28/316 | 9% | 58/315 | 18% ^a |
| <i>Biologic-naïve[‡]</i> | 21/148 | 14% | 33/140 | 24% |
| <i>Not biologic failure</i> | 22/155 | 14% | 36/149 | 24% |
| <i>Prior biological failure</i> | 6/161 | 4% | 22/166 | 13% |
| Symptomatic Remission[¶] | 72 | 23% | 144 | 45% ^b |
| Combined Symptomatic Remission and Endoscopic Healing[‡] | 25 | 8% | 67 | 21% ^b |

[†] Infusion dose of ustekinumab using the weight-based dosage regimen specified in Table 1.

[‡] An additional 7 patients on placebo and 9 patients on ustekinumab (6mg/kg) had been exposed to, but had not failed, biologics.

^{*}Clinical remission is defined as Mayo score ≤2 points, with no individual subscore > 1.

[§] Clinical response is defined as a decrease from baseline in the Mayo score by ≥30% and ≥3 points, with either a decrease from baseline in the rectal bleeding subscore ≥1 or a rectal bleeding subscore of 0 or 1.

[€] Endoscopic healing is defined as a Mayo endoscopic subscore of 0 or 1 determined by central review of the endoscopy.

[‡] Histo-endoscopic mucosal healing is defined as combined endoscopic healing (Mayo endoscopy subscore of 0 or 1) and histologic healing of the colon tissue (neutrophil infiltration in <5% of crypts, no crypt destruction, and no erosions, ulcerations, or granulation tissue).

[€] Symptomatic remission is defined as a stool frequency subscore of 0 or 1 and a rectal bleeding subscore of 0.

[‡] Combined symptomatic remission and endoscopic healing is defined as remission based on a stool frequency subscore of 0 or 1, a rectal bleeding subscore of 0, and an endoscopy subscore of 0 or 1.

^a p < 0.001

^b Nominally significant (p < 0.001)

UNIFI-maintenance evaluated 523 patients who achieved clinical response with single IV administration of ustekinumab in UNIFI-induction. Patients were randomized to receive a subcutaneous maintenance regimen of either 90 mg ustekinumab every 8 weeks, 90 mg ustekinumab every 12 weeks or placebo for 44 weeks.

Significantly greater proportions of patients were in clinical remission at Week 44 and maintained clinical response through Week 44 in both ustekinumab treated groups compared to the placebo group (see Table 37).

| Table 37: Summary of Key Efficacy Measures in UNIFI-Maintenance (Week 44; 52 weeks from initiation of the induction dose) | | | |
|--|----------------------------|---|--|
| | Placebo* N = 175 | Ustekinumab 90 mg every 8 Weeks N = 176 | Ustekinumab 90 mg every 12 Weeks N = 172 |
| Clinical Remission** | 24% | 44% ^a | 38% ^b |
| <i>Biologic-naïve[‡]</i> | 32% (27/84) | 51% (40/79) | 47% (45/95) |
| <i>Not biologic failure</i> | 31% (27/87) | 48% (41/85) | 49% (50/102) |
| <i>Prior biologic failure</i> | 17% (15/88) | 40% (36/91) | 23% (16/70) |
| Maintenance of Clinical Response through Week 44[§] | 45% | 71% ^a | 68% ^a |
| <i>Biologic-naïve[‡]</i> | 52% (44/84) | 77% (61/79) | 77% (73/95) |
| <i>Not biologic failure</i> | 51% (44/87) | 78% (66/85) | 76% (78/102) |
| <i>Prior biologic failure</i> | 39% (34/88) | 65% (59/91) | 56% (39/70) |
| Corticosteroids Free Clinical Remission[‡] | 23% | 42% ^a | 38% ^b |
| <i>Biologic-naïve[‡]</i> | 32% (27/84) | 49% (39/79) | 46% (44/95) |
| <i>Not biologic failure</i> | 31% (27/87) | 47% (40/85) | 48% (49/102) |
| <i>Prior biologic failure</i> | 16% (14/88) | 37% (34/91) | 23% (16/70) |
| Endoscopic Healing at Week 44[†] | 29% | 51% ^a | 44% ^b |
| <i>Biologic-naïve[‡]</i> | 36% (30/84) | 58% (46/79) | 55% (52/95) |
| <i>Not biologic failure</i> | 34% (30/87) | 58% (49/85) | 56% (57/102) |
| <i>Prior biologic failure</i> | 23% (20/88) | 45% (41/91) | 26% (18/70) |
| Maintenance of Clinical Remission through Week 44[€] | 38% | 58% | 65% ^b |

| | | | |
|---|---------------|------------------|------------------|
| <i>Biologic-naïve^a</i> | 36% (9/25) | 75% (12/16) | 70% (21/30) |
| <i>Not biologic failure</i> | 36% (9/25) | 67% (12/18) | 72% (23/32) |
| <i>Prior biologic failure</i> | 40% (8/20) | 50% (10/20) | 38% (3/8) |
| Durable Partial Mayo Remission through Week 44 | 35% | 57% ^c | 48% ^c |
| Symptomatic Remission at Week 44[£] | 45% | 68% ^c | 62% ^c |
| Combined Symptomatic Remission and Endoscopic Healing at Week 44[€] | 28% | 48% ^c | 41% ^c |

* The placebo group consisted of patients who were in response to ustekinumab IV and were randomized to receive placebo at the start of maintenance therapy.

**Clinical remission is defined as Mayo score ≤ 2 points, with no individual subscore > 1 .

§ Clinical response is defined as a decrease from baseline in the Mayo score by $\geq 30\%$ and ≥ 3 points, with either a decrease from baseline in the rectal bleeding subscore ≥ 1 or a rectal bleeding subscore of 0 or 1.

^a An additional 3 patients on placebo and 6 patients on Q8W, 7 patients on Q12W ustekinumab had been exposed to, but had not failed, biologics

[†] Endoscopic healing is defined as a Mayo endoscopic subscore of 0 or 1 determined by central review of the endoscopy

[‡] Corticosteroid-free clinical remission is defined as patients in clinical remission and not receiving corticosteroids at Week 44.

[£] Maintenance of clinical remission is defined as patients in clinical remission at maintenance baseline through Week 44 among patients in clinical remission at maintenance baseline.

^{||} Durable partial Mayo remission is defined as partial Mayo remission (i.e. a partial Mayo score of ≤ 2) at $\geq 80\%$ of all visits prior to Week 44 and in partial Mayo remission at last visit (Week 44).

[€] Symptomatic remission is defined as a stool frequency subscore of 0 or 1 and a rectal bleeding subscore of 0.

[£] Combined symptomatic remission and endoscopic healing is defined as a stool frequency subscore of 0 or 1, a rectal bleeding subscore of 0, and an endoscopy subscore of 0 or 1.

^a p< 0.001

^b p<0.05

^c Nominally significant (p<0.05)

The beneficial effect of ustekinumab on clinical response, mucosal healing and clinical remission was observed in induction and in maintenance both in patients who failed conventional therapy but not a biologic therapy, as well as in those who had failed at least one prior TNF α antagonist therapy, and/or vedolizumab including in patients with a primary non-response to TNF α antagonist therapy.

Delayed Responders to Ustekinumab Induction

Ustekinumab treated patients who were not in response at Week 8 of UNIFI-induction received an administration of 90 mg SC ustekinumab at Week 8 (36% of patients). Of those patients, 9% of patients who were initially randomized to the recommended induction dose achieved clinical remission and 58% achieved clinical response at Week 16. When combining the delayed responders with the initial responders, 80% of subjects randomized to the recommended induction dose in UNIFI-I achieved clinical response and 18% achieved clinical remission within 16 weeks after initiating treatment with ustekinumab.

Patients who were not in clinical response to ustekinumab induction at Week 8 of the UNIFI-induction study but were in response at Week 16 (157 patients) entered in the non-randomized portion of UNIFI-maintenance and continued to receive maintenance dosing every 8 weeks; among these patients, a majority (62%) maintained response and 30% achieved remission at Week 44.

Long-Term Maintenance

In UNIFI (UCO3001), patients who completed the study through week 44 were eligible to continue treatment in a study extension. Among 400 patients who entered in the study extension and were treated with ustekinumab, symptomatic remission was generally maintained through week 200 for patients who failed conventional therapy (but not a biologic therapy) and those who failed biologic therapy, including those who failed both anti-TNF and vedolizumab. Of those patients who were assessed using the full Mayo score at week 200, mucosal healing and clinical remission were generally maintained.

The safety analysis including 457 patients (1289.9 person-years) followed up to 220 weeks showed a safety profile between week 44 and 220 that was comparable with that observed up to week 44.

No new safety concerns were identified in this study extension with up to 4 years of treatment in patients with ulcerative colitis.

Endoscopic Normalization

Normalization of endoscopic appearance of the mucosa was defined as a Mayo endoscopic subscore of 0 and was observed as early as Week 8 of UNIFI-induction. At Week 44 of UNIFI-maintenance, it was achieved in 24% and 29% of patients treated with ustekinumab every 12 or 8 weeks, respectively, as compared to 18% of patients in the placebo group.

Histologic & Histo-Endoscopic Mucosal Healing

Histologic healing (defined as neutrophil infiltration in <5% of crypts, no crypt destruction, and no erosions, ulcerations, or granulation tissue) was assessed at Week 8 of UNIFI-induction and Week 44 of UNIFI-maintenance. At Week 8, after a single intravenous induction dose, significantly greater proportions of patients in the recommended dose group achieved histologic healing (36%) compared with patients in the placebo group (22%). At Week 44 maintenance of this effect was maintained with significantly more patients in histologic healing in the every 12 week (54%) and every 8 week (59%) ustekinumab groups as compared to placebo (33%).

A combined endpoint of histo-endoscopic mucosal healing defined as subjects having both mucosal healing and histologic healing was evaluated at week 8 of UNIFI-induction and Week 44 of UNIFI-maintenance. Patients receiving ustekinumab at the recommended dose showed significant improvements on the histo-endoscopic mucosal healing endpoint at Week 8 in the ustekinumab group (18%) as compared to the placebo group (9%). At Week 44, maintenance of this effect was observed with significantly more patients in histo-endoscopic mucosal healing in the every 12 week (39%) and every 8 week (46%) ustekinumab groups as compared to placebo (24%).

Health-related quality of life

Health-related quality of life was assessed by Inflammatory Bowel Disease Questionnaire (IBDQ), SF-36 and EuroQoL-5D (EQ-5D) questionnaires. At Week 8 of UNIFI-induction, patients receiving ustekinumab showed significantly greater and clinically meaningful improvements on IBDQ total score, EQ-5D and EQ-5D VAS, and SF-36 Mental Component Summary Score and SF-36 Physical Component Summary Score when compared to placebo. These improvements were maintained in ustekinumab-treated patients in UNIFI-maintenance through Week 44.

Patients receiving ustekinumab experienced significantly more improvements in work productivity as assessed by greater reductions in overall work impairment and in activity impairment as assessed by the WPAI-GH questionnaire than patients receiving placebo.

Long-term maintenance of health-related quality of life measures

Improvement in health-related quality of life as measured by IBDQ and SF-36 was generally maintained during the extension through week 200.

Hospitalizations and Ulcerative Colitis related surgeries

Through Week 8 of UNIFI-induction, the proportions of subjects with ulcerative colitis disease related hospitalizations were significantly lower for subjects in the ustekinumab recommended dose group (1.6%, 5/322) compared with subjects in the placebo group (4.4%, 14/319) and no subjects underwent ulcerative colitis disease related surgeries in subjects receiving ustekinumab at the recommended induction dose compared to 0.6% (2/319) subjects in the placebo group.

Through Week 44 of UNIFI-maintenance, a significantly lower number of ulcerative colitis disease related hospitalizations was observed in subjects in the combined ustekinumab group (2.0%, 7/348) as compared with subjects in the placebo group (5.7%, 10/175). A numerically lower number of subjects in the ustekinumab group (0.6%, 2/348) underwent ulcerative colitis disease related surgeries compared with subjects in the placebo group (1.7%, 3/175) through Week 44.

Pharmacokinetic Properties

Absorption

The median time to reach the maximum serum concentration (t_{max}) was 8.5 days after a single 90 mg subcutaneous administration in healthy subjects. The median t_{max} values of ustekinumab following a single subcutaneous administration of either 45 mg or 90 mg in patients with psoriasis were comparable to that observed in healthy subjects.

The absolute bioavailability of ustekinumab following a single subcutaneous administration was estimated to be 57.2% in patients with psoriasis.

Following the recommended intravenous induction dose, median peak serum ustekinumab concentration was 126.1 mcg/mL in patients with Crohn's disease, and 127.0 mcg/mL in patients with ulcerative colitis.

Distribution

Median volume of distribution during the terminal phase (V_z) following a single intravenous administration to patients with psoriasis ranged from 57 to 83 mL/kg.

In a population pharmacokinetic analysis of ustekinumab, the volume of distribution at steady-state was 4.62 L in patients with Crohn's disease and 4.44 L in patients with ulcerative colitis.

Metabolism

The exact metabolic pathway for ustekinumab is unknown.

Elimination

Median systemic clearance (CL) following a single intravenous administration to patients with psoriasis ranged from 1.99 to 2.34 mL/day/kg.

Median half-life ($t_{1/2}$) of ustekinumab was approximately 3 weeks in patients with ulcerative colitis, Crohn's disease, psoriasis and/or psoriatic arthritis, ranging from 15 to 32 days across all psoriasis and psoriatic arthritis studies.

In a population pharmacokinetic analysis of ustekinumab, the clearance was 0.19 L/day while the half-life was approximately 19 days in patients with Crohn's disease and ulcerative colitis.

Dose Linearity

The systemic exposure of ustekinumab (C_{max} and AUC) increased in an approximately dose-proportional manner after a single intravenous administration at doses ranging from 0.09 mg/kg to 4.5 mg/kg or following a single subcutaneous administration at doses ranging from approximately 24 mg to 240 mg in patients with psoriasis.

Single Dose vs. Multiple Doses

Serum concentration-time profiles of ustekinumab were generally predictable after single or multiple subcutaneous dose administrations. In patients with psoriasis, steady-state serum concentrations of ustekinumab were achieved by Week 28 after initial subcutaneous doses at Weeks 0 and 4, followed by doses every 12 weeks. The median steady-state trough concentration ranged from 0.21 mcg/mL to 0.26 mcg/mL (45 mg) and from 0.47 mcg/mL to 0.49 mcg/mL (90 mg).

Following the recommended IV induction dose, median peak serum ustekinumab concentration was 126.1 mcg/mL in patients with Crohn's disease and 127.0 mcg/mL in patient with ulcerative colitis. Starting at Week 8, subcutaneous maintenance dosing of 90 mg ustekinumab was administered every 8 or 12 weeks. Steady state ustekinumab concentration was achieved by the start of the second maintenance dose. There was no apparent accumulation in serum ustekinumab concentration over time when given subcutaneously every 8 or 12 weeks.

Following subcutaneous maintenance dosing of 90 mg ustekinumab every 8 weeks, median steady-state trough concentrations ranged from 1.97 mcg/mL to 2.24 mcg/mL in patients with Crohn's disease and 2.69 mcg/mL to 3.09 mcg/mL in patients with ulcerative colitis. Following subcutaneous maintenance dosing of 90 mg ustekinumab every 12 weeks, median steady state trough concentrations ranged from 0.61 mcg/mL to 0.76 mcg/mL in patients with Crohn's disease and 0.92 mcg/mL to 1.19 mcg/mL in patients with ulcerative colitis. The steady-state trough ustekinumab levels resulting from 90 mg ustekinumab every 8 weeks were associated with higher clinical remission rates as compared to the steady-state trough levels following 90 mg every 12 weeks.

Dosing Frequency Adjustment

In patients with Crohn's disease and ulcerative colitis, based on observed data and population PK analyses, randomized subjects who lost response to treatment had lower serum ustekinumab concentrations over time compared with subjects who did not lose response. In Crohn's disease, dose adjustment from 90 mg every 12 weeks to 90 mg every 8 weeks was associated with an increase in trough serum ustekinumab concentrations and an accompanying increase in efficacy. In ulcerative colitis, population PK model based simulations demonstrated that adjusting dosing from 90 mg every 12 weeks to every 8 weeks would be expected to result in a 3-fold increase in steady-state trough ustekinumab concentrations. Additionally on the basis of clinical trial data in patients with ulcerative colitis, a positive exposure-response relationship was established between trough concentrations and clinical response, clinical remission, and mucosal healing.

Impact of Weight on Pharmacokinetics

Serum ustekinumab concentrations were affected by weight in patients with psoriasis and/or psoriatic arthritis. Within each dose (45 mg or 90 mg), patients of higher weight (> 100 kg) had lower median serum ustekinumab concentrations compared with those in patients of lower weight (≤ 100 kg). However, across doses, the median trough serum concentrations of ustekinumab in patients with higher weight (> 100 kg) in the 90 mg group were comparable to those in patients with lower weight (≤ 100 kg) in the 45 mg group.

Population Pharmacokinetic Analysis

In a population pharmacokinetic analysis using data from patients with psoriasis, the apparent clearance (CL/F) and apparent volume of distribution (V/F) were 0.465 L/d and 15.7 L, respectively, and the t_{1/2} was approximately 3 weeks in patients with psoriasis. The CL/F of ustekinumab was not impacted by sex, age, or race. The CL/F was impacted by body weight, with a trend toward higher CL/F in patients with higher body weight. The median CL/F in patients with weight > 100 kg was approximately 55% higher compared with patients with weight ≤ 100 kg. The median V/F in patients with weight > 100 kg was approximately 37% higher as compared with patients with weight ≤ 100 kg. Similar results were obtained from a confirmatory population pharmacokinetic analysis using data from patients with psoriatic arthritis.

In the population pharmacokinetic analysis using data from patients with psoriasis, the effect of comorbidities (past and current history of diabetes, hypertension, and hyperlipidemia) on pharmacokinetics of ustekinumab was evaluated. The pharmacokinetics of ustekinumab were impacted by the comorbidity of diabetes, with a trend towards higher CL/F in patients with diabetes. The mean CL/F in patients with diabetes was approximately 29% higher compared with patients without diabetes.

Population pharmacokinetic analyses showed that there was a trend towards a higher clearance of ustekinumab in patients with positive immune response.

No specific drug-drug interaction studies have been conducted in healthy subjects or patients with psoriasis, psoriatic arthritis, Crohn's disease or ulcerative colitis.

In the population pharmacokinetic analysis, the effect of the most frequently used concomitant medications in patients with psoriasis (including paracetamol/acetaminophen, ibuprofen, acetylsalicylic acid, metformin, atorvastatin, naproxen, levothyroxine, hydrochlorothiazide, and influenza vaccine) on pharmacokinetics of ustekinumab was explored and none of the concomitant medications exerted significant impact. The pharmacokinetics of ustekinumab was not impacted by the prior use of MTX, cyclosporine, or other biological therapeutics for the treatment of psoriasis. The pharmacokinetics of ustekinumab was not impacted by concomitant use of NSAIDs or prior exposure to anti-TNF α agents in patients with psoriatic arthritis; or by the use of MTX, and oral corticosteroids, 6-MP, AZA in patients with psoriatic arthritis or Crohn's disease, or by prior exposure to biologics (i.e. anti-TNF α agents and/or vedolizumab) in patients with ulcerative colitis.

The effects of IL-12 or IL-23 on the regulation of CYP450 enzymes were evaluated in an *in vitro* study using human hepatocytes, which showed that IL-12 and/or IL-23 at levels of 10 ng/mL did not alter human CYP450 enzyme activities (CYP1A2, 2B6, 2C9, 2C19, 2D6, or 3A4 (see *Drug Interactions*).

No clinically significant changes in exposure of caffeine (CYP1A2 substrate), warfarin (CYP2C9 substrate), omeprazole (CYP2C19 substrate), dextromethorphan (CYP2D6 substrate), or midazolam (CYP3A substrate) were observed when used concomitantly with ustekinumab at the approved recommended dosing in subjects with Crohn's disease (the same dose as ulcerative colitis and higher than approved recommended dosing for plaque psoriasis and psoriatic arthritis) (see *Interactions*).

Special populations

Pediatrics (17 years of age and younger)

The pharmacokinetics of ustekinumab in pediatric psoriasis patients, 6 to 17 years of age, treated with the recommended dose was generally comparable to that in the adult psoriasis population. No pharmacokinetic data are available in pediatric patients with Crohn's disease or ulcerative colitis

Elderly (65 years of age and older)

No specific studies have been conducted in elderly patients. The population pharmacokinetic analysis indicated there were no apparent changes in CL/F and V/F estimates in patients ≥ 65 years.

Renal impairment

No pharmacokinetic data are available in patients with renal insufficiency.

Hepatic impairment

No pharmacokinetic data are available in patients with impaired hepatic function.

Other populations

The pharmacokinetics of ustekinumab were generally comparable between Asian and non-Asian patients with psoriasis, Crohn's disease, or ulcerative colitis.

The pharmacokinetics of ustekinumab were not impacted by the use of tobacco or alcohol.

NON-CLINICAL INFORMATION

In repeated-dose toxicity studies in cynomolgus monkeys, ustekinumab was well-tolerated following IV doses up to 45 mg/kg/week for up to 1 month and following twice-weekly SC doses up to 45 mg/kg for 6 months. There were no ustekinumab -related findings in the immunotoxicity and cardiovascular safety pharmacology evaluations. In histopathology evaluations there were no preneoplastic changes observed.

There were no adverse effects in monkeys at exposures that were 179-fold higher than the peak serum concentration in humans following 90 mg weekly subcutaneous injection and 29-fold higher than the peak serum concentration in humans following 6 mg/kg IV administration.

Carcinogenicity and Mutagenicity

Carcinogenicity studies were not performed with ustekinumab due to the lack of appropriate models for an antibody with no cross-reactivity to rodent IL-12/23 p40.

Reproductive Toxicology

Three developmental toxicity studies were conducted in cynomolgus monkeys. No ustekinumab -related maternal toxicity, abortions, still-births, embryotoxicity, developmental delays, malformations or birth defects were observed at doses up to 45 mg/kg following weekly or twice weekly administration of ustekinumab via the IV or SC routes, respectively. In neonates born from pregnant monkeys treated with ustekinumab no adverse effects on growth or functional development were observed and no deficits were observed in immunotoxicity evaluations. In a male fertility study in cynomolgus monkeys no ustekinumab -related effects on mating behavior, sperm parameters, or serum concentrations of male hormones were observed following twice weekly subcutaneous administration of ustekinumab at doses up to 45 mg/kg.

A female fertility toxicity study was conducted in mice using an analogous antibody that binds to and inhibits IL-12 and IL-23 activity in mice. Twice weekly subcutaneous administration of the anti-mouse IL-12/23 antibody was well tolerated at doses up to 50 mg/kg and no adverse effects on female fertility parameters were observed.

PHARMACEUTICAL PARTICULARS

List of Excipients

45 mg or 90 mg Pre-filled syringe/vial

L-histidine
L-histidine monohydrochloride monohydrate
Polysorbate 80
Sucrose
Water for injection

130 mg vial

EDTA disodium salt dihydrate
L-histidine
L-histidine hydrochloride monohydrate
L-methionine
Polysorbate 80
Sucrose
Water for injection

Incompatibilities

Not applicable

Shelf Life

45 mg vial: 24 months
Prefilled Syringe: 36 months
130 mg vial: 36 months

Special Precautions for Storage

- Store in a refrigerator
 - 2°C to 8°C
- Store in original carton until time of use
- Protect from light.

- Do not freeze
- Do not shake

Keep out of reach of children

If needed, individual STELARA pre-filled syringes may be stored at room temperature up to 30°C for a maximum single period of up to 30 days in the original carton with protection from light. Record the date when the pre-filled syringe is first removed from the refrigerator and the new expiry date on the carton in the spaces provided. The new expiry date must not exceed the original expiry date printed on the carton. Once a syringe has been stored at room temperature, it should not be returned to the refrigerator. Discard the syringe if not used within 30 days at room temperature storage.

Nature and Contents of Container

For subcutaneous injection

STELARA is supplied as a sterile solution in a single-use (Type 1) glass vial. The vial is stoppered with a coated stopper.

STELARA is also supplied as a single-use, sterile solution in a Type 1 glass **pre-filled** syringe with a fixed 27G, half-inch needle and needle cover. The needle cover is manufactured using a dry natural rubber (a derivative of latex) (see *Warnings and Precautions*). The syringe is fitted with a passive safety guard.

The solution is clear to slightly opalescent, colorless to light yellow with a pH of approximately 6.0. Each mL of STELARA contains 90 mg of ustekinumab, 1.0 mg L-histidine and L-histidine hydrochloride, 76 mg sucrose, 0.04 mg polysorbate 80, and Water for Injection, USP. STELARA does not contain preservatives.

There are two strengths of STELARA available: 45 mg of ustekinumab in 0.5 mL or 90 mg of ustekinumab in 1.0 mL. STELARA is available in the following packaging presentation:

- 1 single-use pre-filled syringe
- 1 single use vial

For intravenous infusion only

STELARA 130 mg vial is supplied as a sterile solution in a single-use (Type 1) glass vial. The vial is stoppered with a coated stopper. The solution is clear, colorless to light yellow with a pH of approximately 6.0. Each mL of STELARA contains 5.0 mg of ustekinumab, 0.8 mg L-histidine, 1.1 mg L-histidine hydrochloride monohydrate, 85 mg sucrose, 0.40 mg polysorbate 80, 0.40 mg L-methionine, and 0.02 mg EDTA disodium salt dihydrate. STELARA does not contain preservatives. STELARA is available for intravenous infusion in one strength, 130 mg in 26 mL, and packaged as 1 single use vial.

Instructions for Use and Handling and Disposal

Following administration of STELARA, discard any unused portion. The syringe should be disposed of with accepted medical practices for used syringes. The syringe, needle and vial must never be re-used.

Instructions for dilution of STELARA 130 mg for IV infusion (Crohn's disease)

STELARA 130 mg solution must be diluted and prepared for IV infusion by a healthcare professional using aseptic technique.

1. Calculate the dose and the number of STELARA vials needed based on patient's body weight (see Table 1). Each 26 mL vial of STELARA contains 130 mg of ustekinumab.
2. Withdraw and then discard a volume of the 0.9% w/v sodium chloride solution from the 250 mL infusion bag equal to the volume of STELARA to be added. (discard 26 mL sodium chloride for each vial of STELARA needed, for 2 vials-discard 52 mL, for 3 vials- discard 78 mL, for 4 vials- discard 104 mL). Alternatively, a 250 mL infusion bag containing 0.45% w/v sodium chloride solution may be used.
3. Withdraw 26 mL of STELARA from each vial needed and add it to the 250 mL infusion bag. The final volume in the infusion bag should be 250 mL. Gently mix.
4. Visually inspect the diluted solution before administration. Do not use if visibly opaque particles, discolouration or foreign particles are observed.
5. Administer the diluted solution over a period of at least one hour. Once diluted, the infusion should be completed within eight hours of the dilution in the infusion bag.
6. Use only an infusion set with an in-line, sterile, non-pyrogenic, low protein-binding filter (pore size 0.2 micrometer).
7. Do not infuse STELARA concomitantly in the same intravenous line with other agents.
8. Each vial is for single use only and any unused medicinal product should be disposed of in accordance with local requirements.

Storage

If necessary, the diluted infusion solution may be stored at room temperature. The infusion should be completed within 8 hours of the dilution in the infusion bag. Do not freeze. Discard any unused portion of the infusion solution.

HOW SUPPLIED

- Stelara 45 mg in 0.5 ml Pre-Filled Syringe
Box, 1 Pre-Filled Syringe @ 0.5mL
Reg. No.: DKI2060001643A1
- Stelara 90 mg in 1.0 ml Pre-Filled Syringe
Box, 1 Pre-Filled Syringe @ 1 mL
Reg. No.: DKI2360001643C1
- Stelara 45 mg in 0.5 mL vial
Box, 1 vial @ 45 mg/0.5 mL
Reg. No.: DKI2060001643A2
- Stelara 130 mg in 26mL vial
Box, 1 vial @ 130 mg/26 mL
Reg. No.: DKI2160001643B1

DATE OF FIRST AUTHORISATION

- **Stelara 45 mg in 0.5 ml Pre-Filled Syringe**
21 January 2014
- **Stelara 90 mg in 1.0 ml Pre-Filled Syringe**
20 August 2023
- **Stelara 45 mg in 0.5 mL vial**
1 May 2020
- **Stelara 130 mg in 26mL vial**
22 July 2021

HARUS DENGAN RESEP DOKTER

Manufactured, packed and released by Cilag AG, Hochstrasse 201, 8200, Schaffhausen, Switzerland

Registered by PT Integrated Healthcare Indonesia, Jakarta – Indonesia

For adverse event and product quality complaint please contact drugsafety@jacid.jni.com or Phone (021) 2935 3935

Based on CCDS 17-Oct-22 ver.50 + CCDS 1-Jun-23 ver.51 + CCDS v.52 4Mar24

INFORMASI PRODUK UNTUK PASIEN

Stelara (ustekinumab)

Tinjau informasi pasien ini sebelum Anda mulai menggunakan STELARA dan setiap kali Anda mendapatkannya lagi. Ada kemungkinan terdapat informasi baru. Informasi ini tidak menggantikan percakapan dengan penyedia layanan kesehatan Anda mengenai kondisi atau perawatan medis Anda. Penting untuk tetap berada di bawah perawatan penyedia layanan kesehatan selama menggunakan STELARA.

Apa informasi terpenting yang harus saya ketahui tentang STELARA?

STELARA adalah obat yang dapat mengurangi aktivitas sistem kekebalan tubuh Anda. Ini dapat meningkatkan peluang Anda mendapatkan efek samping yang serius termasuk:

Infeksi serius:

- STELARA dapat menurunkan kemampuan Anda untuk melawan infeksi. Beberapa infeksi dapat menjadi serius dan menyebabkan rawat inap. Jika Anda memiliki infeksi atau mengalami luka terbuka, beri tahu penyedia layanan kesehatan Anda sebelum Anda mulai menggunakan STELARA. Jika Anda mendapatkan infeksi, mengalami tanda-tanda infeksi seperti demam, merasa sangat lelah, batuk, gejala seperti flu, atau kulit atau luka yang hangat, merah, atau nyeri pada tubuh Anda, segera beri tahu penyedia layanan kesehatan Anda. Ini mungkin tanda-tanda infeksi seperti infeksi dada, atau infeksi kulit atau herpes zoster yang dapat menyebabkan komplikasi serius.
- Dokter Anda akan memeriksa Anda untuk Tuberkulosis (TB) dan melakukan tes untuk mengetahui apakah Anda menderita TB. Jika dokter Anda merasa bahwa Anda berisiko untuk TB, Anda akan diobati dengan obat untuk TB sebelum Anda memulai pengobatan dengan STELARA dan selama perawatan dengan STELARA.

Kanker

- Banyak obat seperti STELARA yang dapat menurunkan aktivitas sistem kekebalan tubuh, dapat meningkatkan risiko kanker. Beri tahu dokter Anda jika Anda pernah menderita kanker jenis apa pun.

Lihat juga, "Apa efek samping yang mungkin dari STELARA?" di bawah ini.

Apakah STELARA?

STELARA adalah obat resep yang telah disetujui untuk pengobatan:

Psoriasis plak pada dewasa

STELARA diindikasikan untuk pengobatan psoriasis plak sedang hingga berat pada orang dewasa yang gagal merespons, atau yang memiliki kontraindikasi, atau tidak toleran terhadap terapi sistemik lainnya termasuk siklosporin, metotreksat atau PUVA.

Psoriasis plak pada anak

Stelara, sebagai monoterapi, diindikasikan untuk pengobatan psoriasis plak sedang hingga berat pada pasien anak-anak dan remaja dari usia 6 tahun ke atas, yang merupakan kandidat untuk fototerapi atau pengobatan psoriasis sistemik (baik naif atau dengan riwayat pengobatan sebelumnya) atau memiliki psoriasis yang dianggap tidak terkontrol secara memadai dengan terapi topikal setelah dosis dan durasi terapi yang memadai.

Psoriatic Arthritis (PsA)

STELARA, sebagai monoterapi atau kombinasi dengan metotreksat (MTX), diindikasikan untuk pengobatan *psoriatic arthritis* sedang hingga berat pada pasien dewasa ketika respons terhadap terapi obat anti rematik (DMARD) nonbiologis sebelumnya tidak memadai,

- Mengurangi tanda dan gejala
- Meningkatkan fungsi fisik

- Menghambat perkembangan kerusakan struktural
- Memperbaiki enthesitis
- Memperbaiki psoriasis
- Meningkatkan kualitas hidup yang berhubungan dengan kesehatan pada orang dewasa dengan *psoriatic arthritis*.

Penyakit Crohn

Stelara diindikasikan untuk pengobatan pasien dewasa dengan penyakit Crohn sedang hingga berat yang memiliki respon yang tidak memadai, kehilangan respon, atau tidak toleran terhadap terapi konvensional atau antagonis TNF α atau memiliki kontraindikasi medis terhadap terapi tersebut.

Kolitis ulseratif

STELARA diindikasikan untuk pengobatan pasien dewasa dengan kolitis ulserativa sedang hingga berat yang memiliki respon yang tidak memadai, kehilangan respon, atau tidak toleran terhadap terapi konvensional atau terapi biologis atau memiliki kontraindikasi medis terhadap terapi tersebut.

STELARA memblokir aksi dua protein dalam tubuh Anda yang disebut interleukin 12 (IL-12) dan interleukin 23 (IL-23). IL-12 dan IL-23 dibuat oleh sistem kekebalan tubuh Anda. Pada orang dengan psoriasis, psoriasis arthritis atau penyakit Crohn, IL 12 dan IL-23 dapat menyebabkan sistem kekebalan tubuh mereka menyerang bagian tubuh normal yang sehat. STELARA dapat memblokir IL-12 dan IL-23 dari menyebabkan sistem kekebalan tubuh menyerang kulit, kuku, sendi atau saluran pencernaan.

Siapa yang tidak seharusnya menggunakan STELARA?

Anda seharusnya tidak mendapatkan STELARA jika Anda:

- memiliki reaksi alergi terhadap STELARA, atau salah satu bahan lain di STELARA. Lihat akhir label ini untuk melihat daftar lengkap pada produk STELARA.

Apa yang harus saya katakan kepada dokter saya sebelum memulai pengobatan dengan STELARA?

Dokter Anda akan menilai kesehatan Anda sebelum memulai setiap pengobatan.

Beri tahu dokter Anda tentang semua kondisi medis Anda sebelum setiap pengobatan, termasuk jika Anda:

- pernah mengalami reaksi alergi terhadap STELARA. Tanyakan kepada dokter Anda jika Anda tidak yakin.
- menderita infeksi apa pun walaupun infeksi yang sangat kecil.
- memiliki infeksi yang tidak akan hilang atau riwayat infeksi yang terus muncul.
- menderita TB (tuberkulosis), atau jika Anda baru saja berada di dekat siapa saja yang mungkin menderita TB.
- memiliki atau pernah menderita kanker jenis apa pun.
- memiliki lesi baru atau perubahan dalam area psoriasis atau pada kulit normal.
- baru-baru ini menerima atau dijadwalkan untuk menerima vaksin. Beri tahu dokter Anda jika ada orang di rumah Anda yang membutuhkan vaksin. Virus dalam beberapa vaksin dapat menyebar ke orang dengan sistem kekebalan yang melemah, dan dapat menyebabkan masalah serius.
- Baru menerima atau telah menerima 'suntikan alergi', terutama untuk reaksi alergi serius.
- hamil, berencana hamil, atau menyusui. **Konsultasikan kondisi Anda dengan dokter Anda sebelum Anda menggunakan obat ini.** STELARA **dapat** digunakan selama kehamilan jika diperlukan. STELARA dapat masuk ke dalam ASI Anda dalam jumlah yang sangat kecil. Wanita yang sedang menyusui harus berbicara dengan dokter mereka tentang apakah perlu menggunakan atau tidak menggunakan STELARA.
- alergi terhadap lateks

Beri tahu dokter Anda tentang semua obat yang Anda gunakan, termasuk obat resep dan non-resep, vitamin, dan suplemen herbal.

Ketahui obat-obatan yang Anda minum. Simpan daftar obat-obatan Anda dan tunjukkan ke dokter dan apoteker ketika Anda mendapatkan obat baru.

Bagaimana STELARA digunakan?

- Untuk pengobatan psoriasis dan psoriasis arthritis, STELARA diberikan dengan suntikan di bawah kulit.

- Untuk pengobatan penyakit Crohn atau kolitis ulserativa, dosis pertama STELARA diberikan melalui infus intravena, yang berarti bahwa obat tersebut akan diberikan kepada Anda melalui jarum yang diletakkan di pembuluh darah. Dosis ini diberikan oleh penyedia layanan kesehatan Anda. Setelah itu, STELARA diberikan melalui suntikan di bawah kulit.
- STELARA dimaksudkan untuk digunakan di bawah bimbingan dan pengawasan dokter Anda. Pada anak dan remaja usia 6 tahun atau lebih dengan psoriasis, disarankan pemakaian STELARA dilakukan oleh tenaga kesehatan. Atas penilaian dokter, Anda mungkin dapat memberikan STELARA kepada diri Anda sendiri, setelah pelatihan yang tepat dalam teknik injeksi menggunakan jenis jarum suntik yang tepat dan jumlah (volume) yang akan disuntikkan (lihat **“Petunjuk untuk Menyuntikkan STELARA di bawah kulit sendiri”**).
- Dokter Anda akan menentukan dosis STELARA yang tepat untuk Anda, jumlah untuk setiap suntikan dan seberapa sering Anda harus menerimanya. Pastikan untuk berdiskusi dengan dokter Anda kapan Anda akan menerima suntikan dan datang untuk semua tindak lanjut yang Anda jadwalkan.

Apa yang harus saya hindari saat menggunakan STELARA?

Anda tidak boleh menerima vaksin langsung saat menggunakan STELARA.

Jika Anda menggunakan STELARA saat hamil, beri tahu dokter bayi Anda tentang penggunaan STELARA Anda sebelum bayi menerima vaksin apa pun, termasuk vaksin langsung, seperti vaksin BCG (digunakan untuk mencegah tuberkulosis), vaksin rotavirus, atau vaksin hidup (*live vaccine*) lainnya.

Pemberian vaksin kepada bayi yang lahir dari ibu yang menggunakan Stelara saat hamil tidak dianjurkan sampai usia bayi enam bulan atau sampai kadar serum ustekinumab pada bayi tidak terdeteksi.

Apa efek samping yang mungkin dari STELARA?

Infeksi serius. (lihat **“Apa informasi paling penting yang harus saya ketahui tentang STELARA?”**).

Kanker. (lihat **“Apa informasi paling penting yang harus saya ketahui tentang STELARA?”**).

Efek samping serius lainnya dapat terjadi termasuk reaksi alergi. Tanda-tanda reaksi alergi yang serius mungkin termasuk ruam kulit, Bengkak pada wajah, bibir, mulut atau tenggorokan atau mengi, pusing, kesulitan menelan atau bernapas. Beri tahu dokter Anda atau dapatkan bantuan medis darurat segera jika Anda merasa mengalami alergi.

Dalam kasus yang jarang terjadi, gejala seperti batuk, sesak napas, dan demam juga bisa menjadi tanda reaksi alergi pada paru-paru terhadap STELARA.

Alergi terhadap lateks

Penutup jarum pada jarum suntik mengandung karet alam kering (suatu bentuk lateks). Ini dapat menyebabkan reaksi alergi pada orang yang sensitif terhadap lateks. Beri tahu dokter Anda jika Anda pernah mengalami reaksi alergi terhadap lateks dan reaksi alergi apa pun terhadap injeksi STELARA.

Efek samping yang paling umum dari STELARA adalah:

- Infeksi saluran pernapasan atas seperti flu biasa
- Infeksi pada hidung dan tenggorokan
- Pusing
- Sakit kepala
- Sakit tenggorokan
- Diare
- Mual
- Muntah
- Gatal
- Sakit punggung
- Nyeri otot
- Nyeri sendi
- Merasa sangat lelah

- Kemerahan pada kulit tempat injeksi diberikan
- Nyeri saat injeksi diberikan
- Infeksi sinus

Efek samping diatas tidak mencakup semua efek samping yang berkaitan dengan STELARA. Beri tahu dokter Anda tentang efek samping apa pun yang mengganggu Anda atau tidak hilang. Secara umum, efek samping STELARA pada anak dan remaja usia 6 tahun atau lebih sama dengan dengan efek samping pada pasien dewasa. Tanyakan kepada dokter atau apoteker Anda untuk informasi lebih lanjut.

Informasi umum tentang STELARA

Obat-obatan kadang-kadang diresepkan untuk tujuan yang tidak disebutkan dalam informasi produk untuk pasien. Jangan menggunakan STELARA tidak sesuai dengan yang diresepkan.

Bagaimana STELARA disimpan

Jika Anda menggunakan STELARA di rumah, penting untuk menyimpan produk di lemari es Anda meskipun tidak di kompartemen freezer. STELARA tidak boleh dibekukan. Simpan produk dalam kemasan karton asli agar terlindung dari cahaya sampai saat digunakan.

Jika dibutuhkan, setiap *pre-filled syringe* Stelara dapat disimpan pada suhu kamar hingga 30°C untuk jangka waktu tunggal maksimum hingga 30 hari dalam karton aslinya dengan perlindungan dari cahaya. Catat tanggal ketika *pre-filled syringe* dikeluarkan dari lemari es dan tanggal kedaluwarsa yang baru di tempat yang tersedia pada karton. Tanggal kedaluwarsa yang baru tidak boleh melebihi tanggal kedaluwarsa asli yang dicetak di karton. Setelah jarum suntik disimpan pada suhu kamar, sebaiknya tidak dikembalikan lagi ke lemari es. Buang jarum suntik jika tidak digunakan dalam kurun waktu 30 hari pada penyimpanan suhu kamar.

Apa kandungan dari STELARA?

Bahan aktif adalah ustekinumab.

STELARA 45 mg atau 90 mg untuk injeksi di bawah kulit: Bahan-bahan tambahan mengandung L histidin, L-histidin monohidroklorida monohidrat, polisorbat 80, sukrosa dan air untuk injeksi. Tidak ada bahan pengawet.

STELARA 130 mg untuk infus intravena: Bahan-bahan tambahan mengandung EDTA disodium salt dihydrate, L histidin, L-histidine hidroklorida monohidrat, L-metionin, polisorbat 80, sukrosa dan air untuk injeksi. Tidak ada bahan pengawet.

Informasi penting lainnya:

STELARA berupa cairan yang tidak berwarna hingga berwarna kuning cerah dan mungkin berisi beberapa partikel protein kecil yang tembus cahaya atau putih. Penampilan ini wajar untuk larutan yang mengandung protein.

STELARA tidak boleh digunakan:

- setelah tanggal kedaluwarsa pada label;
- jika segel rusak;
- jika cairannya berubah warna, keruh atau Anda dapat melihat partikel lain mengambang di dalamnya;
- jika Anda tahu, atau berpikir bahwa itu mungkin telah terkena suhu ekstrem (seperti tidak sengaja beku atau dipanaskan);

Selalu jauhkan obat dari jangkauan anak-anak.

Bagaimana Cara Membuang STELARA setelah pemakaian?

Jarum suntik bekas harus ditempatkan dalam wadah tahan tusukan, seperti wadah benda tajam. Buang wadah benda tajam Anda sesuai dengan peraturan yang berlaku. Botol kosong, tisu antiseptik, dan persediaan lain dapat ditempatkan di tempat sampah biasa. Jarum suntik, alat suntikan, dan botol vial tidak boleh digunakan kembali. Buang bagian STELARA yang tidak digunakan.

Petunjuk untuk Menyuntikkan STELARA di bawah kulit sendiri

Suntikan STELARA pertama Anda dapat diberikan oleh penyedia layanan kesehatan Anda. Pada anak dan remaja usia 6 tahun atau lebih, direkomendasikan untuk semua dosis STELARA diberikan oleh tenaga kesehatan. Namun, tenaga kesehatan Anda mungkin memutuskan bahwa Anda atau perawat Anda dapat mempelajari cara menyuntikkan STELARA di bawah kulit (secara subkutan) sendiri. **Sebelum Anda menyuntikkan sendiri**


STELARA, Anda harus dilatih oleh tenaga kesehatan profesional. Jika Anda belum dilatih, silakan hubungi penyedia layanan kesehatan Anda untuk menjadwalkan sesi pelatihan. Hubungi penyedia layanan kesehatan Anda jika Anda memiliki pertanyaan tentang cara menyuntikkan pada diri sendiri.

INSTRUKSI CARA PENYUNTIKAN STELARA MENGGUNAKAN “PRE-FILLED SYRINGE”

Untuk mengurangi risiko kecelakaan jarum yang tidak disengaja bagi pengguna, setiap pre-filled syringe dilengkapi dengan pelindung jarum yang secara otomatis diaktifkan untuk menutupi jarum setelah jarum suntik selesai digunakan.

Jangan mengocok STELARA sembarangan. Kocokan yang kuat dalam waktu lama dapat merusak produk. Jika produk telah dikocok dengan keras, jangan gunakan lagi.

LANGKAH 1: MEMPERSIAPKAN PENGGUNAAN PRE-FILLED SYRINGE



Keluarkan Syringe dari Kulkas

Periksa kemasannya untuk memastikan bahwa itu adalah dosis yang tepat, baik 45 mg atau 90 mg seperti yang ditentukan oleh penyedia layanan kesehatan Anda. Anak dengan berat badan 60kg atau lebih, dapat menggunakan pre-filled syringe.

Periksa Tanggal Kedaluwarsa

Buka kotak dan keluarkan prefilled syringe. Periksa tanggal kedaluwarsa pada prefilled syringe dan label kotak. Jika tanggal kedaluwarsa telah berlalu, jangan gunakan lagi.

Merakit Peralatan Tambahan

Pasang peralatan tambahan yang Anda perlukan untuk injeksi Anda. Ini termasuk tisu antiseptik, bola kapas atau kain kasa, dan wadah benda tajam untuk pembuangan jarum suntik.

Periksa larutan di dalam Jarum Suntik

Pegang prefilled syringe dengan jarum tertutup mengarah ke atas. Pastikan jarum suntik tidak rusak. Lihatlah larutan atau cairan di dalam jarum suntik untuk memastikan terlihat jelas sedikit opalescent dan tidak berwarna hingga agak kuning. **JANGAN** gunakan jika beku, berubah warna, berawan atau mengandung partikel besar dan hubungi penyedia layanan kesehatan Anda untuk mendapatkan bantuan.


JANGAN melepas penutup jarum dari prefilled syringe.

JANGAN pernah menarik kembali kepala pendorong.

LANGKAH 2: MEMILIH DAN MENYIAPKAN TEMPAT PENYUNTIKAN

Pilih tempat Penyuntikan*

Tempat penyuntikan yang baik adalah bagian atas paha dan sekitar perut (abdomen) tetapi sekitar 2 inci jauhnya dari pusar (pusar). Hindari, jika mungkin, kulit yang terkena psoriasis. Jika perawat Anda memberi Anda suntikan, mereka mungkin juga menggunakan lengan bagian atas.



*Areas in gray are recommended injection sites.


Siapkan tempat Penyuntikan

Cuci tangan Anda dengan sabun dan air hangat. Usap tempat penyuntikan dengan antiseptik. JANGAN menyentuh area ini lagi sebelum memberikan suntikan.

LANGKAH 3: PENYUNTIKAN OBAT

Lepaskan Penutup Jarum


Saat Anda siap menyuntikkan, ambil *prefilled syringe*, pegang tubuh jarum suntik dengan satu tangan dan tarik penutup jarum. Buang tutup jarum ke tempat sampah. Anda mungkin melihat gelembung udara kecil di *prefilled syringe*. Anda tidak perlu menghilangkan gelembung udara tersebut. Anda juga dapat melihat setetes cairan di ujung jarum - ini normal. Jangan menyentuh jarum atau membiarkannya menyentuh permukaan apa pun.



Catatan: Penutup jarum **TIDAK** boleh dilepas sampai Anda siap untuk menyuntikkan obat. Jangan menggunakan jarum suntik jika jarum terjatuh tanpa penutup jarum pada tempatnya. Jika Anda menjatuhkan jarum suntik tanpa penutup jarum, harap hubungi penyedia layanan kesehatan Anda untuk mendapatkan bantuan.

Suntikkan Obat

Jepit kulit yang dibersihkan dengan lembut di antara ibu jari dan jari telunjuk.




Jangan meremasnya.

Dorong jarum suntik ke kulit yang terjepit.

Dorong plunger dengan ibu jari sejauh mungkin untuk menyuntikkan semua cairan.
Dorong perlahan dan merata, jaga kulit terjepit.

Ketika plunger memenuhi ujung tabung jarum suntik, dan semua obat telah disuntikkan, lepaskan kulit yang terjepit dan lepaskan jarum dengan lembut. Setelah injeksi lengkap, pelindung jarum akan secara otomatis menjangkau jarum dan mengunci saat Anda melepaskan plunger.



LANGKAH 4: SETELAH PENYUNTIKAN

Buang Jarum Suntik Kosong

Segera buang jarum suntik kosong ke dalam wadah benda tajam. Untuk keselamatan dan kesehatan Anda dan untuk keselamatan orang lain, alat suntikan dan jarum suntik TIDAK BOLEH digunakan kembali. Buang wadah benda tajam sesuai dengan peraturan yang berlaku.



Gunakan Bola Kapas atau Kasa

Mungkin ada sejumlah kecil darah atau cairan di tempat suntikan, dimana hal ini adalah normal. Anda dapat menekan bola kapas atau kain kasa di atas tempat injeksi dan tahan selama 10 detik. Jangan menggosok tempat penyuntikan. Anda dapat menutup tempat penyuntikan dengan perban perekat kecil, jika perlu.

INSTRUKSI CARA PENYUNTIKKAN STELARA MENGGUNAKAN VIAL

Jangan mengocok STELARA sembarangan. Kocokan yang kuat dalam waktu lama dapat merusak produk. Jika produk dikocok dengan keras, jangan gunakan lagi. STELARA tidak boleh dicampur dengan cairan lain untuk injeksi.

LANGKAH 1: PERIKSA VIAL DAN ALAT-ALAT PELENGKAP

Keluarkan Vial dari Kulkas

Periksa paket untuk memastikan bahwa itu adalah dosis yang tepat. Jika dosis Anda 45 mg, Anda akan menerima satu botol 45 mg.

Anak-anak dengan berat kurang dari 60 kg memerlukan dosis lebih rendah dari 45 mg. Pastikan Anda tahu jumlah (volume) yang tepat dan jenis jarum suntik yang diperlukan untuk dosis ini. Jika Anda tidak tahu jumlah atau jenis jarum suntik yang diperlukan, hubungi tenaga kesehatan Anda untuk instruksi lebih lanjut.

Periksa Tanggal Kedaluwarsa


Buka kotak dan keluarkan vial. Periksa tanggal kedaluwarsa pada vial dan label kotak. Jika tanggal kedaluwarsa telah berlalu, jangan gunakan lagi.

Periksa larutan dalam Vial

Pastikan vial tidak rusak. Lihatlah larutan atau cairan dalam vial untuk memastikan bahwa larutan jelas, sedikit opalescent dan tidak berwarna hingga agak kuning. **JANGAN** gunakan jika beku, berubah warna, berawan atau mengandung partikel besar dan hubungi penyedia layanan kesehatan Anda untuk bantuan.

Mempersiapkan alat Tambahan


Siapkan alat tambahan yang Anda perlukan untuk injeksi Anda. Yaitu tisu antiseptik, bola kapas atau kain kasa, dan wadah benda tajam untuk pembuangan jarum suntik.



LANGKAH 2: MEMILIH DAN MENYIAPKAN TEMPAT PENYUNTIKAN

Pilih Tempat Penyuntikan *

Tempat penyuntikan yang baik adalah bagian atas paha dan sekitar perut (abdomen) sekitar 2 inci jauhnya dari pusar. Hindari, jika mungkin, kulit yang terkena psoriasis. Jika perawat Anda memberi Anda suntikan, mereka mungkin juga menggunakan lengan bagian atas.




*Areas in gray are recommended injection sites.

Siapkan tempat Penyuntikan

Cuci tangan Anda dengan sabun dan air hangat. Usap tempat penyuntikan dengan antiseptik. **JANGAN** menyentuh area ini lagi sebelum memberikan suntikan.

LANGKAH 3: MEMPERSIAPKAN OBAT


Lepaskan tutup dari bagian atas botol tetapi jangan lepaskan sumbatnya. Bersihkan sumbat dengan tisu antiseptik.



Lepaskan penutup jarum dari jarum suntik. Jangan menyentuh jarum atau membiarkan jarum menyentuh apa pun.

Letakkan vial pada permukaan yang rata dan dorong jarum suntik melalui penghenti karet.

Balikkan vial dan jarum suntik di bagian bawah vial.




Untuk pasien dewasa dan anak-anak (anak dan remaja) usia 6 tahun atau lebih, dengan berat badan 60 kg atau lebih, tarik alat suntik untuk mengisi jarum suntik dengan jumlah (volume) cairan yang ditentukan oleh tenaga kesehatan Anda (0,5 mL atau 1,0 mL).

Untuk anak dan remaja usia 6 tahun atau lebih dengan berat badan kurang dari 60 kg, jumlah cairan yang diresepkan oleh tenaga kesehatan Anda mungkin kurang dari 0,5 mL. Tenaga kesehatan Anda akan merekomendasikan berapa banyak cairan yang dibutuhkan.

Penting bahwa jarum selalu berada dalam cairan untuk mencegah terbentuknya gelembung udara di jarum suntik.

Lepaskan jarum dari botol. Pegang jarum suntik dengan jarum mengarah ke atas untuk melihat apakah ada gelembung udara di dalamnya. Jika ada gelembung udara ketuk sisi dengan lembut sampai gelembung udara hilang ke atas jarum suntik dan tekan pendorong sampai semua udara (tetapi tidak ada cairan) keluar. Jangan meletakkan jarum suntik ke bawah atau membiarkan jarum menyentuh apa pun.



LANGKAH 4: PENYUNTIKAN OBAT

Jepit kulit yang dibersihkan dengan lembut di antara ibu jari dan jari telunjuk. Jangan meremasnya.

Dorong jarum suntik ke kulit yang terjepit.



Dorong plunger dengan ibu jari sejauh mungkin untuk menyuntikkan semua cairan.
Dorong perlahan dan merata, jaga kulit dijepit dengan lembut.

Saat pendorong didorong sejauh mungkin, keluarkan jarum dan lepaskan kulit.

Buang Jarum Suntik Kosong dan vial

Buang bagian STELARA yang tidak digunakan. Segera buang jarum suntik kosong ke dalam wadah benda tajam. Untuk keselamatan dan kesehatan Anda dan untuk keselamatan orang lain, botol, alat suntikan dan jarum suntik harus TIDAK BOLEH digunakan kembali. Buang wadah benda tajam sesuai dengan peraturan yang berlaku. Botol dapat dibuang di tempat sampah biasa.

Gunakan Bola Kapas atau Kasa

Mungkin ada sejumlah kecil darah atau cairan di tempat suntikan, dimana hal ini adalah normal. Anda dapat menekan bola kapas atau kain kasa di atas tempat penyuntikan dan tahan selama 10 detik. Jangan menggosok tempat penyuntikan. Anda dapat menutup tempat penyuntikan dengan perban perekat kecil, jika perlu.

LANGKAH 5: APA YANG HARUS DILAKUKAN JIKA ANDA LUPA MENGGUNAKAN STELARA?

Lakukan injeksi berikutnya segera setelah Anda ingat. Jangan menggandakan dosis suntikan.

Lembar informasi ini merangkum informasi paling penting mengenai STELARA. Anda dapat meminta informasi kepada penyedia layanan kesehatan Anda tentang STELARA yang ditulis untuk profesional kesehatan.

Kemasan:

- Stelara 45 mg in 0.5 ml Pre-Filled Syringe
Dus, 1 pre-filled syringe @ 0.5 mL
No. Reg: DKI2060001643A1
- Stelara 90 mg in 1 ml Pre-Filled Syringe
Dus, 1 pre-filled syringe @ 1.0 mL
No. Reg: DKI2360001643C1
- Stelara 45 mg in 0.5 mL vial
Dus, 1 vial @ 45mg/0.5mL
No. Reg: DKI2060001643A2
- Stelara 130 mg in 26mL vial
Dus, 1 vial @ 130mg/26mL
No. Reg: DKI2160001643B1

HARUS DENGAN RESEP DOKTER

Diproduksi oleh:

Cilag AG, Schaffhausen, Switzerland.

Didaftarkan oleh:

PT Integrated Healthcare Indonesia, Jakarta – Indonesia

Untuk pelaporan efek samping dan keluhan kualitas produk, dapat menghubungi drugsafety@jacid.jnj.com atau telp. (021) 2935-3935

Based on CPPI 6-Apr-2022 + CPPI 1-Jun-2023