

## Directions for Use

B. Braun Melsungen AG - 34209 Melsungen, Germany

### Composition

The ready to use emulsion for infusion contains after mixing of the contents of the individual chambers:

#### Active ingredients

	in 1250 ml	in 1875 ml	in 2500 ml
<b>- from the upper, left chamber</b>			
Glucose monohydrate	88.0 g	137.0 g	176.0 g
equivalent to anhydrous glucose	80.0 g	120.0 g	160.0 g
Sodium dihydrogen phosphate dihydrate	1.170 g	1.755 g	2.340 g
Zinc acetate dihydrate	6.625 mg	9.9 mg	13.2 mg
<b>- from the upper, right chamber</b>			
Soya-bean oil	25.0 g	37.5 g	50.0 g
Medium-chain triglycerides	25.0 g	37.5 g	50.0 g
<b>- from the lower chamber</b>			
Isoleucine	2.34 g	3.51 g	4.68 g
Leucine	3.13 g	4.70 g	6.26 g
Lysine hydrochloride	2.84 g	4.26 g	5.68 g
eq. to Lysine	2.26 g	3.39 g	4.52 g
Methionine	1.96 g	2.94 g	3.92 g
Phenylalanine	3.51 g	5.27 g	7.02 g
Threonine	1.82 g	2.73 g	3.64 g
Tryptophan	0.57 g	0.86 g	1.14 g
Valine	2.60 g	3.90 g	5.20 g
Arginine	2.70 g	4.05 g	5.40 g
Histidin hydrochloride monohydrate	1.69 g	2.54 g	3.38 g
eq. to Histidine	1.25 g	1.88 g	2.50 g
Alanine	4.85 g	7.28 g	9.70 g
Aspartic acid	1.50 g	2.25 g	3.00 g
Glutamic acid	3.50 g	5.25 g	7.00 g
Glycine	1.65 g	2.48 g	3.30 g
Proline	3.40 g	5.10 g	6.80 g
Serine	3.00 g	4.50 g	6.00 g
Sodium hydroxide	0.800 g	1.200 g	1.600 g
Sodium chloride	1.081 g	1.622 g	2.162 g
Sodium acetate trihydrate	0.544 g	0.816 g	1.088 g
Potassium acetate	2.943 g	4.415 g	5.886 g
Magnesium acetate tetrahydrate	0.644 g	0.966 g	1.288 g
Calcium chloride dihydrate	0.441 g	0.662 g	0.882 g
Amino acid content [g]	40	60	80
Total nitrogen content [g]	5.7	8.6	11.4
Carbohydrate content [g]	80	120	160
Lipid content [g]	50	75	100
Energy in the form of lipid [kJ/(kcal)]	1990 (475)	2985 (715)	3980 (950)
Energy in the form of carbohydrate [kJ/(kcal)]	1340 (320)	2010 (480)	2680 (640)
Energy in the form of amino acids [kJ/(kcal)]	670 (160)	1005 (240)	1340 (320)
Non-protein energy [kJ/(kcal)]	3330 (795)	4995 (1195)	6660 (1590)
Total energy [kJ/(kcal)]	4000 (955)	6000 (1435)	8000 (1910)
	<b>in 1250 ml</b>	<b>in 1875 ml</b>	<b>in 2500 ml</b>
Osmolality [mOsm/kg]	920	920	920
pH	5.0 - 6.0	5.0 - 6.0	5.0 - 6.0
<b>Electrolyte content (mmol)</b>			
Sodium	50	75	100
Potassium	30	45	60
Magnesium	3.0	4.5	6.0
Calcium	3.0	4.5	6.0
Zinc	0.03	0.045	0.06
Chloride	48	72	96
Acetate	40	60	80
Phosphate	7.5	11.25	15

#### Excipients:

Citric acid monohydrate, egg lecithin, glycerol, sodium oleate, water for injections

#### Pharmaceutical form

Emulsion for infusion in three-chamber bags containing 1250 ml, 1875 ml and 2500 ml.

#### Pharmaco-therapeutic group

Emulsion for intravenous supply of amino acids, carbohydrates, fat and electrolytes.

## NuTRIflex® Lipid peri Amino Acids + Glucose + Lipid Emulsion for Infusion. Especially suitable for Infusion into peripheral veins.

### Indications

Supply of energy, essential fatty acids, amino acids, electrolytes and fluids during parenteral nutrition for patients with mild to moderately severe catabolism when oral or enteral nutrition is impossible, insufficient or contraindicated.

### Contraindications

This product must not be administered in the following conditions

- disturbances of amino acid metabolism,
- disturbances of lipid metabolism,
- hyperkalaemia; hypernatraemia,
- unstable metabolism (e.g. severe postaggression syndrome, unstabilized diabetic metabolic situation, coma of unknown origin),
- hyperglycaemia not responding to insulin doses of up to 6 units insulin/hour,
- acidosis,
- intrahepatic cholestasis,
- severe hepatic insufficiency,
- severe renal insufficiency,
- manifest cardiac insufficiency,
- aggravating haemorrhagic diatheses,
- acute phases of cardiac infarction and stroke,
- acute thrombo-embolic events, lipid embolism,
- known hypersensitivity to egg or soya-bean protein, peanut oil or to any of the excipients

On account of its composition NuTRIflex® Lipid peri should not be used for neonates, infants and children under 2 years of age.

General contra-indications to parenteral nutrition are:

- unstable circulatory status with vital threat (states of collapse and shock),
- inadequate cellular oxygen supply,
- states of hyperhydration,
- disturbances of the electrolyte and fluid balance, acute pulmonary oedema, de-compensated cardiac insufficiency

### Special Warnings and Special Precautions for Use

Due to the individual needs of paediatric patients, NuTRIflex® Lipid peri may not cover sufficiently the total energy requirements. In such cases carbohydrates and / or lipids must be provided in addition, as appropriate.

Caution should be exercised in cases of increased serum osmolality.

As for all large-volume infusion solutions NuTRIflex® Lipid peri should be administered with caution to patients with impaired cardiac or renal function. Disturbances of the fluid, electrolyte or acid-base balance, e.g. hyperhydration, hyperkalaemia, acidosis, should be corrected before the start of infusion. Too rapid infusion can lead to fluid overload with pathological serum electrolyte concentrations, hyperhydration and pulmonary oedema.

The serum triglyceride concentration should be monitored when infusing NuTRIflex® Lipid peri. Fasting lipaemia should be excluded in patients with suspected disturbances of lipid metabolism before starting infusion. The administration of lipids is contra-indicated if there is fasting lipaemia. The presence of hypertriglyceridaemia 12 hours after lipid administration also indicates a disturbance of lipid metabolism. NuTRIflex® Lipid peri should be administered cautiously to patients with disturbances of lipid metabolism, e.g. renal insufficiency, diabetes mellitus, pancreatitis, impaired hepatic function, hypothyroidism (with hypertriglyceridemia) and sepsis. If NuTRIflex® Lipid peri is given to patients with these conditions, close monitoring of serum triglycerides is mandatory.

Any sign or symptom of anaphylactic reaction (such as fever, shivering, rash or dyspnoea) should lead to immediate interruption of the infusion.

Depending on the patient's metabolic condition, occasional hypertriglyceridaemia or increases of the blood glucose concentration may occur. If the plasma triglyceride concentration rises to more than 2 mmol/l during administration of lipid it is recommended that the infusion rate should be reduced. Should the plasma triglyceride concentration remain above 3 mmol/l the administration should be stopped until the level normalizes.

A dose reduction or interruption of administration is also indicated if the blood glucose concentration rises to more than 14 mmol/l (250 mg/dl) when administering the product.

As with all solutions containing carbohydrates the administration of NuTRIflex® Lipid peri can lead to hyperglycaemia. The blood glucose level should be monitored. If there is hyperglycaemia the rate of infusion should be reduced or insulin should be administered.

Intravenous infusion of amino acids is accompanied by increased urinary excretion of the trace elements, especially copper and, in particular, zinc. This should be considered in the dosing of trace elements, especially during long-term intravenous nutrition.

NuTRIflex® Lipid peri should not be given simultaneously with blood in the same infusion set due to the risk of pseudoagglutination.

Moreover controls of the serum electrolytes, the water balance, the acid-base

RENEWA

**B | BRAUN**

balance and - during long-term administration - of blood cell counts, coagulation status and hepatic function are necessary.

The fat content may interfere with certain laboratory measurements (e.g. bilirubin, lactate dehydrogenase, oxygen saturation). If blood is sampled before fat has been adequately cleared from the blood stream.

Substitution of electrolytes, vitamins and trace elements may be necessary as required.

As NuTRiflex® Lipid per contains zinc and magnesium, care should be taken when it is coadministered with solutions containing these elements.

As with all intravenous solutions strict aseptic precautions are necessary for the infusion of NuTRiflex® Lipid per.

NuTRiflex® Lipid per is a preparation of complex composition. It is, therefore, strongly advisable not to add other solutions.

#### Pregnancy and Lactation

Preclinical studies have not been performed with NuTRiflex® Lipid per. The prescriber should consider the benefit/ risk relationship before administering NuTRiflex® Lipid per to pregnant women.

Breast-feeding is not recommended if women need parenteral nutrition in that time.

#### Interactions

Some drugs, like insulin, may interfere with the body's lipase system. This kind of interaction seems, however, to be of only limited clinical importance.

Heparin given in clinical doses causes a transient release of lipoprotein lipase into the circulation. This may result initially in increased plasma lipolysis followed by a transient decrease in triglyceride clearance.

Soya-bean oil has a natural content of vitamin K<sub>1</sub>. This may interfere with the therapeutic effect of coumarin derivatives which should be closely monitored in patients treated with such drugs.

#### Dosage

The dosage is adapted to the individual patients' requirements.

##### Adults:

The maximum daily dose is 40 ml per kg body weight, corresponding to

- 1.28 g amino acids /kg body weight per day

- 2.56 g glucose /kg body weight per day

- 1.6 g fat /kg body weight per day

It is recommended that NuTRiflex® Lipid per be administered continuously. A stepwise increase of the infusion rate over the first 30 minutes up to the desired infusion rate avoids possible complications.

The maximum infusion rate is 2.5 ml/kg body weight per hour, corresponding to

- 0.08 g amino acids /kg body weight per hour

- 0.16 g glucose /kg body weight per hour

- 0.1 g fat /kg body weight per hour

For a patient weighing 70 kg this corresponds to an infusion rate of 175 ml/hour. The amount of amino acid administered is then 5.6 g/hour, of glucose 11.2 g/hour and of lipid 7 g/hour.

##### Children over 2 years of age:

The given dosage recommendations are guiding data based on average requirements. The dosage should be individually adapted, according to age, development stage and illness. For calculation of dosage account must be taken of the hydration status of the paediatric patient.

For children, it might be necessary to start the nutritional therapy with half of the target dosage. The dosage should be increased stepwise according to the individual metabolic capacity up to the maximum dosage.

##### Daily dose during 3<sup>rd</sup> - 5<sup>th</sup> year of life:

45 ml/kg body weight, corresponding to

- 1.44 g amino acids /kg body weight per day

- 2.88 g glucose /kg body weight per day

- 1.8 g lipid /kg body weight per day.

##### Daily dose during 6<sup>th</sup> - 14<sup>th</sup> year of life:

30 ml/kg body weight, corresponding to

- 0.96 g amino acids /kg body weight per day

- 1.92 g glucose /kg body weight per day

- 1.2 g lipid /kg body weight per day

The maximum rate of infusion is 2.5 ml/kg body weight per hour, corresponding to

- 0.08 g amino acids /kg body weight per hour

- 0.16 g glucose /kg body weight per hour

- 0.1 g lipid /kg body weight per hour

Additional energy that may be required for paediatric patients should be administered in the form of glucose solutions or fat emulsions, as appropriate.

#### Method of administration:

For intravenous infusion. Especially suitable for infusion into peripheral veins.

#### Preparation of the mixed solution:

Remove the bag from its protective pack and proceed as follows:

- open out the bag and lay on a solid surface
- open the peel seals to the two upper chambers by using pressure with both hands
- briefly mix the contents of the bag together

#### Preparation for infusion:

- fold the two empty chambers backwards
- hang the mixing bag on the infusion stand by the centre hanging loop
- remove the protective cap from the run-out port and carry out infusion using the normal technique

#### Duration of use

The duration of treatment for the indications stated should not exceed 7 days.

#### Overdose

Overdose of NuTRiflex® Lipid per is not to be expected on proper administration.

#### Symptoms of fluid and electrolyte overdose

Hypertonic hyperhydration, electrolyte imbalance and pulmonary oedema.

#### Symptoms of amino acid overdose:

Renal amino acid losses with consecutive amino acid imbalances, sickness, vomiting and shivering.

#### Symptoms of glucose overdose:

Hyperglycaemia, glucosuria, dehydration, hyperosmolality, hyperglycaemic and hyperosmolar coma.

#### Symptoms of lipid overdose:

Lipid overdose may lead to the overload syndrome, characterised (for example) by fever, headache, abdominal pain, fatigue, hyperlipaemia, hepatomegaly with or without jaundice, splenomegaly, pathological disturbances of liver function, anaemia, reduction in platelet count, reduction in white cell count, haemorrhagic diathesis and haemorrhage, alteration or depression of blood coagulation factors (bleeding time, coagulation time, prothrombin time etc.). The plasma triglyceride concentration should not exceed 3 mmol/l during infusion.

#### Emergency treatment, antidotes:

Immediate cessation of infusion is indicated for overdose. Further therapeutic measures depend on the particular symptoms and their severity. When infusion is recommenced after the symptoms have declined it is recommended that the infusion rate be raised gradually with monitoring at frequent intervals.

#### Undesirable effects

Possible early reactions on the administration of lipid emulsions are: slight increase in temperature, flush, cold feeling, shivering, loss of appetite, nausea, vomiting, respiratory distress, headache, pain in the back, bones, chest and lumbar region, fall or increase in blood pressure (hypotension, hypertension), hypersensitivity reactions (e.g. anaphylactic reactions, dermal eruptions).

Hot flushes or bluish discoloration of the skin due to reduced oxygen content of the blood (cyanosis) can occur as side effects.

If these side effects occur the infusion should be discontinued or, if appropriate, the infusion should be continued at a lower dose level.

Attention should be paid to the possibility of an overloading syndrome. This can occur as a result of individually varying, genetically determined metabolic conditions and can occur at different rates and after differing doses depending on previous disorders.

Overloading syndrome is associated with the following symptoms: enlargement of the liver (hepatomegaly) with and without jaundice (icterus), enlargement of the spleen (splenomegaly), fatty infiltration of the organs, pathological hepatic function parameters, anaemia, reduction of white cell count (leucopenia), reduction of platelet count (thrombocytopenia), a tendency to haemorrhage and haemorrhages, alterations or reduction in the blood coagulation factors (bleeding time, coagulation time, prothrombin time etc.), fever, hyperlipaemia, headache, stomach-ache, fatigue.

If signs of vein wall irritation, phlebitis, or thrombophlebitis occur, change of the infusion site should be considered.

Please inform your doctor or pharmacist if you notice any undesirable effect that is not mentioned in this leaflet.

#### Instructions for storage / use / handling

Do not use the product beyond the expiry date stated on the labelling.

The emulsion is to be used immediately after mixing. It can be stored at 2 - 8 °C over 4 days, plus 48 hours at 25 °C.

The ready-to-use emulsion can be stored for 4 days at 2 - 8 °C plus 48 hours at 25 °C.

The emulsion is to be used immediately after connecting the container to the giving set.

NuTRiflex® Lipid per is supplied in single dose containers. Unused residues must be discarded.

If filters are used they must be lipid-permeable.

Do not store above 25 °C.

Do not freeze. If accidentally frozen, discard the bag.

Only use bags that are undamaged and in which the amino acid and glucose solutions are clear. Do not use bags where there is discernible phase separation (oil drops) in the chamber containing lipid emulsion.

Keep bags in the outer carton in order to protect from light.

Date of last revision 01/2006

#### Presentation:

NuTRiflex® Lipid per: 1250 ml, 1875 ml, 2500 ml

3 chamber compartment bag

Reg. No.: DK11386701849A1

Harus dengan resep dokter

Licence Holder:

PT. Pliapros Tbk., Indonesia

Imported by

PT. B. Braun Medical Indonesia

Jakarta, Indonesia.

Manufactured by:

B. Braun Melsungen AG

34209 Melsungen, Germany

**B | BRAUN**

B. Braun Melsungen AG  
34209 Melsungen, Germany

**RENEWAL**

1250 ml of the ready to use emulsion contain:  
- from the upper, left chamber

Glucose monohydrate	88.0 g
eq. to anhydrous glucose	80.0 g
Sodium dihydrogen phosphate dihydrate	1.170 g
Zinc acetate dihydrate	6.625 mg

Reg. No.: DK11386701849A1

HET : Rp. 654.000



Manuf. date:                      Batch no.:                      Expiry date:




## NuTRiflex® Lipid peri Amino Acids + Glucose + Lipid Emulsion for Infusion.

Especially suitable for Infusion  
into peripheral veins.

1250 ml of the ready to use  
emulsion contain:  
- from the upper, right chamber

Soya-bean oil	25.0 g
Medium-chain triglycerides	25.0 g



HKS 47  
schwarz (passer, druckt nicht)

Dokument = 214 x 330 mm

Lätus 2110



ID\_\_\_176  
176/FF79602/0114  
3KB (1250ml)  
Standort Melsungen

Fontsize: 8,0 pt.

2000

## NuTRiflex® Lipid peri Amino Acids + Glucose + Lipid Emulsion for Infusion.

Especially suitable for Infusion into peripheral veins.

Content of amino acids	3.20 %
Content of carbohydrates	6.40 %
Content of fat	4 %

1250 ml of the ready to use emulsion  
contain after mixing of the three cham-  
ber contents:

- from the lower chamber

Isoleucine	2.34 g
Leucine	3.13 g
Lysine hydrochloride eq. to Lysine 2.26 g	2.84 g
Methionine	1.96 g
Phenylalanine	3.51 g
Threonine	1.82 g
Tryptophan	0.57 g
Valine	2.60 g
Arginine	2.70 g
Histidine hydrochloride monohydrate eq. to Histidine 1.25 g	1.69 g
Alanine	4.85 g
Aspartic acid	1.50 g
Glutamic acid	3.50 g
Glycine	1.65 g

Proline	3.40 g
Serine	3.00 g
Sodium hydroxide	0.800 g
Sodium chloride	1.081 g
Sodium acetate trihydrate	0.544 g
Potassium acetate	2.943 g
Magnesium acetate tetrahydrate	0.644 g
Calcium chloride dihydrate	0.441 g
Excipients: Citric acid monohydrate, egg lecithin, glycerol, sodium oleate, water for injections	
Electrolytes:	
Sodium	50 mmol
Potassium	30 mmol
Magnesium	3.0 mmol
Calcium	3.0 mmol
Chloride	48 mmol
Acetate	40 mmol
Phosphate	7.5 mmol
Zinc	0.03 mmol

Amino acid content	40 g
Total nitrogen content	5.7 g
Carbohydrate content	80 g
Lipid content	50 g
Energy in the form of lipid	1990 kJ (= 475 kcal)
Energy in the form of carbohydrate	1340 kJ (= 320 kcal)
Energy in the form of amino acids	670 kJ (= 160 kcal)
Non-protein energy	3330 kJ (= 795 kcal)
Total energy	4000 kJ (= 955 kcal)
Osmolality	920 mOsm/kg
pH	5.0 - 6.0

Only use bags that are undamaged and  
in which the amino acid and glucose so-  
lutions are clear. Do not use bags where  
there is discernible phase separation (oil  
drops) in the chamber containing lipid  
emulsion.

Sterile and apyrogenic.  
Keep bags in the outer carton in order  
to protect from light.  
Keep out of the reach and sight of  
children.

Harus dengan resep dokter

Content: 1250 ml

1500

1000

500

**B | BRAUN**



**RENEWAL**

Read Directions for Use carefully.  
Mix contents of the chambers before  
start of the infusion.  
Do not store above 25 °C.  
For single use only. Discard unused con-  
tents.  
Do not freeze. If accidentally frozen, dis-  
card the bag.

176/12606779/0114

# NuTRIflex® Lipid peri

## Amino Acids + Glucose + Lipid Emulsion for Infusion.

Especially suitable for Infusion into peripheral veins.

Content of amino acids 3.20 %  
 Content of carbohydrates 6.40 %  
 Content of fat 4 %

1250 ml of the ready to use emulsion contain after mixing of the three chamber contents:

- from the upper, left chamber

Glucose monohydrate 88.0 g  
 eq. to anhydrous glucose 80.0 g  
 Sodium dihydrogen phosphate dihydrate 1.170 g  
 Zinc acetate dihydrate 6.625 mg

- from the upper, right chamber

Soya-bean oil 25.0 g  
 Medium-chain triglycerides 25.0 g

- from the lower chamber

Isoleucine 2.34 g  
 Leucine 3.13 g  
 Lysine hydrochloride eq. to Lysine 2.26 g  
 Methionine 1.96 g  
 Phenylalanine 3.51 g  
 Threonine 1.82 g  
 Tryptophan 0.57 g  
 Valine 2.60 g  
 Arginine 2.70 g  
 Histidine hydrochloride monohydrate eq. to Histidine 1.25 g

Alanine 4.85 g  
 Aspartic acid 1.50 g  
 Glutamic acid 3.50 g  
 Glycine 1.65 g  
 Proline 3.40 g  
 Serine 3.00 g  
 Sodium hydroxide 0.800 g  
 Sodium chloride 1.081 g  
 Sodium acetate trihydrate 0.544 g  
 Potassium acetate 2.943 g  
 Magnesium acetate tetrahydrate 0.644 g  
 Calcium chloride dihydrate 0.441 g

Excipients: Citric acid monohydrate, egg lecithin, glycerol, sodium oleate, water for injections

Electrolytes:  
 Sodium 50 mmol  
 Potassium 30 mmol  
 Magnesium 3.0 mmol  
 Calcium 3.0 mmol  
 Chloride 48 mmol  
 Acetate 40 mmol  
 Phosphate 7.5 mmol  
 Zinc 0.03 mmol

Amino acid content 40 g  
 Total nitrogen content 5.7 g  
 Carbohydrate content 80 g  
 Lipid content 50 g  
 Energy in the form of lipid 1990 kJ (= 475 kcal)  
 Energy in the form of carbohydrate 1340 kJ (= 320 kcal)  
 Energy in the form of amino acids 670 kJ (= 160 kcal)  
 Non-protein energy 3330 kJ (= 795 kcal)  
 Total energy 4000 kJ (= 955 kcal)  
 Osmolality 920 mOsm/kg  
 pH 5.0 - 6.0

Read Directions for Use carefully. Mix contents of the chambers before start of the infusion. Do not store above 25°C. For single use only. Discard unused contents. Do not freeze. If accidentally frozen, discard the bag.

Only use bags that are undamaged and in which the amino acid and glucose solutions are clear. Do not use bags where there is discernible phase separation (oil drops) in the chamber containing lipid emulsion. Sterile and apyrogenic. Keep bags in the outer carton in order to protect from light. Keep out of the reach and sight of children.

Harus dengan resep dokter

Reg. No. : DK11386701849A1  
 HET : Rp. 5 x 654.000



Art. No.: 353 3140

Content: 5 x 1250 ml

Manuf. date:

Batch-no.:

Expiry date:

Licence Holder:  
 PT. Phapros Tbk., Indonesia  
 Imported by:  
 PT. B. Braun Medical Indonesia  
 Jakarta, Indonesia  
 Manufactured by:  
 B. Braun Melsungen AG  
 34209 Melsungen, Germany

**RENEWAL**



**B | BRAUN**



12606779\_NuTRIflexLipid-peri\_KET5x1250\_ID.indd 1

03.01.14 09:40

Schwarz

HKS 47

HKS 13

Blank (M03; var.Feld, druckt nicht)

Label = 162 x 116 mm

Lätus 400



400

ID\_\_176

176/12606779/0114

3KB-KET 5x1250 ml

Standort Melsungen

Font size: 7,0 pt.