

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
Trade Name: Zavicefta®
CDS Effective Date: February 28, 2024
Supersedes: November 18, 2021
Approved by BPOM: January 31, 2025

PT. Pfizer Indonesia
Local Product Document

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
Trade Name: Zavicefta®
CDS Effective Date: February 28, 2024
Supersedes: November 18, 2021

1. NAME OF THE MEDICINAL PRODUCT

1.1 Product name

Zavicefta®

1.2 Strength

2 g/0.5 g

1.3 Pharmaceutical dosage form

Powder for solution for infusion.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

2.1 Qualitative declaration

Each vial contains ceftazidime (as pentahydrate) equivalent to 2 g and avibactam (as sodium salt) equivalent to 0.5 g.

After reconstitution, 1 ml of solution contains 167.3 mg of ceftazidime and 41.8 mg of avibactam (See section 6.6).

2.2 Quantitative declaration

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder for solution for infusion.

A white to yellow sterile powder.

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
Trade Name: Zavicefta®
CDS Effective Date: February 28, 2024
Supersedes: November 18, 2021
Approved by BPOM: January 31, 2025

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Zavicefta® is indicated in adults for the treatment of the following infections of ESBL (Extended Spectrum Beta Lactamase) producing bacteria and *Pseudomonas aeruginosa* resistance to ceftazidime as follow:

- Complicated Intra-Abdominal Infection (cIAI): require surgical intervention and extend beyond the hollow viscus of the intestinal tract into the peritoneal space (laparotomy, percutaneous drainage of an abscess, or laparoscopic surgery). For treatment of cIAI use in combination with metronidazole.
- Complicated Urinary Tract Infection, including Pyelonephritis (cUTI): acute pyelonephritis indicated by flank pain or costovertebral angle tenderness on examination and at least one of the following: fever, nausea, vomiting; or complicated lower UTI without pyelonephritis indicated by qualifying symptoms (dysuria, urgency, frequency, suprapubic pain, fever, nausea, vomiting) plus at least 1 complicating factor (history of chronic urinary retention, obstructive uropathy, functional or anatomical abnormality of the urogenital tract, use of bladder catheterization, urogenital procedures).
- Hospital-acquired Pneumonia (HAP), including ventilator associated pneumonia (VAP): pneumonia that arises after ≥ 48 hours in an inpatient acute or chronic care facility or that developed within 7 days after being discharged; parenchymal lung infection that arises ≥ 48 hours after endotracheal intubation and mechanical ventilation.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

The recommended dosage of Zavicefta® is 1 vial where each vial contains 2 g ceftazidime and 0.5 g avibactam administered by intravenous (IV) infusion in a volume of 100 ml at a constant rate over 120 minutes in patients aged 18 years or older. Treatment is repeated every 8 hours. For patients with renal impairment where $\text{CrCl} \leq 50$ ml/min, see dose recommendations in Table 2.

Treatment duration

Table 1 Summary of The Treatment Duration by Indication or Condition

Indication	Treatment Duration
Complicated Intra-Abdominal Infection (cIAI)	5-14 days
Complicated Urinary Tract Infection (cUTI), including Pyelonephritis	5-10 days ¹
Hospital-acquired Pneumonia, including ventilator associated pneumonia	7-14 days

¹ Treatment duration includes intravenous plus oral treatment. The time to switch from intravenous Zavicefta® to oral treatment with another antibiotic depends on the clinical situation, but is normally after about 5 days (the minimum duration of treatment with ceftazidime-avibactam in clinical trials was 5 days).

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium

Trade Name: Zavicefta®

CDS Effective Date: February 28, 2024

Supersedes: November 18, 2021

Approved by BPOM: January 31, 2025

For Complicated Urinary Tract Infection (cUTI) including Pyelonephritis, the total duration of treatment could be increased to 14 days for patients with bacteraemia.

The duration of treatment should be guided by the severity of the infection, the pathogen(s) and the patient's clinical and bacteriological progress.

Special populations

Elderly patients

No dosage adjustment is considered necessary in elderly patients (≥ 65 years). The dose regimen should be adjusted if renal impairment is present (see section 5.2).

Patients with renal impairment

The following dose adjustment is recommended in patients with renal impairment (see sections 4.4 and 5.2).

Dose adjustments for Zavicefta® for patients with an estimated creatinine clearance (CrCl) ≤ 50 ml/min are outlined in Table 2 below. The only information on dosing of Zavicefta® for patients requiring dialysis is in the setting of intermittent haemodialysis. For other types of dialysis, it is suggested that the dose/frequency of ceftazidime-avibactam should follow local label/local guidelines for dosing of ceftazidime. For example, for a dose of 500 mg ceftazidime the dose of ceftazidime-avibactam would be 500 mg ceftazidime/125 mg avibactam.

Table 2 Recommended Dose for Patients with Renal Impairment*

Estimated CrCl (ml/min) ^a	Recommended Dosage Regimen Ceftazidime/Avibactam	Infusion Time (hours)	Frequency of Dosing (hourly)
50-31	1000 mg/250 mg	2	Every 8 hours
30-16	750 mg/187.5 mg	2	Every 12 hours
15 to 6	750 mg/187.5 mg ^b	2	Every 24 hours
<6	750 mg/187.5 mg ^b	2	Every 48 hours

^a Creatinine Clearance (CrCl) calculated using the Cockcroft-Gault formula.

^b Both ceftazidime and avibactam are haemodialyzable; thus, Zavicefta® should be administered after haemodialysis on haemodialysis day.

* Dose recommendations are based on PK modelling.

In patients with impaired renal function, regular monitoring of estimated creatinine clearance is advised as in some patients, especially early in the course of their infection, the creatinine clearance estimated from serum creatinine can change quickly.

Haemodialysis

Both ceftazidime and avibactam are haemodialyzable; thus, Zavicefta® should be administered after haemodialysis on haemodialysis day.

Haemofiltration

There is insufficient data to make specific dosage adjustment recommendations for patients undergoing continuous veno-venous haemofiltration.

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
Trade Name: Zavicefta®
CDS Effective Date: February 28, 2024
Supersedes: November 18, 2021
Approved by BPOM: January 31, 2025

Peritoneal dialysis

There is insufficient data to make specific dosage adjustment recommendations for patients undergoing peritoneal dialysis.

Patients with hepatic impairment

No dosage adjustment is considered necessary in patients with hepatic impairment (see section 5.2). Close clinical monitoring for safety and efficacy is advised.

Paediatric patients

Safety and efficacy in paediatric patients (<18 years of age) have not been established (see section 5.2).

Method of administration

Zavicefta® is administered by intravenous infusion over 120 minutes in an infusion volume of 100 ml (see section 6.6).

Constitution and compatibility

For instructions on reconstitution and dilution of the medicinal product before administration (see section 6.6).

4.3 Contraindications

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.
Hypersensitivity to the cephalosporin class of antibacterials.

Immediate and severe hypersensitivity (e.g. anaphylactic reaction) to any other type of β -lactam antibacterial agent (e.g. penicillins, monobactams or carbapenems).

4.4 Special warnings and precautions for use

Hypersensitivity reactions

As with all β -lactam antibacterial agents, serious and occasionally fatal hypersensitivity reactions have been reported. In case of severe hypersensitivity reactions, treatment with Zavicefta® must be discontinued immediately and adequate emergency measures must be initiated.

Before beginning treatment, it should be established whether the patient has a history of severe hypersensitivity reactions to ceftazidime, to other cephalosporins or to any other type of β -lactam agent. Caution should be used if ceftazidime-avibactam is given to patients with a history of non-severe hypersensitivity to other β -lactam agents.

***Clostridium difficile*-associated diarrhoea**

Antibacterial agent-associated colitis and pseudo-membranous colitis have been reported with nearly all anti-bacterial agents, including ceftazidime-avibactam, and may range in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhoea during or subsequent to the administration of Zavicefta® (see section 4.8). Discontinuation of therapy with Zavicefta® and the administration of specific treatment for *Clostridium difficile* should be considered. Medicinal products that inhibit peristalsis should not be given.

Patients with renal impairment

Ceftazidime and avibactam are eliminated via the kidneys, therefore the dose should be reduced according to the degree of renal impairment. Patients with renal impairment should be closely monitored for both safety and efficacy. Neurological sequelae, including tremor, myoclonus, nonconvulsive status epilepticus, convulsion, encephalopathy and coma, have occasionally been reported with ceftazidime when the dose has not been reduced in patients with renal impairment (see section 4.2).

In patients with renal impairment, close monitoring of estimated creatinine clearance is advised. In some patients, the creatinine clearance estimated from serum creatinine can change quickly, especially early in the course of treatment for the infection.

Nephrotoxicity

Concurrent treatment with high doses of cephalosporins and nephrotoxic medicinal products such as aminoglycosides or potent diuretics (e.g. furosemide) may adversely affect renal function.

Non-susceptible organisms

Prolonged use may result in the overgrowth of non-susceptible organisms (e.g. enterococci, fungi), which may require interruption of treatment or other appropriate measures.

Non-drug interference

Ceftazidime does not interfere with enzyme-based tests for glycosuria, but slight interference (false-positive) may occur with copper reduction methods (Benedict's, Fehling's, Clinitest).

Direct antiglobulin test (DAGT or Coombs test) seroconversion and potential risk of haemolytic anaemia

Cephalosporin use may cause development of a positive direct antiglobulin test (DAGT, or Coombs test), which may interfere with the cross-matching of blood and/or may cause drug induced immune hemolytic anemia. While DAGT seroconversion in patients receiving Zavicefta® was frequent in clinical studies, there was no evidence of haemolysis in patients who developed a positive DAGT on treatment (see section 4.8). However, the possibility that haemolytic anaemia could occur in association with Zavicefta® treatment cannot be ruled out. Patients experiencing anaemia during or after treatment with Zavicefta® should be investigated for this possibility.

Spectrum of activity of ceftazidime/avibactam

Ceftazidime has little or no activity against the majority of Gram-positive organisms and anaerobes. Additional antibacterial agents should be used when these pathogens are known or suspected to be contributing to the infectious process.

The inhibitory spectrum of avibactam includes many of the enzymes that inactivate ceftazidime, including Ambler class A β -lactamases and class C β -lactamases. Avibactam does not inhibit class B enzymes (metallo- β -lactamases) and is not able to inhibit many of the class D enzymes.

Controlled sodium diet

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium

Trade Name: Zavicefta®

CDS Effective Date: February 28, 2024

Supersedes: November 18, 2021

Approved by BPOM: January 31, 2025

For patients who are on a controlled sodium diet, the following important information about the ingredients of ceftazidime and avibactam should be considered:

- 2 g powder for solution for infusion

Ceftazidime 2 g contains 4.52 mmol of sodium per vial; and

- 500 mg powder for solution for infusion

Avibactam 500 mg contains 1.92 mmol of sodium per vial.

4.5 Interaction with other medicinal products and other forms of interaction

Concurrent treatment with high doses of cephalosporins and nephrotoxic medicinal products such as aminoglycosides or potent diuretics (e.g. furosemide) may adversely affect renal function (see section 4.4).

Chloramphenicol is antagonistic *in vitro* with ceftazidime and other cephalosporins. The clinical relevance of this finding is unknown, but if concurrent administration of ceftazidime-avibactam with chloramphenicol is proposed, the possibility of antagonism should be considered.

Avibactam showed no significant inhibition of cytochrome P450 enzymes. Avibactam and ceftazidime showed no *in vitro* cytochrome P450 induction in the clinically relevant exposure range. Avibactam and ceftazidime do not inhibit the major renal or hepatic transporters in the clinically relevant exposure range, therefore the drug-drug interaction potential via these mechanisms is considered low.

In vitro, avibactam is a substrate of OAT1 and OAT3 transporters which might contribute to the active uptake from the blood compartment and, thereby its excretion. Probenecid (a potent OAT inhibitor) inhibits this uptake by 56% to 70% *in vitro* and, therefore, has the potential to alter the elimination of avibactam when co-dosed. Since a clinical interaction study of avibactam and probenecid has not been conducted, co-dosing of avibactam with probenecid is not recommended.

4.6 Pregnancy and lactation

Pregnancy

There is limited clinical data from the use of ceftazidime-avibactam in pregnant women. Animal embryofetal development studies conducted with ceftazidime or avibactam do not indicate harmful effects at exposures equivalent to therapeutic concentrations. Following administration of avibactam throughout pregnancy and lactation in the rat at maternal exposures greater than or equal to approximately 1.5 times human therapeutic exposures, there were minor changes in the morphology of the kidney and ureters in the rat pups (see section 5.3).

Ceftazidime-avibactam should not be used during pregnancy unless clearly necessary and only if the potential benefit outweighs the possible risk.

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
Trade Name: Zavicefta®
CDS Effective Date: February 28, 2024
Supersedes: November 18, 2021
Approved by BPOM: January 31, 2025

Lactation

There are no data on human milk excretion of ceftazidime-avibactam. Ceftazidime is excreted in human milk in small quantities. It is unknown whether avibactam is excreted in human milk. Women who are breast-feeding should be treated with ceftazidime-avibactam only if clearly indicated. Interruption of breast-feeding is recommended.

Fertility

The effects of ceftazidime-avibactam on fertility in humans have not been studied. Animal studies with ceftazidime or avibactam do not indicate harmful effects with respect to fertility (see section 5.3).

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. However, undesirable effects may occur (e.g. dizziness), which may influence the ability to drive and use machines (see section 4.8).

4.8 Undesirable effects

In seven Phase 2 and Phase 3 clinical trials, 2024 adult patients were treated with Zavicefta®. The most common adverse reactions occurring in $\geq 5\%$ of patients treated with Zavicefta® were Coombs direct test positive, nausea, and diarrhoea. These were usually mild or moderate in intensity. No clinically significant differences were observed in the safety profile across indications.

The following adverse reactions have been reported with ceftazidime alone and/or identified during all Phase 2 and Phase 3 clinical trials with Zavicefta® (N=2024). Adverse reactions are classified according to frequency and System Organ Class. Frequency categories are derived from adverse reactions and/or potentially clinically significant laboratory abnormalities, and are defined according to the following conventions:

Very common ($\geq 1/10$)
Common ($\geq 1/100$ and $< 1/10$)
Uncommon ($\geq 1/1,000$ and $< 1/100$)
Rare ($\geq 1/10,000$ and $< 1/1,000$)
Very rare ($< 1/10,000$)
Unknown (cannot be estimated from the available data)

If an event was not seen in the overall Phase 2 and Phase 3 pool but was a known ADR for ceftazidime alone, the frequency category for ceftazidime alone was used (including the category Unknown).

Table 3 Frequency of adverse reactions by system organ class

System Organ Class	Very Common	Common	Uncommon	Very Rare	Unknown
Infections and infestations		Candidiasis (including Vulvovaginal candidiasis and	Clostridium difficile colitis Pseudomembranous		

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium

Trade Name: Zavicefta®

CDS Effective Date: February 28, 2024

Supersedes: November 18, 2021

Approved by BPOM: January 31, 2025

System Organ Class	Very Common	Common	Uncommon	Very Rare	Unknown
		Oral candidiasis)	colitis		
Blood and lymphatic system disorders	Coombs direct test positive ¹	Eosinophilia Thrombocytosis Thrombocytopenia	Neutropenia Leukopenia Lymphocytosis		Agranulocytosis Haemolytic anemia
Immune system disorders					Anaphylactic reaction Kounis syndrome ²
Nervous system disorders		Headache Dizziness	Paraesthesia		
Gastrointestinal disorders		Diarrhoea Abdominal pain Nausea Vomiting	Dysgeusia		
Hepatobiliary disorders		Alanine aminotransferase increased Aspartate aminotransferase increased Blood alkaline phosphatase increased Gamma-glutamyltransferase increased Blood lactate dehydrogenase increased			Jaundice
Skin and subcutaneous tissue disorders		Rash maculo-papular Urticaria Pruritus			Toxic epidermal necrolysis Stevens-Johnson syndrome

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
Trade Name: Zavicefta®
CDS Effective Date: February 28, 2024
Supersedes: November 18, 2021
Approved by BPOM: January 31, 2025

System Organ Class	Very Common	Common	Uncommon	Very Rare	Unknown
					Erythema multiforme Angioedema Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)
Renal and urinary disorders			Blood creatinine increased Blood urea increased Acute kidney injury	Tubulointerstitial nephritis	
General disorders and administration site conditions		Infusion site thrombosis Infusion site phlebitis Pyrexia			

¹ See section 4.4.

² Acute coronary syndrome associated with an allergic reaction

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via:

Pusat Farmakovigilans/MESO Nasional
Direktorat Pengawasan Keamanan, Mutu, dan Ekspor Impor Obat, Narkotika, Psikotropika, Prekursor dan Zat Adiktif
Badan Pengawas Obat dan Makanan
Jl. Percetakan Negara No. 23, Jakarta Pusat, 10560
Email: pv-center@pom.go.id
Phone: +62-21-4244691 Ext.1079
Website: <https://e-meso.pom.go.id/ADR>

PT Pfizer Indonesia
Email: IDN.AEReporting@pfizer.com
Website: www.pfizersafetyreporting.com

4.9 Overdose

Overdosage of ceftazidime-avibactam is unlikely, although overdosing could potentially occur in patients with moderate to severe renal impairment, and in end stage renal disease

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium

Trade Name: Zavicefta®

CDS Effective Date: February 28, 2024

Supersedes: November 18, 2021

Approved by BPOM: January 31, 2025

including patients undergoing haemodialysis (see sections 4.4 and 5.2). Overdosing with ceftazidime-avibactam can lead to neurological sequelae including encephalopathy, convulsions and coma, due to the ceftazidime component.

Treatment for overdose should follow local standard medical practice. Both ceftazidime and avibactam can be partially removed by haemodialysis.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Mechanism of action

Ceftazidime inhibits bacterial peptidoglycan cell wall synthesis following attachment to penicillin binding proteins (PBPs), which leads to bacterial cell lysis and death. This broad spectrum cephalosporin is active against many important Gram-negative and Gram-positive bacterial pathogens *in vitro*. Avibactam is a non β -lactam, β -lactamase inhibitor that acts by forming a covalent adduct with the enzyme that is stable to hydrolysis. It inhibits both Ambler class A and class C β -lactamases, including extended-spectrum β -lactamases (ESBLs), KPC carbapenemases, and AmpC enzymes. Avibactam also inhibits the class D carbapenemase OXA-48, which does not significantly hydrolyze ceftazidime. Avibactam has no clinically relevant *in vitro* antibacterial activity. Avibactam did not induce transcription of *bla*_{AmpC} in *Enterobacter cloacae*, *Citrobacter freundii* or *Pseudomonas aeruginosa in vitro* at concentrations used to treat patients.

Mechanism of resistance

Ceftazidime-avibactam is not active against metallo- β -lactamase-producing bacteria. Bacterial resistance mechanisms that could potentially affect ceftazidime-avibactam include mutant or acquired PBPs, decreased outer membrane permeability to either compound, active efflux of either compound, mutated or acquired β -lactamase enzymes insensitive to avibactam and able to hydrolyze ceftazidime.

Cross-resistance

An absence of cross-resistance between ceftazidime-avibactam and fluoroquinolones or aminoglycosides has been demonstrated *in vitro* using molecularly-characterized clinical isolates. Some isolates resistant to ceftazidime (and other cephalosporins) or to carbapenems are susceptible to ceftazidime-avibactam. There is cross-resistance with β -lactam antibacterial agents, including carbapenems, when the mechanism is production of metallo- β -lactamases, such as VIM-2.

Interaction with other antimicrobial agents

In vitro interaction tests with ceftazidime-avibactam show ceftazidime-avibactam has little potential to antagonize or be antagonized by other antibiotics of various classes (e.g. metronidazole, tobramycin, levofloxacin, vancomycin, linezolid, colistin, tigecycline).

Susceptibility testing

Minimum Inhibitory Concentration (MIC) breakpoints established by the European Committee on Antimicrobial Susceptibility Testing (EUCAST) for ceftazidime/avibactam are as follows:

Organisms	Susceptible	Resistant
-----------	-------------	-----------

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
Trade Name: Zavicefta®
CDS Effective Date: February 28, 2024
Supersedes: November 18, 2021
Approved by BPOM: January 31, 2025

<i>Enterobacteriaceae</i>	≤8 mg/L	>8 mg/L
<i>Pseudomonas aeruginosa</i>	≤8 mg/L	>8 mg/L

Pharmacokinetic/pharmacodynamic relationship

The antimicrobial activity of ceftazidime-avibactam against specific pathogens has been shown to best correlate with the percent time of free-drug concentration above the ceftazidime-avibactam minimum inhibitory concentration over the dose interval ($\%fT > MIC$ of ceftazidime-avibactam) for ceftazidime, and the percent time of the free drug concentration above a threshold concentration over the dose interval ($\%fT > C_T$) for avibactam.

Clinical efficacy against specific pathogens

Efficacy has been demonstrated in clinical studies against the pathogens, listed under each indication, that were susceptible to ceftazidime-avibactam *in vitro*.

Complicated intra-abdominal infections

Gram-negative micro-organisms

- *Citrobacter freundii*
- *Enterobacter cloacae*
- *Escherichia coli*
- *Klebsiella oxytoca*
- *Klebsiella pneumoniae*
- *Pseudomonas aeruginosa*

Complicated urinary-tract infections

Gram-negative micro-organisms

- *Escherichia coli*
- *Klebsiella pneumoniae*
- *Proteus mirabilis*
- *Enterobacter cloacae*
- *Pseudomonas aeruginosa*

Hospital-acquired pneumonia including ventilator-associated pneumonia

Gram-negative micro-organisms

- *Enterobacter cloacae*
- *Escherichia coli*
- *Klebsiella pneumoniae*
- *Proteus mirabilis*
- *Serratia marcescens*
- *Pseudomonas aeruginosa*

Clinical efficacy has not been established against the following pathogens that are relevant to the approved indications although *in vitro* studies suggest that they would be susceptible to ceftazidime-avibactam in the absence of acquired mechanisms of resistance.

Gram-negative micro-organisms

- *Citrobacter koseri*
- *Enterobacter aerogenes*

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
 Trade Name: Zavicefta®
 CDS Effective Date: February 28, 2024
 Supersedes: November 18, 2021
 Approved by BPOM: January 31, 2025

- *Morganella morganii*
- *Proteus vulgaris*
- *Providencia rettgeri*

Ceftazidime-avibactam is active *in vitro* against *Streptococcus pyogenes* and *Streptococcus agalactiae*, but not generally active against other clinically-important Gram-positive bacteria including methicillin-resistant *Staphylococcus aureus* (MRSA).

Clinical efficacy and safety

Complicated intra-abdominal infections

A total of 1058 adults with complicated intra-abdominal infections (defined as infections that require surgical intervention and extend beyond the hollow viscus into the intraperitoneal space) were randomised and received treatment in two identical randomised, multi-centre, multinational, double-blind studies (RECLAIM 1 and RECLAIM 2) comparing Ceftazidime-avibactam (2000 mg of ceftazidime and 500 mg of avibactam) administered intravenously over 120 minutes every 8 hours plus metronidazole (500 mg) to meropenem (1000 mg) administered intravenously over 30 minutes. Treatment duration was 5 to 14 days. The modified intent-to-treat (MITT) population included all patients who met the disease definition of cIAI and received at least 1 dose of the study drug. The clinically evaluable (CE) population included patients who had an appropriate diagnosis of cIAI and excluded patients with a bacterial species typically not expected to respond to both study drugs (i.e. *Acinetobacter baumannii* or *Stenotrophomonas* spp.) and/or who had an important protocol deviation impacting the assessment of efficacy.

The primary efficacy endpoint was the clinical response at the Test of Cure (TOC) visit in the co-primary populations of the CE and MITT patients in Table 4 below.

Table 4 Clinical Cure Rate at TOC (RECLAIM MITT and CE Analysis Sets)

Analysis set	Number (%) of patients		
	CAZ-AVI + MTZ	Meropenem	Difference (%) 95% CI
MITT	(N=520)	(N=523)	
Clinical cure	429 (82.5)	444 (84.9)	-2.4 (-6.90, 2.10)
CE	(N=410)	(N=416)	
Clinical cure	376 (91.7)	385 (92.5)	-0.8 (-4.61, 2.89)

Clinical cure rates at TOC by pathogen in the microbiologically Modified Intent to Treat (mMITT) population for Gram-negative aerobes are shown in Table 5 below.

Table 5 Clinical cure rate at TOC by common (combined frequency of ≥10) Gram-negative baseline pathogen (RECLAIM mMITT analysis set)

Pathogen	Number of patients					
	CAZ-AVI + MTZ (N=413)			Meropenem (N=410)		n
	Cure rate (%)	Number of clinical cures	N	Cure rate (%)	Number of clinical cures	
<i>Enterobacteriaceae</i>	81.4	272	334	86.4	305	353
<i>Citrobacter freundii</i> complex	77.8	14	18	75.0	9	12
<i>Enterobacter aerogenes</i>	80.0	4	5	100	5	5
<i>Enterobacter cloacae</i>	84.6	11	13	84.2	16	19
<i>Escherichia coli</i>	80.4	218	271	87.0	248	285
<i>Klebsiella oxytoca</i>	77.8	14	18	80.0	12	15
<i>Klebsiella pneumoniae</i>	78.4	40	51	75.5	37	49
<i>Proteus mirabilis</i>	62.5	5	8	77.8	7	9
<i>Pseudomonas aeruginosa</i>	85.7	30	35	94.4	34	36

A further 432 adults with complicated intra-abdominal infections were randomised and received treatment in a multi-centre, double-blind study (RECLAIM 3) conducted in 3 Asian countries (China, Republic of Korea and Vietnam). The patient population and key aspects of the study design were identical to RECLAIM apart from the primary efficacy endpoint of clinical response at the TOC visit being solely in the CE population (see Table 6 below).

Table 6 Clinical cure rates at TOC (RECLAIM 3 CE at TOC analysis set)

	Number (%) of patients		
	CAZ-AVI + MTZ	Meropenem	Difference (%) 95% CI
	(N=177)	(N=184)	
Clinical cure	166 (93.8)	173 (94.0)	-0.2 (-5.53, 4.97)

Clinical cure rates at TOC by pathogen in the microbiologically modified Intent to Treat (mMITT) population for Gram-negative aerobes are shown in Table 7 below.

Table 7 Clinical cure rates at TOC by common (combined frequency of ≥7) Gram- negative baseline pathogen (RECLAIM 3 mMITT analysis set)

Pathogen	Number of patients					
	CAZ-AVI + MTZ (N=143)			Meropenem (N=152)		
	Cure rate (%)	Number of clinical cures	N	Cure rate (%)	Number of clinical cures	n
<i>Enterobacteriaceae</i>	80.9	93	115	92.7	115	124
<i>Citrobacter freundii</i> complex	62.5	5	8		0	0
<i>Enterobacter cloacae</i>	100	5	5	66.7	2	3
<i>Escherichia coli</i>	83.3	70	84	94.4	84	89
<i>Klebsiella oxytoca</i>	100	5	5	100	5	5
<i>Klebsiella pneumoniae</i>	82.1	23	28	88.6	31	35
<i>Proteus mirabilis</i>	66.7	2	3	100	5	5
<i>Pseudomonas aeruginosa</i>	82.4	14	17	85.0	17	20

Complicated urinary tract infections

A total of 1020 adults with documented complicated urinary tract infection (cUTI) (737 with acute pyelonephritis and 283 with cUTI without acute pyelonephritis) were randomised and received treatment in a phase III multicentre, double-blind, comparative study. Treatment was with either Ceftazidime-avibactam (2000 mg/500 mg) IV over 120 mins every 8 hours or doripenem 500 mg IV over 60 mins every 8 hours. There was an optional switch to oral therapy for patients who had clinical improvement as defined in the study protocol after a minimum of 5 days IV treatment. Total duration of antibiotic therapy (IV plus oral) was 10 days (optionally up to 14 if bacteraemic). The mMITT population included all patients with a confirmed cUTI diagnosis, received at least 1 dose of study treatment and had a study-qualifying pre-treatment urine culture containing 10⁵ CFU/mL of a Gram-negative pathogen and no more than 2 species of microorganisms. Any patient with a Gram-positive pathogen, or a bacterial species not expected to respond to both study drugs was excluded.

The primary efficacy endpoint was per-patient microbiological response at the TOC visit in the mMITT analysis set.

Table 8 Favourable per-patient microbiological response rate at TOC (RECAPTURE mMITT analysis set)

		CAZ-AVI (N=393)	Doripenem (N=417)	Difference (%) (95% CI)
Per patient microbiological response	Favourable	304 (77.4)	296 (71.0)	6.4 (0.33, 12.36)

Favourable microbiological response rates at TOC by pathogen in the mMITT population are shown in Table 9 below.

Table 9 Favourable per-pathogen microbiological response rate at TOC by common (combined frequency of ≥ 10) baseline pathogen (RECAPTURE mMITT)

Pathogen	Number of patients					
	CAZ-AVI (N=393)			Doripenem (N=417)		
	Favourable response rate (%)	Number of favourable responses	N	Favourable response rate (%)	Number of favourable responses	n
<i>Enterobacteriaceae</i>	78.3	299	382	70.6	281	398
<i>Enterobacter cloacae</i>	54.5	6	11	69.2	9	13
<i>Escherichia coli</i>	78.4	229	292	71.9	220	306
<i>Klebsiella pneumoniae</i>	75.0	33	44	62.5	35	56
<i>Proteus mirabilis</i>	94.1	16	17	69.2	9	13
<i>Pseudomonas aeruginosa</i>	66.7	12	18	75.0	15	20

Hospital-acquired pneumonia

A total of 808 adults with nosocomial pneumonia (35% with VAP) were randomised and received treatment in a phase III double-blind, comparative study of Ceftazidime-avibactam (2000 mg/500 mg) IV over 120 mins every 8 hours or meropenem 1g IV over 30 mins every 8 hours. Treatment duration was 7 to 14 days. The clinically modified intent to treat (cMITT) population included patients who met the minimum disease criteria, received at least 1 dose of study treatment and who had properly obtained baseline respiratory or blood cultures demonstrating Gram-negative pathogens excluding patients with monomicrobial Gram-negative infections with species not expected to respond to both study drugs (e.g. *Acinetobacter* species or *Stenotrophomonas* species). The cMITT also included patients in whom no etiologic pathogens were identified from respiratory or blood cultures at baseline. The CE at TOC analyses set was the clinically evaluable subset of the cMITT.

The primary efficacy endpoint was the clinical response at the TOC visit in the co-primary populations of the cMITT and CE at TOC. See Table 10 below.

Table 10 Clinical cure rates at TOC (REPROVE cMITT and CE at TOC analysis sets)

Number (%) of patients				
Analysis set	Response	CAZ-AVI	Meropenem	Difference (%) 95% CI
cMITT		(N=356)	(N=370)	
	Clinical cure	245 (68.8)	270 (73.0)	-4.2 (-10.76, 2.46)
CE at TOC		(N=257)	(N=270)	

Number (%) of patients				
Analysis set	Response	CAZ-AVI	Meropenem	Difference (%) 95% CI
	Clinical cure	199 (77.4)	211 (78.1)	-0.7 (-7.86, 6.39)

All-cause mortality rates at Day 28 (cMITT) was 8.4% (30/356) and 7.3% (27/370) ceftazidime-avibactam and meropenem treated patients, respectively.

Clinical cure rate and favourable microbiological response rate at TOC by pathogen in mMITT for Gram-negative aerobes are shown in Tables 11 and 12.

Table 11 Clinical cure rate at TOC by common (combined frequency of ≥ 10) Gram-negative baseline pathogen (REPROVE mMITT)

Pathogen	Number of patients					
	CAZ-AVI (N=171)			Meropenem (N=184)		
	Cure rate (%)	Number of clinical cures	N	Cure rate (%)	Number of clinical cures	n
<i>Enterobacteriaceae</i>	73.6	89	121	75.4	104	138
<i>Enterobacter aerogenes</i>	62.5	5	8	50.0	4	8
<i>Enterobacter cloacae</i>	92.3	24	26	54.5	12	22
<i>Escherichia coli</i>	64.7	11	17	75.0	15	20
<i>Klebsiella pneumoniae</i>	72.9	43	59	77.5	55	71
<i>Proteus mirabilis</i>	85.7	12	14	75.0	9	12
<i>Serratia marcescens</i>	73.3	11	15	92.3	12	13
<i>Pseudomonas aeruginosa</i>	60.3	35	58	74.5	35	47
<i>Haemophilus influenzae</i>	81.3	13	16	80.0	20	25

Table 12 Per-pathogen microbiological response at TOC by common (combined frequency of ≥ 10) Gram-negative baseline pathogen (REPROVE mMITT)

Pathogen	Number of patients					
	CAZ-AVI (N=171)			Meropenem (N=184)		
	Favourable response rate (%)	Number of favourable responses	N	Favourable response rate (%)	Number of favourable responses	n
<i>Enterobacteriaceae</i>						
<i>Enterobacter aerogenes</i>	62.5	5	8	62.5	5	8
<i>Enterobacter cloacae</i>	80.8	21	26	59.1	13	22
<i>Escherichia coli</i>	76.5	13	17	80.0	16	20
<i>Klebsiella pneumoniae</i>	62.7	37	59	74.6	53	71
<i>Proteus mirabilis</i>	78.6	11	14	66.7	8	12
<i>Serratia marcescens</i>	66.7	10	15	61.5	8	13

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
Trade Name: Zavicefta®
CDS Effective Date: February 28, 2024
Supersedes: November 18, 2021
Approved by BPOM: January 31, 2025

Table 12 Per-pathogen microbiological response at TOC by common (combined frequency of ≥ 10) Gram-negative baseline pathogen (REPROVE mMITT)

Pathogen	Number of patients					
	CAZ-AVI (N=171)			Meropenem (N=184)		
	Favourable response rate (%)	Number of favourable responses	N	Favourable response rate (%)	Number of favourable responses	n
<i>Pseudomonas aeruginosa</i>	37.9	22	58	38.3	18	47
<i>Haemophilus influenzae</i>	87.5	14	16	92.0	23	25

5.2 Pharmacokinetic properties

Distribution

The human protein binding of both ceftazidime and avibactam is low, approximately 10% and 8%, respectively. The steady-state volumes of distribution of ceftazidime and avibactam were comparable, about 22 L and 18 L, respectively in healthy adults following multiple doses of 2000 mg/500 mg ceftazidime-avibactam infused over 2 hours every 8 hours. Pharmacokinetic parameters of ceftazidime and avibactam following single and multiple dose administration of Zavicefta® were similar to those determined when ceftazidime or avibactam were administered alone. Both ceftazidime and avibactam penetrate into human bronchial epithelial lining fluid (ELF) to the same extent with concentrations around 30% that of plasma, and a similar concentration time profile between ELF and plasma.

Ceftazidime and avibactam plasma exposure were comparable across patients with different indications, cIAI, cUTI and NP.

Penetration of ceftazidime into the intact blood-brain barrier is poor, resulting in low levels of ceftazidime in the CSF in the absence of inflammation. However, concentrations of 4 to 20 mg/L or more are achieved in the CSF when the meninges are inflamed. Avibactam penetration of the blood brain barrier has not been studied clinically, however, in rabbits with inflamed meninges, CSF exposures of ceftazidime and avibactam were 43% and 38% of plasma AUC, respectively. For ceftazidime, concentrations in excess of the MIC of ceftazidime-avibactam for common pathogens can be achieved in tissues such as bone, heart, bile, sputum, aqueous humour, synovial, pleural and peritoneal fluids. Ceftazidime crosses the placenta readily, and is excreted in the breast milk. Avibactam penetrates into the subcutaneous tissue at the site of skin infections, with tissue concentrations approximately equal to free drug concentrations in plasma.

Biotransformation

Ceftazidime is not metabolized. No metabolism of avibactam was observed in human liver preparations (microsomes and hepatocytes). Unchanged avibactam was the major drug-related component in human plasma and urine following dosing with [¹⁴C]-avibactam.

Elimination

The terminal half-life ($t_{1/2}$) of both ceftazidime and avibactam is about 2 h after IV administration. Ceftazidime is excreted unchanged into the urine by glomerular filtration; approximately 80 - 90% of the dose is recovered in the urine within 24 h. Avibactam is

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium

Trade Name: Zavicefta®

CDS Effective Date: February 28, 2024

Supersedes: November 18, 2021

Approved by BPOM: January 31, 2025

excreted unchanged into the urine with a renal clearance of approximately 158 ml/min, suggesting active tubular secretion in addition to glomerular filtration; approximately 97% of the dose is recovered in the urine, 95% within 12 h. Less than 1% of ceftazidime is excreted via the bile and less than 0.25% of avibactam is excreted into faeces.

Linearity/non-linearity

The pharmacokinetics of both ceftazidime and avibactam are approximately linear across the dose range studied (50 mg to 2000 mg) for a single IV administration. No appreciable accumulation of ceftazidime or avibactam was observed following multiple IV infusions of 2000 mg/500 mg of ceftazidime-avibactam administered every 8 hours for up to 11 days in healthy adults with normal renal function.

Special populations

Patients with renal impairment

Elimination of ceftazidime and avibactam is decreased in patients with moderate or severe renal impairment, the average increases in avibactam AUC are 3.8-fold and 7-fold in subjects with moderate and severe renal impairment, and end stage renal disease including patients undergoing haemodialysis; the dose should be reduced in patients with CrCl ≤50 ml/min) (see section 4.2).

Patients with hepatic impairment

Mild to moderate hepatic impairment had no effect on the pharmacokinetics of ceftazidime in individuals administered 2 g IV every 8 hours for 5 days, provided renal function was not impaired. The pharmacokinetics of ceftazidime in patients with severe hepatic impairment has not been established. The pharmacokinetics of avibactam in patients with any degree of hepatic impairment has not been studied.

As ceftazidime and avibactam do not appear to undergo significant hepatic metabolism, the systemic clearance of either drug is not expected to be significantly altered by hepatic impairment. Therefore, no dosage adjustment of ceftazidime-avibactam is recommended for patients with hepatic impairment (see section 4.2).

Elderly patients

The reduced clearance observed in elderly patients was primarily due to age-related decrease in renal clearance of ceftazidime. The mean elimination half-life ranged from 3.5 to 4 hours following single or 7 days repeated every 12 hours dosing of 2 g IV bolus injections in elderly patients 80 years or older.

Following single dose IV administration of 500 mg avibactam as a 30-minute IV infusion, the elderly had a slower terminal half-life of avibactam, which may be attributed to age related decrease in renal clearance. Dosage adjustment for ceftazidime-avibactam is not required in elderly subjects (≥65 years of age) with CrCl >50 ml/min.

Paediatric patients

The safety and efficacy of Zavicefta® in paediatric patients (<18 years of age) have not been established.

Gender

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium

Trade Name: Zavicefta®

CDS Effective Date: February 28, 2024

Supersedes: November 18, 2021

Approved by BPOM: January 31, 2025

The pharmacokinetics of ceftazidime-avibactam was similar between males and females. No dose adjustment is required based on sex.

Race

Based on a population pharmacokinetic analysis, no dose adjustment of ceftazidime-avibactam is required based on race.

5.3 Preclinical safety data

Genetic toxicology

For ceftazidime a mouse Micronucleus test and an Ames test were both negative for mutagenic effects. Carcinogenicity studies have not been conducted. In genotoxicity assays with avibactam, there was no induction of gene mutation in the *in vitro* bacterial reverse mutation tests, nor were there any indications of genotoxicity in an *in vitro* unscheduled DNA synthesis test in rat liver cells or an *in vitro* micronucleus test in mouse lymphoma cells. In cultured human lymphocytes, statistically significant increases in chromosomal aberrations were observed under a single treatment condition (44 h harvest time, -S9). As these findings were not replicated in an independent study, the results are considered to be of limited biological relevance. When administered up to the limit dose of 2 g/kg IV, avibactam was negative in a rat *in vivo* micronucleus assay. Carcinogenicity studies have not been conducted. No genetic toxicology studies have been conducted on ceftazidime-avibactam.

Reproductive toxicology

Reproduction studies have been performed with ceftazidime in mice and rats at doses up to 40 times the human dose and have revealed no evidence of impaired fertility or harm to the fetus. In pregnant rabbits at exposures of avibactam approximately 8 fold higher than those observed in humans at 0.5 g three times daily there was a significant effect on maternal food consumption and a slight effect on fetal weight and slight retardation of ossification of a few bones in the fetus. In the rat, no adverse effects were observed on embryofetal development or fertility. Following administration of avibactam throughout pregnancy and lactation in the rat, there was no effect on pup survival, growth or development, however there was an increase in incidence of dilation of the renal pelvis and ureters in less than 10% of the rat pups at maternal exposures greater than or equal to approximately 1.5 times human therapeutic exposures. No reproductive toxicology studies have been conducted on ceftazidime-avibactam.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium carbonate

6.2 Incompatibilities

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

6.3 Shelf life

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium

Trade Name: Zavicefta[®]

CDS Effective Date: February 28, 2024

Supersedes: November 18, 2021

Approved by BPOM: January 31, 2025

After reconstitution:

The reconstituted vial should be used immediately.

After dilution:

Infusion bags:

If the intravenous solution is prepared with diluents listed in section 6.6 (ceftazidime concentration 8 mg/mL), the chemical and physical in-use stability has been demonstrated (from initial vial puncture) for up to 12 hours at 2-8°C, followed by up to 4 hours below 25°C.

If the intravenous solution is prepared with diluents listed in section 6.6 (ceftazidime concentration > 8 mg/mL to 40 mg/mL), the chemical and physical in-use stability has been demonstrated (from initial vial puncture) for up to 4 hours below 25°C.

From a microbiological point of view, the medicinal product should be used immediately, unless reconstitution and dilution have taken place in controlled and validated aseptic conditions. If not used immediately, in use storage times and conditions prior to use are the responsibility of the user and must not exceed those above.

6.4 Special precautions for storage

Store below 30°C.

Store in the original package in order to protect from light.

For storage conditions of the reconstituted and diluted medicinal product, see section 6.3.

6.5 Nature and contents of container

20 ml glass vial (Type 1) closed with a rubber (halobutyl) stopper and aluminium seal with flip-off cap.

The medicinal product is supplied in packs of 10 vials.

6.6 Special precautions for disposal and other handling

The powder must be reconstituted with sterile water for injections and the resulting concentrate must then be immediately diluted prior to use. The reconstituted solution is a pale yellow solution that is free of any particles.

Standard aseptic techniques should be used for solution preparation and administration.

1. Introduce the syringe needle through the vial closure and inject 10 ml of sterile water for injection.
2. Withdraw the needle and shake the vial to give a clear solution.
3. Do not insert a gas relief needle until the product has dissolved. Insert a gas relief needle through the vial closure to relieve the internal pressure.

Generic Name: Ceftazidime Pentahydrate, Avibactam Sodium
Trade Name: Zavicefta®
CDS Effective Date: February 28, 2024
Supersedes: November 18, 2021
Approved by BPOM: January 31, 2025

4. Transfer the entire contents (approximately 12.0 ml) of the resultant solution to an infusion bag immediately. Reduced doses may be achieved by transfer of an appropriate volume of the resultant solution to an infusion bag, based upon ceftazidime and avibactam content of 167.3 mg/ml and 41.8 mg/ml, respectively. A dose of 1000 mg/250 mg or 750 mg/187.5 mg is achieved with 6.0 ml or 4.5 ml aliquots, respectively.

Note: To preserve product sterility, it is important that the gas relief needle is not inserted through the vial closure before the product is dissolved.

Vials of ceftazidime-avibactam powder should be reconstituted with 10 ml of sterile water for injections, followed by shaking until the content dissolves. An infusion bag may contain any of the following: sodium chloride 9 mg/ml (0.9%) solution for injection, dextrose 50 mg/ml (5%) solution for injection, or Lactated Ringer's solution. A 100 ml infusion bag can be used to prepare the infusion, based on the patient's volume requirements. The total time interval between starting reconstitution and completing preparation of the intravenous infusion should not exceed 30 minutes.

Each vial is for single use only.

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER NAME AND ADDRESS

Imported by:
PT. Pfizer Indonesia.
Jakarta, Indonesia

Manufactured by:
ACS DOBFAR S.p.A
VIA A. Fleming, 2
Verona 37135
Italy

8. MARKETING AUTHORISATION NUMBER

Box, 10 vial @ 2.5 g; Reg. No.: DKI2043900344A1

HARUS DENGAN RESEP DOKTER

9. DATE OF REVISION OF THE TEXT

01/2025

Brosur kemasan: Informasi bagi pengguna

Zavicefta® 2 g/0,5 g Serbuk untuk konsentrat larutan infus Ceftazidim/Avibactam

Baca semua bagian brosur ini dengan cermat sebelum mulai menggunakan obat ini karena berisi informasi penting bagi Anda.

- Simpan brosur ini. Anda mungkin perlu membacanya kembali.
- Jika ada pertanyaan lebih lanjut, hubungi dokter, apoteker, atau perawat Anda.
- Obat ini telah diresepkan hanya untuk Anda. Jangan berikan kepada orang lain, karena dapat membahayakan mereka sekali pun tanda-tanda penyakit mereka sama dengan Anda.
- Jika Anda mengalami efek samping apa pun, konsultasikan dengan dokter atau perawat Anda. Ini termasuk segala bentuk efek samping yang tidak tercantum di dalam brosur ini. Lihat bagian 13.

Isi brosur ini:

1. Nama obat
2. Bentuk sediaan
3. Deskripsi obat
4. Apa kandungan obat ini?
5. Kekuatan obat
6. Apa kegunaan obat ini?
7. Berapa banyak dan seberapa sering Anda seharusnya menggunakan obat ini? Apa yang harus dilakukan jika ada dosis terlewat?
8. Kapan seharusnya Anda tidak menggunakan obat ini?
9. Apa yang perlu dipertimbangkan saat menggunakan obat ini?
10. Apa saja obat atau makanan lain yang harus dihindari selama menggunakan obat ini?
11. Apakah obat ini aman untuk perempuan hamil dan menyusui?
12. Apakah pasien diizinkan untuk mengemudi atau menjalankan mesin selama menggunakan obat ini?
13. Apakah potensi efek yang tidak diharapkan jika menggunakan obat ini?
14. Tanda-tanda dan gejala-gejala overdosis
15. Apa yang harus dilakukan jika Anda menggunakan dosis melebihi anjuran?
16. Bagaimana cara menyimpan obat ini?
17. Berapa lama umur simpan obat setelah kemasannya pertama kali dibuka?
18. Bagaimana cara melarutkan obat ini?
19. Nomor izin pemasaran
20. Nama dan alamat pemohon dan/atau pemilik obat sesuai dengan ketentuan yang berlaku
21. Tanggal revisi PIL
22. Peringatan khusus

1. Nama obat

Zavicefta®

2. Bentuk sediaan

Serbuk untuk larutan infus.

3. Deskripsi obat

Zavicefta® adalah bubuk steril berwarna putih hingga kuning untuk larutan infus.

4. Apa kandungan obat ini?

Zavicefta® merupakan obat antibiotik yang mengandung dua bahan aktif yaitu ceftazidim dan avibactam.

- Ceftazidim termasuk dalam kelompok antibiotik yang disebut “sefalosporin”. Bahan ini dapat membunuh banyak jenis bakteri.
- Avibactam adalah “penghambat beta-laktamase” yang membantu ceftazidim membunuh sebagian bakteri yang tidak dapat dibunuh oleh ceftazidim jika diberikan sendirian.

5. Kekuatan obat

2 g/0,5 g

6. Apa kegunaan obat ini?

Zavicefta® diindikasikan bagi orang dewasa untuk mengobati infeksi bakteri yang memproduksi ESBL (Beta Laktamase Spektrum Luas) dan resistansi *Pseudomonas aeruginosa* terhadap ceftazidime sebagai berikut:

- Infeksi Intraabdomen dengan Komplikasi (cIAI): memerlukan intervensi bedah dan meluas melampaui viskus berongga pada saluran usus ke dalam rongga peritoneal (laparotomi, drainase perkutan suatu abses, atau pembedahan laparoskopi). Untuk mengobati cIAI, digunakan dalam kombinasi dengan metronidazol.
- Infeksi Saluran Kemih dengan Komplikasi, termasuk Pielonefritis (cUTI): pielonefritis akut diindikasikan dengan nyeri panggul atau nyeri tekan sudut kostovertebral pada saat pemeriksaan dan setidaknya salah satu gejala berikut ini: demam, muntah, mual, atau infeksi saluran kemih bawah dengan komplikasi tanpa pielonefritis yang diindikasikan dengan gejala-gejala yang memperkuat (disuria, dorongan untuk buang air kecil, frekuensi buang air kecil, nyeri suprapubis, demam, mual, muntah) ditambah setidaknya 1 faktor komplikasi (riwayat retensi urin kronis, uropati obstruktif, abnormalitas fungsional atau anatomi saluran urogenital, penggunaan kateterisasi kandung kemih, prosedur urogenital).
- Pneumonia dapatan rumah sakit (HAP), termasuk pneumonia yang berhubungan dengan ventilator (VAP): pneumonia yang muncul setelah ≥ 48 jam di fasilitas rawat inap akut atau kronis atau yang berkembang dalam waktu 7 hari setelah keluar rumah sakit; infeksi parenkim paru-paru yang muncul ≥ 48 jam setelah intubasi endotrakea dan ventilasi mekanis.

Pertimbangan harus diberikan terhadap panduan resmi tentang penggunaan agen antibakteri yang tepat.

7. Berapa banyak dan seberapa sering Anda seharusnya menggunakan obat ini? Apa yang harus dilakukan jika ada dosis yang terlewat?

Dosis yang disarankan adalah satu vial (2 g ceftazidim dan 0,5 g avibactam), setiap 8 jam.

Obat ini diberikan melalui pembuluh darah vena – ini akan membutuhkan waktu sekitar 2 jam.

Rangkaian pengobatan biasanya membutuhkan 5 hingga 14 hari, bergantung pada jenis infeksi yang Anda alami dan respons Anda terhadap pengobatan.

Pasien dengan masalah ginjal

Jika Anda mengalami masalah ginjal, dokter dapat mengurangi dosis Anda. Hal ini karena Zavicefta® dikeluarkan dari tubuh Anda melalui ginjal.

Jika Anda merasa telah melewatkan satu dosis, segera beri tahu dokter atau perawat Anda.

8. Kapan seharusnya Anda tidak menggunakan obat ini?

Jangan menggunakan Zavicefta® jika:

- Anda alergi terhadap ceftazidim, avibactam, atau bahan lain yang terkandung dalam obat ini

Nama Generik: Avibactam Natrium, Ceftazidim Pentahidrat
Nama Dagang: Zavicefta®
Tanggal Efektif CDS: 28 Februari, 2024
Menggantikan: 18 November 2021
Disetujui oleh BPOM: 31 Januari 2025

- Anda alergi terhadap antibiotik sefalosporin
- Anda pernah mengalami reaksi alergi parah terhadap antibiotik lain yang termasuk kelompok penisilin atau karbapenem.

Jangan menggunakan Zavicefta® jika ada di antara kondisi di atas yang berlaku untuk Anda. Jika kurang yakin, konsultasikan dengan dokter atau perawat Anda sebelum menggunakan Zavicefta®.

9. Apa yang perlu dipertimbangkan saat menggunakan obat ini?

Konsultasikan dengan dokter atau perawat Anda sebelum menggunakan Zavicefta® jika:

- Anda pernah mengalami reaksi alergi (sekali pun hanya ruam kulit) terhadap antibiotik lain yang termasuk kelompok penisilin atau karbapenem
- Anda mempunyai masalah ginjal - dokter Anda mungkin akan memberi Anda dosis yang lebih rendah untuk memastikan Anda tidak diberi obat terlalu banyak. Ini dapat menyebabkan gejala seperti kejang (lihat bagian 13)

Jika ada di antara kondisi di atas pernah terjadi pada Anda (atau Anda kurang yakin), sampaikan kepada dokter atau perawat Anda sebelum menggunakan Zavicefta®.

Beri tahu dokter atau perawat Anda jika Anda menderita diare selama pengobatan Anda.

Infeksi lain

Ada kemungkinan kecil bahwa Anda dapat terkena infeksi yang berbeda yang disebabkan oleh bakteri lain selama atau setelah pengobatan dengan Zavicefta®. Ini termasuk kandidiasis (infeksi jamur di mulut atau area genital).

Tes lab

Beri tahu dokter Anda bahwa Anda menggunakan Zavicefta® jika Anda akan menjalani suatu tes. Ini karena Anda mungkin mendapatkan hasil yang tidak normal untuk tes yang disebut “DAGT” atau “Coombs”. Tes ini digunakan untuk menemukan antibodi yang melawan sel darah merah Anda.

Zavicefta® juga dapat berpengaruh pada hasil dari beberapa tes urine untuk gula. Beri tahu pengambil sampel Anda bahwa Anda telah diberi Zavicefta®.

Anak-anak dan remaja

Zavicefta® tidak boleh digunakan pada anak-anak dan remaja. Hal ini dikarenakan keamanan obat yang tidak diketahui pada kelompok usia ini.

Zavicefta® mengandung natrium

Untuk pasien dengan diet terkontrol natrium, setiap vial mengandung sekitar 148 mg natrium.

10. Apa saja obat atau makanan lain yang harus dihindari selama menggunakan obat ini?

Beri tahu dokter atau perawat Anda jika Anda sedang menggunakan, baru-baru ini menggunakan, atau mungkin akan menggunakan obat-obatan lain.

Beri tahu dokter Anda sebelum menggunakan Zavicefta® jika Anda sedang meminum obat-obatan berikut ini:

- antibiotik kloramfenikol
- sejenis antibiotik aminoglikosida – seperti gentamisin, tobramisin
- tablet air bernama furosemid
- obat untuk gout bernama probenesid

Nama Generik: Avibactam Natrium, Ceftazidim Pentahidrat
Nama Dagang: Zavicefta®
Tanggal Efektif CDS: 28 Februari, 2024
Menggantikan: 18 November 2021
Disetujui oleh BPOM: 31 Januari 2025

Beri tahu dokter Anda sebelum menggunakan Zavicefta® jika ada di antara kondisi di atas yang sama dengan Anda.

11. Apakah obat ini aman untuk perempuan hamil dan menyusui?

Jika Anda hamil atau sedang menyusui, menduga bahwa diri Anda sedang hamil, atau berencana untuk hamil, mintalah saran dokter Anda sebelum menggunakan obat ini.

12. Apakah pasien diizinkan untuk mengemudi atau menjalankan mesin selama menggunakan obat ini?

Zavicefta® dapat membuat Anda merasa pusing. Ini dapat memengaruhi kemampuan Anda dalam mengemudi, menggunakan alat atau mesin.

13. Apakah potensi efek yang tidak diharapkan jika menggunakan obat ini?

Seperti semua obat-obatan yang ada, obat ini dapat menimbulkan efek samping, meskipun tidak semua orang mengalaminya. Efek samping berikut dapat terjadi dengan obat ini:

Efek samping serius

Segera beri tahu dokter Anda jika Anda mengamati adanya efek samping serius berikut - Anda mungkin membutuhkan perawatan medis segera:

- reaksi alergi berat – tanda-tandanya termasuk pembengkakan mendadak pada bibir, wajah, tenggorokan, atau lidah, ruam parah atau reaksi kulit parah lain, kesulitan menelan atau bernapas, nyeri dada tiba-tiba (yang mungkin merupakan tanda sindrom Kounis). Reaksi-reaksi ini dapat mengancam nyawa.
 - diare yang semakin memburuk atau tidak berhenti, atau feses yang berdarah atau berlendir – ini dapat terjadi selama atau setelah pengobatan Zavicefta® dihentikan. Jika ini terjadi jangan mengonsumsi obat yang menghentikan atau memperlambat gerakan usus.
- Segera beri tahu dokter Anda jika teramati adanya efek samping serius di atas.

Efek samping lainnya

Beri tahu dokter atau perawat Anda jika teramati adanya efek samping berikut ini:

Sangat umum terjadi: (dapat terjadi pada lebih dari 1 di antara 10 orang)

- hasil yang tidak normal untuk tes yang disebut “DAGT” atau “Coombs”. Tes ini digunakan untuk menemukan antibodi yang melawan sel darah merah Anda. Ini mungkin dapat menyebabkan anemia (yang dapat membuat Anda merasa lelah) dan sakit kuning (kekuningan pada kulit dan mata)

Umum: (dapat dialami hingga 1 di antara 10 orang)

- infeksi jamur, termasuk pada mulut dan vagina
- perubahan jumlah pada jenis sel darah tertentu (“eosinofil” dan “trombosit”) – tampak dalam tes darah
- sakit kepala
- merasa pusing
- merasa mual atau mengalami muntah-muntah
- sakit perut
- diare
- peningkatan jumlah enzim tertentu yang dihasilkan oleh hati Anda - tampak dalam hasil tes darah
- ruam kulit yang gatal dan menebal (“urtikaria”)
- gatal-gatal
- kemerahan, nyeri, atau pembengkakan di tempat Zavicefta® diberikan ke dalam vena
- demam
- trombosis lokasi infus
- flebitis lokasi infus

Nama Generik: Avibactam Natrium, Ceftazidim Pentahidrat
Nama Dagang: Zavicefta®
Tanggal Efektif CDS: 28 Februari, 2024
Menggantikan: 18 November 2021
Disetujui oleh BPOM: 31 Januari 2025

Tidak umum: (dapat dialami hingga 1 di antara 100 orang)

- peningkatan jumlah pada jenis sel darah tertentu (“limfosit”) – tampak dalam tes darah
- penurunan jumlah pada jenis sel darah tertentu (“leukosit”) – tampak dalam tes darah
- kesemutan atau mati rasa
- rasa tidak enak pada mulut
- peningkatan jumlah zat tertentu di dalam darah (“kreatinin” dan “urea”). Ini menunjukkan seberapa baik ginjal Anda berfungsi.
- kolitis *Clostridium difficile*
- kolitis pseudomembranosa
- gagal ginjal akut

Sangat jarang: (dapat dialami hingga 1 di antara 10.000 orang)

- pembengkakan sebagian dari ginjal yang menyebabkan penurunan fungsi kerja normalnya

Tidak diketahui: (frekuensi tidak dapat diestimasi dari data yang tersedia)

- penurunan signifikan pada jenis sel darah putih yang digunakan untuk melawan infeksi – tampak dalam tes darah
- penurunan jumlah sel darah merah (anemia hemolitik) – tampak dalam tes darah
- reaksi alergi berat (lihat **Efek samping serius**, di atas)
- kekuningan pada bagian putih mata atau kulit
- timbulnya ruam berat atau lecet atau terkelupasnya kulit secara mendadak, yang dapat disertai demam tinggi atau nyeri persendian (ini mungkin merupakan tanda-tanda kondisi medis yang lebih serius seperti nekrolisis epidermal beracun, sindrom Stevens-Johnson, eritema multiform, atau kondisi yang dikenal sebagai DRESS (Drug Reaction with Eosinophilia and Systemic Symptoms/Reaksi obat dengan eosinofilia dan gejala sistemik)
- pembengkakan di bawah kulit, khususnya di bibir dan sekitar mata

Beri tahu dokter atau perawat Anda jika teramati adanya efek samping yang tercantum di atas. Untuk melaporkan efek samping, hubungi www.pfizersafetyreporting.com atau email di IDN.AEReporting@pfizer.com.

14. Tanda dan gejala kelebihan dosis

Overdosis dengan Zavicefta® dapat menyebabkan sekuelae neurologis termasuk ensefalopati, konvulsi, dan koma, akibat komponen ceftazidim.

15. Apa yang harus dilakukan jika Anda menggunakan dosis melebihi anjuran?

Zavicefta® akan diberikan kepada Anda oleh dokter atau perawat, jadi kecil kemungkinan terjadi kesalahan pemberian dosis. Namun, jika Anda mengalami efek samping atau beranggapan telah diberi terlalu banyak Zavicefta®, segera beri tahu dokter atau perawat Anda. Jika Anda menggunakan terlalu banyak Zavicefta® itu dapat berpengaruh pada otak dan menyebabkan kejang atau koma.

16. Bagaimana cara menyimpan obat ini?

Jauhkan obat ini dari penglihatan dan jangkauan anak-anak.

Jangan gunakan obat ini setelah melewati tanggal kedaluwarsa yang tertera pada wadahnya. Tanggal kedaluwarsa mengacu pada hari terakhir bulan tersebut.

Simpan di bawah suhu 30 °C.

Simpan dalam kemasan aslinya untuk melindunginya dari cahaya.

Nama Generik: Avibactam Natrium, Ceftazidim Pentahidrat
Nama Dagang: Zavicefta®
Tanggal Efektif CDS: 28 Februari, 2024
Menggantikan: 18 November 2021
Disetujui oleh BPOM: 31 Januari 2025

Jangan membuang obat melalui saluran pembuangan air atau bersama sampah rumah tangga. Tanyakan kepada apoteker cara membuang obat yang sudah tidak dibutuhkan. Langkah-langkah ini akan membantu melindungi lingkungan.

17. Berapa lama umur simpan obat setelah kemasannya pertama kali dibuka?

Setelah rekonstitusi:

Vial yang direkonstitusi harus segera digunakan.

Setelah pengenceran:

Kantong Infus:

Jika larutan intravena disiapkan dengan larutan untuk injeksi natrium klorida 9 mg/mL, larutan untuk injeksi dekstrosa 50 mg/mL, atau larutan Ringer Laktat (ceftazidim dengan konsentrasi 8 mg/mL), stabilitas kimia dan fisika selama penggunaan telah terbukti (sejak vial pertama kali ditusuk) hingga maksimal 12 jam pada suhu 2–8 °C, diikuti dengan maksimal 4 jam dibawah 25°C.

Jika larutan intravena disiapkan dengan larutan untuk injeksi natrium klorida 9 mg/mL, larutan untuk injeksi dekstrosa 50 mg/mL, atau larutan Ringer Laktat (ceftazidim dengan konsentrasi > 8 mg/mL hingga 40 mg/mL), stabilitas kimia dan fisika selama penggunaan telah terbukti (sejak vial pertama kali ditusuk) hingga maksimal 4 jam dibawah 25°C.

Dari sudut pandang mikrobiologi, produk medisinal harus digunakan dengan segera, kecuali telah dilakukan rekonstitusi dan pengenceran dalam kondisi aseptik terkontrol dan tervalidasi. Jika tidak segera digunakan, waktu penyimpanan selama penggunaan dan kondisi sebelum penggunaan menjadi tanggung jawab pengguna dan tidak boleh melebihi yang disebutkan di atas.

18. Bagaimana cara melarutkan obat ini?

Serbuk harus direkonstitusi dengan air steril untuk injeksi dan konsentrat yang dihasilkan harus segera diencerkan sebelum digunakan.

19. Nomor izin pemasaran

Dus, 10 vial @ 2,5 g; No. Reg.: DK12043900344A1

20. Nama dan alamat pemohon dan/atau pemilik obat sesuai dengan ketentuan yang berlaku

Diimpor oleh:

PT. Pfizer Indonesia.

Jakarta, Indonesia

Diproduksi oleh:

ACS DOBFAR S.p.A

VIA A. Fleming, 2

Verona 37135

Italia

21. Tanggal revisi PIL

01/2025

22. Peringatan khusus

HARUS DENGAN RESEP DOKTER

Nama Generik: Avibactam Natrium, Ceftazidim Pentahidrat
Nama Dagang: Zavicefta®
Tanggal Efektif CDS: 28 Februari, 2024
Menggantikan: 18 November 2021
Disetujui oleh BPOM: 31 Januari 2025

Informasi berikut ini ditujukan untuk tenaga kesehatan profesional saja:

Penting: Harap melihat Informasi Produk lengkap sebelum meresepkan.

Teknik aseptik harus diikuti dalam menyiapkan larutan infus. Isi dari vial Zavicefta® harus direkonstitusi dengan 10 mL air steril untuk injeksi. Petunjuk untuk rekonstitusi vial Zavicefta® dirangkum di bawah ini:

Kekuatan dosis ceftazidim/avibactam (mg)	Volume pengencer untuk ditambahkan (mL)	Perkiraan konsentrasi ceftazidim/avibactam (mg/mL)	Jumlah untuk diambil
2000/500	10	167,3/41,8	Volume total

1. Tusukkan jarum suntik menembus penutup vial dan suntikkan 10 mL air steril untuk injeksi.
2. Tarik jarum dan kocok vial hingga larutan jernih.
3. Jangan tusukkan jarum pelepas gas sebelum produk terlarut. Tusukkan jarum pelepas gas melalui penutup vial untuk melepaskan tekanan di dalam vial.
4. Segera pindahkan seluruh isi (sekitar 12,0 mL) larutan hasil ke kantong infus. Dosis yang dikurangi bisa didapatkan dengan memindahkan jumlah yang sesuai dari larutan hasil ke kantong infus, berdasarkan kandungan ceftazidim dan avibactam masing-masing sebesar 167,3 mg/mL dan 41,8 mg/mL. Dosis sebesar 1000 mg/250 mg atau 750 mg/187,5 mg masing-masing dicapai dengan alikuot 6,0 mL atau 4,5 mL.

Catatan: untuk menjaga kesterilan produk, penting bahwa jarum pelepas gas tidak ditusukkan ke penutup vial sebelum produk terlarut.

Larutan yang direkonstitusi harus diencerkan lebih lanjut untuk menghasilkan larutan Zavicefta® untuk infus. Kantong infus 100 mL dapat digunakan untuk menyiapkan infus, berdasarkan kebutuhan volume pasien. Pengencer infus yang sesuai termasuk: larutan injeksi natrium klorida 9 mg/mL (0,9%), larutan injeksi dekstrosa 50 mg/mL (5%) atau larutan Ringer Laktat. Larutan yang dihasilkan harus diberikan selama 120 menit.

Waktu rekonstitusi kurang dari 2 menit. Kocok dengan perlahan untuk merekonstitusi dan periksa apakah isinya sudah terlarut sepenuhnya. Interval waktu total antara dimulainya rekonstitusi hingga selesainya penyiapan infus intravena tidak boleh melebihi 30 menit. Produk obat parenteral harus diperiksa secara visual untuk melihat adanya partikulat sebelum diberikan.

Larutan infus Zavicefta® berwarna kuning pucat dan bebas partikel.

Dari sudut pandang mikrobiologi, produk medisinal harus digunakan dengan segera, kecuali telah dilakukan rekonstitusi dan pengenceran dalam kondisi aseptik terkontrol dan tervalidasi. Jika tidak segera digunakan, waktu penyimpanan selama penggunaan dan kondisi sebelum penggunaan menjadi tanggung jawab pengguna dan tidak boleh melebihi yang disebutkan di atas.

Kompatibilitas Zavicefta® dengan obat-obatan lain belum diketahui. Zavicefta® tidak boleh dicampur dengan atau ditambahkan secara fisik ke larutan yang mengandung produk obat lain.

Setiap vial hanya untuk sekali pakai.

Setiap produk yang tidak digunakan atau bahan limbah harus dibuang sesuai dengan persyaratan setempat.