PRIMPERAN
METOCLOPRAMIDE HCl
Pengatur Fungsi Cerna

Primperan adalah suatu obat yang tersendiri dalam sifat-sifat kimiawi, farmakologi dan terapi. Zat aktifnya yaitu: Metoclopramide.
Primperan memiliki cara kerja khusus pada berbagai kelainan pencernaan yang umum dijumpai dalam bidang kedokteran; mual (nausea), muntah-muntah (vomiting); perasaan kenyang (sensation of heaviness), kembung (meteorism), tidak enak pada ulu hati (epigastric discomfort) atau kehilangan nafsu makan (anorexia), perut merasa tidak enak atau sakit (abdominal discomfort or pain), ceklukan (hiccup) dan sakit kepala karena gangguan pencernaan.

KOMPOSISI :
Tiap 5 ml sirup mengandung Metoclopramide HCl Monohydrate 5,91 mg setara dengan Metoclopramide 5 mg

Bahan tambahan : Metilparaben, Propilparaben, Etil Maltol, Sukralosa, Asam sitrat, Natrium sitrat, Hidroksietil selulosa, FD&C Yellow No. 5, Perisa anggur, Aqua purificata

Primperan sirup merupakan cairan berwarna kuning, bau anggur.

INDIKASI :
- Meringankan/mengurangi gastroparesis akut dan yang kambuh kembali.
- Menghilangkan rasa panas sehubungan dengan reflux esophagitis.
- Menanggulangi mual dan muntah metabolik karena obat atau sesudah operasi.

KONTRA INDIKASI :
Penderita epilepsi, keadaan perdarahan gastrointestinal, obstruksi mekanik atau perforasi, pheochromocytomas.

PERINGATAN DAN PERHATIAN :
Takaran harap dikurangi pada penderita dengan gangguan renal karena dapat meningkatkan gejala extrapiramidal.

EFEK SAMPING :
Pada takaran tinggi dapat menimbulkan pusing, kegelisahan, lelah yang berlebihan, gejala extrapiramidal.

TAKARAN DAN CARA PEMAKAIAN :
Karena Primperan hanya dapat diperoleh melalui resep, takaran yang tercantum di bawah ini diberikan hanyalah sebagai pedoman:

DEWASA :
Sehari tiga kali 1-2 sendok teh (1 sendok teh = 5 ml = 5 mg), 30 menit sebelum makan dan sebelum tidur.

ANAK-ANAK :
0,1 mg per kg berat badan/hari, dibagi dalam beberapa dosis. Diberikan 30 menit sebelum makan dan sebelum tidur.

Primperan dapat dipakai bersama-sama dengan semua obat yang umum dipakai pada gastro-enterologi. Primperan tidak mempengaruhi tekanan darah maupun jantung bila diberikan bersamaan dengan komponen digitalis. Primperan tidak mempengaruhi fungsi ginjal dan sama sekali tidak menyebabkan retensi.

KEMASAN :
Primperan sirup (5 mg per 5 ml) dalam botol @ 50 ml
No. Reg. DKL8924208337A1

Jauhkan obat ini dari jangkauan anak-anak
Harus dengan resep dokter

SIMPAN DI BAWAH SUHU 30˚C
TERLINDUNG DARI CAHAYA

Diproduksi oleh :
PT. SOHO INDUSTRI PHARMASI, JAKARTA - INDONESIA

xxxxxx
x/xxxx
xxxx

PRIMPERAN
METOCLOPRAMIDE HCl
Gastro Intestinal Tract Regulator

Primperan is a drug which is original in its chemical, pharmacological and therapeutic properties. Its active ingredients is Metoclopramide. Primperan resides in the elective character of its actions in various digestive manifestations of nausea, vomiting, sensation of heaviness, meteorism, epigastric discomfort or pains, anorexia, abdominal discomfort of pains, hiccup and headaches associated with disorders of digestion.

COMPOSITION :
Each 5 ml syrup contains Metoclopramide HCl Monohydrate 5.91 mg equal to Metoclopramide 5 mg

Excipients : Methylparaben, Propylparaben, Ethyl maltol, Sucralose, Citric acid, Sodium citrate, Hydroxyethylcellulose, FD&C Yellow No. 5, Grape flavor, Aqua purificata

Primperan syrup is clear and yellowish solution, with grape odor.

INDICATIONS :
- To relieve acute and recurrent gastroparesis.
- Heartburn associated with reflux esophagitis.
- Anaesthetic or post operative nausea and vomiting.

CONTRA INDICATIONS :
- Epilepsi.
- Gastrointestinal bleeding, mechanic obstruction or perforation.
- Pheochromocytoma.

PRECAUTION :
Dosage should be reduced in patients with renal distrubance as extrapiramidal symptoms may be enhanced.

ADVERSE REACTIONS :
Large doses may cause headache, anxiety, fatique, extrapyramidal symptoms.

DOSAGE AND DIRECTIONS FOR USE :
Primperan being a medically prescribed product, the doses mentioned below are merely given as a recommendation :

ADULTS :
1 to 2 teaspoonful three times a day (1 teaspoonful = 5 ml = 5 mg), 30 minutes before meals and before bedtime.

CHILDREN :
0.1 mg per kg of body weight a day, in several divided doses. To be given 30 minutes before meals and before bedtime.

Primperan may be associated with all classical drugs use in gastro-enterology. It does affect neither blood pressure nor the heart, if given simultaneously with digitalic compound. It has no action on the renal function and does not cause retention of any sort.

AVAILABILITY :
Primperan syrup of 5 mg (a teaspoonfull) in bottles of 50 ml
Reg. No. DKL8924208337A1

Keep out of reach of children
With doctor's-prescription only

STORE BELOW 30˚C
PROTECT FROM LIGHT

Manufactured by :
PT. SOHO INDUSTRI PHARMASI, JAKARTA - INDONESIA

SOHO Global Health

PACKAGING DEVELOPMENT - R&D DIVISION

Part Description	**Leaflet Primperan Syrup-2F**		
Kode Pencetakan	**-**		
Supplier	**-**		
Material	**HVS 60 gsm**		
Dimension	**155 x 140 mm**		
Type Folded	**B1 - B2**		
Color	**P Black U** ■		

Finishing	**Lebar leaflet dibagi 4 bagian**	
Software used for TD	**Adobe Illustrator CC**	
Draft by / Date	**Andri S/ 03-03-23**	
FONT	**Jenis Font**	**Size**
Brand Name :	**Arial Bold**	**18,459 pt**
Generik Brand Name :	**Arial Bold**	**14,767 pt**
Komposisi, dll :	**Arial**	**6 pt**

DISETUJUI OLEH BPOM : 04/01/2024

ID : EREG100061VR12300038