


SUTENT

DESCRIPTION

SUTENT[®], an oral multi-kinase inhibitor targeting several receptor tyrosine kinases (RTK), is the malate salt of sunitinib. Sunitinib malate is described chemically as Butanedioic acid, hydroxy-, (2S)-, compound with *N*-[2-(diethylamino)ethyl]-5-[(*Z*)-(5-fluoro-1,2-dihydro-2-oxo-3*H*-indol-3-ylidene)methyl]-2,4-dimethyl-1*H*-pyrrole-3-carboxamide (1:1). The molecular formula is $C_{22}H_{27}FN_4O_2 \cdot C_4H_6O_5$ and the molecular weight is 532.6 Daltons. The chemical structure of sunitinib malate is:



Sunitinib malate is a yellow to orange powder with a pKa of 8.95. The solubility of sunitinib malate in aqueous media over the range pH 1.2 to pH 6.8 is in excess of 25 mg/mL. The log of the distribution coefficient (octanol/water) at pH 7 is 5.2.

SUTENT[®] (sunitinib malate) capsules are supplied as printed hard shell capsules containing sunitinib malate equivalent to 12.5 mg, 25 mg or 50 mg of sunitinib together with mannitol, croscarmellose sodium, povidone (K-25) and magnesium stearate as inactive ingredients.

The orange gelatin capsule shells contain titanium dioxide, and red iron oxide. The caramel gelatin capsule shells also contain yellow iron oxide and black iron oxide. The printing ink contains shellac, propylene glycol, sodium hydroxide, povidone and titanium dioxide.

CLINICAL PHARMACOLOGY

Mechanism of Action

Sunitinib malate is a small molecule that inhibits multiple RTKs, some of which are implicated in tumor growth, pathologic angiogenesis, and metastatic progression of cancer. Sunitinib was evaluated for its inhibitory activity against a variety of kinases (>80 kinases) and was identified as an inhibitor of platelet-derived growth factor receptors (PDGFR α and PDGFR β), vascular endothelial growth factor receptors (VEGFR1, VEGFR2 and VEGFR3), stem cell factor receptor (KIT), Fms-like tyrosine kinase-3 (FLT3), colony stimulating factor receptor Type 1 (CSF-1R), and the glial cellline derived neurotrophic factor receptor (RET). Sunitinib inhibition of the activity of these RTKs has been demonstrated in biochemical and cellular assays, and inhibition of function has been demonstrated in cell proliferation assays. The primary metabolite exhibits similar potency compared to sunitinib in biochemical and cellular assays.

Sunitinib inhibited the phosphorylation of multiple RTKs (PDGFR β , VEGFR2, KIT) in tumor xenografts expressing RTK targets *in vivo* and demonstrated inhibition of tumor growth or tumor regression and/or inhibited metastases in some experimental models of cancer. Sunitinib demonstrated the ability to inhibit growth of tumor cells expressing dysregulated target RTKs (PDGFR, RET, or KIT) *in vitro* and to inhibit PDGFR β - and VEGFR2-dependent tumor angiogenesis *in vivo*.

Pharmacokinetics

The pharmacokinetics of sunitinib and sunitinib malate have been evaluated in 135 healthy volunteers and in 266 patients with solid tumors.

Absorption, Distribution, Metabolism, and Elimination

Maximum plasma concentrations ($C_{\max}$) of sunitinib are generally observed between 6 and 12 hours ($T_{\max}$) following oral administration. Food has no effect on the bioavailability of sunitinib. Sunitinib may be taken with or without food.

Binding of sunitinib and its primary metabolite to human plasma protein *in vitro* was 95% and 90%, respectively, with no concentration dependence in the range of 100 – 4000 ng/mL. The apparent volume of distribution (V_d/F) for sunitinib was 2230 L. In the dosing range of 25 - 100 mg, the area under the plasma concentration-time curve (AUC) and $C_{\max}$ increase proportionately with dose.

Sunitinib is metabolized primarily by the cytochrome P450 enzyme, CYP3A4, to produce its primary active metabolite, which is further metabolized by CYP3A4. The primary active metabolite comprises 23 to 37% of the total exposure.

In-vitro studies indicate that sunitinib neither induces nor inhibits major CYP enzymes, including CYP3A4 (See section Drug-Drug Interaction)

The calculated *in vitro* K_i values for all CYP isoforms tested (CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, CYP3A4/5 AND CYP4A9/11) indicated that sunitinib and its primary active metabolite are unlikely to have any clinically relevant drug-drug interactions with drugs that may be metabolized by these enzymes.

Excretion is primarily via feces. In a human mass balance study of [^{14}C] sunitinib, 61% of the dose was eliminated in feces, with renal elimination accounting for 16% of the administered dose. Sunitinib and its primary active metabolite were the major drug-related compounds identified in plasma, urine, and feces, representing 91.5%, 86.4% and 73.8% of radioactivity in pooled samples, respectively. Minor metabolites were identified in urine and feces but generally not found in plasma. Total oral clearance (CL/F) ranged from 34 to 62 L/hr with an inter-patient variability of 40%.

Following administration of a single oral dose in healthy volunteers, the terminal half-lives of sunitinib and its primary active metabolite are approximately 40 to 60 hours and 80 to 110 hours, respectively. With repeated daily administration, sunitinib accumulates 3- to 4-fold while the primary metabolite accumulates 7- to 10-fold. Steady-state concentrations of sunitinib and its

primary active metabolite are achieved within 10 to 14 days. By Day 14, combined plasma concentrations of sunitinib and its active metabolite ranged from 62.9 – 101 ng/mL. No significant changes in the pharmacokinetics of sunitinib or the primary active metabolite were observed with repeated daily administration or with repeated cycles in the dosing regimens tested.

The pharmacokinetics were similar in healthy volunteers and in the solid tumor patient populations tested, including patients with gastrointestinal stromal tumor (GIST) and metastatic renal cell carcinoma (MRCC) (see CLINICAL STUDIES).

Special Populations

Population pharmacokinetic analyses of demographic data indicate that there are no clinically relevant effects of age, body weight, creatinine clearance, race, gender or ECOG score on the pharmacokinetics of SUTENT or the active metabolite.

The pharmacokinetics of sunitinib have not been evaluated in pediatric patients.

Hepatic Insufficiency

Sunitinib and its primary metabolite are mainly metabolized by the liver. Systemic exposures after a single dose of sunitinib were similar in subjects with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment compared to subjects with normal hepatic function. Sunitinib was not studied in subjects with severe (Child-Pugh Class C) hepatic impairment.

Renal Insufficiency

No clinical studies were conducted in patients with impaired renal function. Studies that were conducted excluded patients with serum creatinine > 2.0 x ULN. Population pharmacokinetic analyses have shown that sunitinib pharmacokinetics were unaltered in patients with calculated creatinine clearances in the range of 42 –347 mL/min.

Cardiac Electrophysiology

QT interval prolongation was investigated in a Phase 1 trial with 24 evaluable patients, aged 20-87 years, with advanced malignancies. At therapeutic plasma concentrations, the maximum QTcF mean change from baseline was 9.6 msec (90% CI 15.1 msec). At approximately twice the therapeutic concentrations, the maximum QTcF mean change from baseline was 15.4 msec (90% CI 22.4 msec). Moxifloxacin (400 mg) used as a positive control showed a 5.6 msec maximum mean QTcF change from baseline. No subjects experienced an effect on the QTc interval greater than Grade 2 (CTCAE v. 3.0). No patient presented with a cardiac arrhythmia (See section WARNING).

Drug-Drug Interactions

In vitro studies indicate that sunitinib does not induce or inhibit major CYP enzymes.

In Vitro Studies of CYP Inhibition and Induction: The *in vitro* studies in human liver

microsomes and hepatocytes of the activity of CYP isoforms CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, CYP3A4/5, and CYP4A9/11 indicated that sunitinib and its primary active metabolite are unlikely to have any clinically relevant drug-drug interactions with drugs that may be metabolized by these enzymes.

CYP3A4 Inhibitors: Concurrent administration of sunitinib malate with the strong CYP3A4 inhibitor, ketoconazole, resulted in 49% and 51% increases in the combined (sunitinib + primary active metabolite) $C_{\max}$ and $AUC_{0-\infty}$ values, respectively, after a single dose of sunitinib malate in healthy volunteers. A dose reduction for SUTENT should be considered when it must be co-administered with strong CYP3A4 inhibitors (see DOSAGE AND ADMINISTRATION). Administration of sunitinib with strong inhibitors of the CYP3A4 family (e.g. ritonavir, itraconazole, erythromycin, clarithromycin, grapefruit juice) may increase sunitinib concentrations. Concomitant administration with inhibitors should therefore be avoided, or the selection of an alternate concomitant medication with no, or minimal potential to inhibit CYP3A4 should be considered. If this is not possible, the dosage of sunitinib may need to be reduced. (See DOSAGE AND ADMINISTRATION)

CYP3A4 Inducers: Concurrent administration of SUTENT with the strong CYP3A4 inducer, rifampin, resulted in a 23% and 46% reduction in the combined (sunitinib + primary active metabolite) $C_{\max}$ and $AUC_{0-\infty}$ values, respectively, after a single dose of SUTENT in healthy volunteers. A dose increase for SUTENT should be considered when it must be co-administered with CYP3A4 inducers (see DOSAGE AND ADMINISTRATION). Administration of sunitinib with strong inducers of the CYP3A4 family (e.g., dexamethasone, phenytoin, carbamazepine, rifampin, phenobarbital or Hypericum perforatum also known as St. John's Wort) may decrease sunitinib concentrations. Concomitant administration with inducers should therefore be avoided, or selection of an alternate concomitant medication with no, or minimal potential to induce CYP3A4 should be considered. If this is not possible, the dosage of sunitinib may need to be increased. (see DOSAGE AND ADMINISTRATION)

CLINICAL STUDIES

The clinical safety and efficacy of SUTENT have been studied in patients with gastrointestinal stromal tumor (GIST) after progression on or intolerance to imatinib mesylate, and in patients with metastatic renal cell carcinoma (MRCC) after failure of cytokine-based therapy.

Efficacy is based on time to tumor progression (TTP) and an increase in survival in GIST.

Efficacy is based on progression-free survival (PFS) and objective response rates (ORR) for treatment-naïve and cytokine-refractory MRCC, respectively.

Gastrointestinal Stromal Tumor (GIST)

Study A

Study A was a two-arm, international, randomized, double-blind, placebo-controlled trial of SUTENT in patients with GIST who had disease progression during prior imatinib mesylate (imatinib) treatment or who were intolerant of imatinib. The primary objective was to compare

time-to-tumor progression (TTP) in patients receiving SUTENT plus best supportive care versus patients receiving placebo plus best supportive care. Secondary objectives included progression-free survival (PFS), objective response rate (ORR), and overall survival (OS). Patients were randomized (2:1) to receive either 50 mg SUTENT or placebo orally, once daily, on a schedule of 4 weeks on treatment followed by 2 weeks off (Schedule 4/2) until disease progression or withdrawal from the study for another reason. Treatment was unblinded at the time of disease progression. Patients randomized to placebo were then offered crossover to open-label SUTENT, and patients randomized to SUTENT were permitted to continue treatment per investigator judgment.

The intent-to-treat (ITT) population included 312 patients. Two-hundred seven patients were randomized to the SUTENT arm, and 105 patients were randomized to the placebo arm. Baseline age, gender, race and ECOG performance status were comparable between the placebo and SUTENT groups. Prior exposure to imatinib was similar between the two study arms. Demographics and patient characteristics are shown in Table 1.

Table 1. Baseline Demographic in Study A

	SUTENT (N=207)	Placebo (N=105)
Gender [N(%)]		
Male	132 (64)	64 (61)
Female	75 (36)	41 (39)
Self-identified Race [N(%)]		
White	183 (88)	92 (88)
Asian	10 (5)	5 (5)
Black	8 (4)	4 (4)
Not reported	6 (3)	4 (4)
Age Group [N (%)]		
< 65 years	143 (69)	76 (72)
≥ 65 years	64 (31)	29 (28)
Performance Status [N (%)]		
0	92 (44)	48 (46)
1	113 (55)	55 (52)
2	2 (1)	2 (2)
Prior Treatment [N (%)]		
Surgery (other than biopsy)	194 (94)	98 (93)
Radiotherapy	16 (8)	16 (15)

Imatinib outcome [N (%)]		
Intolerance	9 (4)	4 (4)
Progression within 6 months	36 (17)	17 (16)
Progression beyond 6 months	162 (78)	84 (80)

A planned interim efficacy and safety analysis was performed after 149 TTP events had occurred. There was a statistically significant advantage for SUTENT over placebo in the primary endpoint of TTP, as well as in the secondary endpoint of progression-free survival. Data were not mature enough to determine the overall survival benefit. Efficacy results are summarized in Table 2.

Table 2. GIST Efficacy Results (interim analysis)

Efficacy Parameter	Study A			
	SUTENT (N = 207)	Placebo (N = 105)	P-value (log-rank test)	HR (95% CI)
Time to Tumor Progression ^a [median, weeks (95 %CI)]	27.3 (16.0, 32.1)	6.4 (4.4, 10.0)	< 0.0001*	0.33 (0.23, 0.47)
Progression Free Survival ^b [median, weeks (95 %CI)]	24.1 (11.1, 28.3)	6.0 (4.4, 9.9)	< 0.0001*	0.33 (0.24, 0.47)
Objective Response Rate (PR) [%; (95% CI)]	6.8 (3.7, 11.1)	0	0.006 ^c	

CI=Confidence interval, HR=Hazard ratio, PR=Partial response

* A comparison is considered statistically significant if the p-value is < 0.0042 (O'Brien Fleming stopping boundary)

^a Time from randomization to progression; deaths prior to documented progression were censored at time of last radiographic evaluation

^b Time from randomization to progression or death due to any cause

^c Pearson chi-square

Figure 1. Kaplan-Meier Curve of TTP in Study A (Intent-to-Treat Population)

The baseline malignancy and prior treatment history of the patients were comparable between Studies 1 and 2. Across the two studies, 95% of the pooled population of patients had at least some component of clear-cell histology. All patients in Study 1 were required to have a histological clear-cell component. Most patients enrolled in the studies (97% of the pooled population) had undergone nephrectomy; prior nephrectomy was required for patients enrolled in Study 1. All patients had received one previous cytokine regimen. Metastatic disease present at the time of study entry included lung metastases in 81% of patients. Liver metastases were more common in Study 1 (27% vs. 16% in Study 2) and bone metastases were more common in Study 2 (51% vs. 25% in Study 1); 52% of patients in the pooled population had at least 3 metastatic

sites. Patients with known brain metastases or leptomeningeal disease were excluded from both studies.

Results of Studies 1 and 2

The ORR and DR data from Studies 1 and 2 are provided in Table 3.

Table 3. MRCC Efficacy Results

Efficacy Parameter	Study 1 (N = 106)	Study 2 (N = 63)
Objective Response Rate (PR) [% (95% CI)]	25.5 ¹ (17.5, 34.9)	36.5 ² (24.7, 49.6)
Duration of Response [median, weeks (95% CI)]	27.1 (24.4, *)	54 (34.3, 70.1)

CI=Confidence interval, PR=Partial response

¹ Assessed by blinded core radiology laboratory

² Assessed by investigators

* Data not mature enough to determine upper confidence limit

There were 27 PRs in Study 1 as assessed by a core radiology laboratory for an ORR of 25.5% (95% CI 17.5, 34.9). There were 23 PRs in Study 2 as assessed by the investigators for an ORR of 36.5% (95% CI 24.7-49.6). The majority (>90%) of objective disease responses were observed during the first four cycles; the latest reported response was observed in cycle 10. DR data from Study 1 is premature as only 4 of 27 patients (15%) responding to treatment had experienced disease progression. At the time of the data cutoff, Study 1 was ongoing with 44 of 106 patients (41.5%) continuing treatment, and 11 of the 63 patients (17.5%) enrolled on Study 2 continued to receive SUTENT[®] on continuation protocols.

INDICATIONS AND USAGE

SUTENT[®] is indicated for the treatment of gastrointestinal stromal tumor after disease progression on or intolerance to imatinib mesylate.

SUTENT[®] is indicated for the treatment of advanced renal cell carcinoma. Approval for advanced renal cell carcinoma is based on partial response rates and duration of responses. There are no randomized trials of SUTENT[®] demonstrating clinical benefit such as increased survival or improvement in disease-related symptoms in renal cell carcinoma.

CONTRAINDICATIONS

Use of SUTENT[®] is contraindicated in patients with hypersensitivity to sunitinib malate or to any other component of SUTENT[®].

WARNINGS

Pregnancy Category D

Sunitinib was evaluated in pregnant rats (0.3, 1.5, 3.0, 5.0 mg/kg/day) and rabbits (0.5, 1, 5, 20 mg/kg/day) for effects on the embryo. Significant increases in the incidence of embryoletality and structural abnormalities were observed in rats at the dose of 5 mg/kg/day (approximately 5.5 times the systemic exposure in patients administered the recommended daily doses [RDD]). Significantly increased embryoletality was observed in rabbits at 5 mg/kg/day while developmental effects were observed at =1 mg/kg/day (approximately 0.3 times the AUC in patients administered the RDD of 50 mg/day). Developmental effects consisted of fetal skeletal malformations of the ribs and vertebrae in rats. In rabbits, cleft lip was observed at 1 mg/kg/day and cleft lip and cleft palate were observed at 5 mg/kg/day (approximately 2.7 times the AUC in patients administered the RDD). Neither fetal loss nor malformations were observed in rats dosed at = 3 mg/kg/day (approximately 2.3 times the AUC in patients administered the RDD).

As angiogenesis is a critical component of embryonic and fetal development, inhibition of angiogenesis following administration of SUTENT[®] should be expected to result in adverse effects on pregnancy. There are no adequate and well-controlled studies of SUTENT[®] in pregnant women. If the drug is used during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant while receiving treatment with SUTENT[®].

PRECAUTIONS

Adverse events described in the following sections for MRCC patients are derived from Study 1 and Study 2. Adverse events discussed for GIST patients are derived from Study A, the randomized, placebo-controlled trial.

Left Ventricular Dysfunction

In the two MRCC studies, twenty-five patients (15%) had decreases in left ventricular ejection fraction (LVEF) to below the lower limit of normal (LLN). In GIST Study A, 22 patients (11%) on SUTENT[®] and 3 patients (3%) on placebo had treatment-emergent LVEF values below the LLN. Nine of twenty-two GIST patients on SUTENT[®] with LVEF changes recovered without intervention. Five patients had documented LVEF recovery following intervention (dose reduction- 1 patient; addition of antihypertensive or diuretic medications- 4 patients). Six patients went off study without documented recovery. Additionally, three patients (1%) on SUTENT[®] had Grade 3 reductions in left ventricular systolic function to LVEF < 40%; two of these patients died without receiving further study drug. No GIST patients on placebo had Grade 3 decreased LVEF. In GIST Study A, 1 patient (<1%) on SUTENT[®] and 1 patient (1%) on placebo died of diagnosed heart failure; 2 patients (1%) on SUTENT[®] and 2 patients (2%) on placebo died of treatment-emergent cardiac arrest.

Patients who presented with cardiac events within 12 months prior to SUTENT[®] administration,

such as myocardial infarction (including severe/unstable angina), coronary/peripheral artery bypass graft, symptomatic congestive heart failure (CHF), cerebrovascular accident or transient ischemic attack, or pulmonary embolism were excluded from SUTENT[®] clinical studies. It is unknown whether patients with these concomitant conditions may be at a higher risk of developing drug-related left ventricular dysfunction. Physicians are advised to weigh this risk against the potential benefits of the drug. **These patients should be carefully monitored for clinical signs and symptoms of CHF while receiving SUTENT[®]. Baseline and periodic evaluations of LVEF should also be considered while the patient is receiving SUTENT[®]. In patients without cardiac risk factors, a baseline evaluation of ejection fraction should be considered.**

In the presence of clinical manifestations of CHF, discontinuation of SUTENT[®] is recommended. The dose of SUTENT[®] should be interrupted and/or reduced in patients without clinical evidence of CHF but with an ejection fraction <50% and >20% below baseline.

Hemorrhagic Events

Bleeding events occurred in 44/169 patients (26%) receiving SUTENT[®] for MRCC and 37/202 patients (18%) receiving SUTENT[®] in GIST Study A, compared to 17/102 patients (17%) receiving placebo. Epistaxis was the most common hemorrhagic adverse event reported. Less common bleeding events in MRCC or GIST patients included rectal, gingival, upper GI, genital, and wound bleeding. Most events in MRCC patients were Grade 1 or 2; there was one Grade 3 event (bleeding foot wound). In GIST Study A, 14/202 patients (7%) receiving SUTENT[®] and 9/102 patients (9%) on placebo had Grade 3 or 4 bleeding events. In addition, one patient in Study A taking placebo had a fatal gastrointestinal bleeding event during cycle 2.

Tumor-related hemorrhage has been observed in patients treated with SUTENT[®]. These events may occur suddenly, and in the case of pulmonary tumors may present as severe and life threatening hemoptysis or pulmonary hemorrhage. Fatal pulmonary hemorrhage occurred in 2 patients receiving SUTENT[®] on a clinical trial of patients with metastatic non-small cell lung cancer (NSCLC). Both patients had squamous cell histology. SUTENT[®] is not approved for use in patients with NSCLC. Treatment-emergent Grade 3 and 4 tumor hemorrhage occurred in 5 of 202 patients (3%) with GIST receiving SUTENT[®] on Study A. Tumor hemorrhages were observed as early as cycle 1 and as late as cycle 6. One of these five patients received no further drug following tumor hemorrhage. None of the other four patients discontinued treatment or experienced dose delay due to tumor hemorrhage. No patients with GIST in the Study A placebo arm were observed to undergo intratumoral hemorrhage. Tumor hemorrhage has not been observed in patients with MRCC. Clinical assessment of these events should include serial complete blood counts (CBCs) and physical examinations.

Serious, sometimes fatal gastrointestinal complications including gastrointestinal perforation, have occurred rarely in patients with intra-abdominal malignancies treated with SUTENT[®].

QT Interval Prolongation

At approximately twice the therapeutic concentrations, sunitinib has been shown to prolong the QTcF (Fridericia's correction) interval (See section Special Population). There were no patients with greater than Grade 2 Common Terminology Criteria for Adverse Events v3.0 (CTCAE) QT/QTc interval prolongation. QT interval prolongation may lead to an increased risk for ventricular arrhythmias including torsade de pointes. Torsade de pointes has been observed in <0.1% of sunitinib-exposed patients. Sunitinib should be used with caution in patients with a known history of QT interval prolongation, patients who are taking antiarrhythmics, or patients with relevant pre-existing cardiac disease, bradycardia, or electrolyte disturbances. Concomitant treatment with strong CYP3A4 inhibitors, which may increase sunitinib plasma concentrations, should be used with caution and the dose of sunitinib reduced (See sections DOSAGE AND ADMINISTRATION and Drug-Drug Interaction).

Hypertension

Hypertension (all grades) was reported in 48/169 MRCC patients (28%), 31/202 GIST patients on SUTENT[®] (15%), and 11/102 GIST patients on placebo (11%). Grade 3 hypertension was reported in 10 MRCC patients (6%), 9 GIST patients on SUTENT[®] (4%), and none of the GIST patients on placebo. No Grade 4 hypertension was reported. SUTENT[®] dosing was reduced or temporarily delayed for hypertension in 6/169 MRCC patients (4%) and none of the patients in GIST Study A. No patients were discontinued from treatment with SUTENT[®] due to systemic hypertension. Severe hypertension (>200 mmHg systolic or 110 mmHg diastolic) occurred in 10/169 MRCC patients (6%), 8/202 GIST patients on SUTENT[®] (4%), and 1/102 GIST patients on placebo (1%).

Patients should be monitored for hypertension and treated as needed with standard antihypertensive therapy. In cases of severe hypertension, temporary suspension of SUTENT[®] is recommended until hypertension is controlled.

Adrenal Function

Adrenal toxicity was noted in non-clinical repeat dose studies of 14 days to 9 months in rats and monkeys at plasma exposures as low as 0.7 times the AUC observed in clinical studies. Histological changes of the adrenal gland were characterized as hemorrhage, necrosis, congestion, hypertrophy and inflammation. In clinical studies, CT/MRI obtained in 336 patients after exposure to one or more cycles of SUTENT[®] demonstrated no evidence of adrenal hemorrhage or necrosis. ACTH stimulation testing was performed in approximately 400 patients across multiple clinical trials of SUTENT[®]. Among patients with normal baseline ACTH stimulation testing, one patient developed consistently abnormal test results during treatment that are unexplained and may be related to treatment with SUTENT[®]. Eleven additional patients with normal baseline testing had abnormalities in the final test performed, with peak cortisol levels of 12-16.4 mcg/dL (normal >18 mcg/dL) following stimulation. None of these patients were reported to have clinical evidence of adrenal insufficiency.

Physicians prescribing SUTENT[®] are advised to monitor for adrenal insufficiency in patients who experience stress such as surgery, trauma or severe infection.

Thyroid Dysfunction

Baseline laboratory measurement of thyroid function is recommended and patients with hypothyroidism or hyperthyroidism should be treated as per standard medical practice prior to the start of sunitinib treatment. All patients should be observed closely for signs and symptoms of thyroid dysfunction on sunitinib treatment. Patients with signs and/or symptoms suggestive of thyroid dysfunction should have laboratory monitoring of thyroid function performed and be treated as per standard medical practice. Treatment-emergent acquired hypothyroidism was noted in 4% of GIST patients on sunitinib versus 1% on placebo. Hypothyroidism was reported as an adverse event in 2% of patients on sunitinib in the treatment-naïve MRCC study and one patient (<1%) in the IFN- α arm, and in 4% of patients across the two cytokine-refractory MRCC studies. Additionally, TSH elevations were reported in 2% of cytokine-refractory MRCC patients. Overall, 7% of the cytokine-refractory MRCC population had either clinical or laboratory evidence of treatment-emergent hypothyroidism.

Rare cases of hyperthyroidism, some followed by hypothyroidism, have been reported in clinical trials and through post-marketing experience.

Information for Patients

Gastrointestinal disorders such as diarrhea, nausea, stomatitis, dyspepsia, and vomiting were the most commonly reported gastrointestinal events occurring in patients who received SUTENT[®]. Supportive care for gastrointestinal adverse events requiring treatment may include anti-emetic or anti-diarrheal medication.

Serious, sometimes fatal gastrointestinal complications including gastrointestinal perforation, have occurred rarely in patients with intra-abdominal malignancies treated with sunitinib.

Increases in serum lipase and amylase were observed in patients with various solid tumors who received sunitinib. Increases in lipase levels were transient and were generally not accompanied by signs or symptoms of pancreatitis in subjects with various solid tumors. Pancreatitis was observed in 0.4% of patients with solid tumors. Hepatic failure was observed in <1% of solid tumor patients treated with sunitinib. If symptoms of pancreatitis or hepatic failure are present, patients should have sunitinib discontinued and be provided with appropriate supportive care.

Skin discoloration possibly due to the drug color (yellow) occurred in approximately 1/3 of patients. Patients should be advised that depigmentation of the hair or skin may occur during treatment with SUTENT[®]. Other possible dermatologic effects may include dryness, thickness or cracking of skin, blister or rash on the palms of the hands and soles of the feet. These events were not cumulative, were typically reversible and generally did not result in treatment discontinuation.

Other commonly reported adverse events included fatigue, high blood pressure, bleeding, swelling, mouth pain/irritation and taste disturbance.

Patients should be advised to inform their health care providers of all concomitant medications, including over-the-counter medications and dietary supplements (see Drug Interactions).

Laboratory Tests

CBCs with platelet count and serum chemistries including phosphate should be performed at the beginning of each treatment cycle for patients receiving treatment with SUTENT[®].

Drug Interactions

Co-administration of SUTENT[®] with strong inhibitors of the CYP3A4 family (e.g., ketoconazole, itraconazole, clarithromycin, atazanavir, indinavir, nefazodone, nelfinavir, ritonavir, saquinavir, telithromycin, voriconazole) may increase sunitinib concentrations. Grapefruit may also increase plasma concentrations of SUTENT[®] (see CLINICAL PHARMACOLOGY). Co-administration of SUTENT[®] with inducers of the CYP3A4 family (e.g., dexamethasone, phenytoin, carbamazepine, rifampin, rifabutin, rifapentin, phenobarbital, St. John's Wort) may decrease sunitinib concentrations (see CLINICAL PHARMACOLOGY). St. John's Wort may decrease SUTENT plasma concentrations unpredictably. Patients receiving SUTENT[®] should not take St. John's Wort concomitantly. SUTENT[®] dose modification is recommended in patients using concomitant CYP3A4 inhibitors or inducers (see DOSAGE AND ADMINISTRATION).

Carcinogenesis, Mutagenesis, Impairment of Fertility

Although definitive carcinogenicity studies with sunitinib have not been performed, carcinoma and hyperplasia of the Brunner's gland of the duodenum have been observed at the highest dose tested in H2ras transgenic mice administered doses of 0, 10, 25, 75, or 200 mg/kg/day for 28 days.

Sunitinib did not cause genetic damage when tested in *in vitro* assays (bacterial mutation [AMES Assay], human lymphocyte chromosome aberration) and an *in vivo* rat bone marrow micronucleus test.

Effects on the female reproductive system were identified in a 3-month repeat dose monkey study (2, 6, 12 mg/kg/day), where ovarian changes (decreased follicular development) were noted at 12 mg/kg/day (approximately 5.1 times the AUC in patients administered the RDD), while uterine changes (endometrial atrophy) were noted at ≥ 2 mg/kg/day (approximately 0.4 times the AUC in patients administered the RDD). With the addition of vaginal atrophy, the uterine and ovarian effects were reproduced at 6 mg/kg/day in the 9-month monkey study (0.3, 1.5 and 6 mg/kg/day administered daily for 28 days followed by a 14 day respite; the 6 mg/kg dose produced a mean AUC that was approximately 0.8 times the AUC in patients administered the RDD). A no effect level was not identified in the 3 month study; 1.5 mg/kg/day represents a no effect level in monkeys administered sunitinib for 9 months.

Although fertility was not affected in rats, SUTENT[®] may impair fertility in humans. In female rats, no fertility effects were observed at doses of ≤ 5.0 mg/kg/day [(0.5, 1.5, 5.0 mg/kg/day) administered for 21 days up to gestational day 7; the 5.0 mg/kg dose produced an AUC that was approximately 5 times the AUC in patients administered the RDD], however significant

embryolethality was observed at the 5.0 mg/kg dose. No reproductive effects were observed in male rats dosed (1, 3 or 10 mg/kg/day) for 58 days prior to mating with untreated females. Fertility, copulation, conception indices, and sperm evaluation (morphology, concentration, and motility) were unaffected by sunitinib at doses ≤ 10 mg/kg/day (the 10-mg/kg/day dose produced a mean AUC that was approximately 25.8 times the AUC in patients administered the RDD).

Pregnancy Category D: see WARNINGS.

Nursing Mothers

Sunitinib and/or its metabolites are excreted in rat milk. In lactating female rats administered 15 mg/kg, sunitinib and its metabolites were extensively excreted in milk at concentrations up to 12-fold higher than in plasma. It is not known whether SUTENT[®] or its primary active metabolite are excreted in human milk. Because drugs are commonly excreted in human milk and because of the potential for serious adverse reactions in nursing infants, women should be advised against breastfeeding while taking SUTENT[®].

Pediatric Use

The safety and efficacy of SUTENT[®] in pediatric patients have not been studied in clinical trials.

Physeal dysplasia was observed in Cynomolgus monkeys with open growth plates treated for ≥ 3 months (3 month dosing 2, 6, 12 mg/kg/day; 8 cycles of dosing 0.3, 1.5, 6.0 mg/kg/day) with sunitinib at doses that were > 0.4 times the RDD based on systemic exposure (AUC). In developing rats treated continuously for 3 months (1.5, 5.0 and 15.0 mg/kg) or 5 cycles (0.3, 1.5, and 6.0 mg/kg/day), bone abnormalities consisted of thickening of the epiphyseal cartilage of the femur and an increase of fracture of the tibia at doses ≥ 5 mg/kg (approximately 10 times the RDD based on AUC). Additionally, caries of the teeth were observed in rats at > 5 mg/kg. The incidence and severity of physeal dysplasia were dose-related and were reversible upon cessation of treatment however findings in the teeth were not. A no effect level was not observed in monkeys treated continuously for 3 months, but was 1.5 mg/kg/day when treated intermittently for 8 cycles. In rats the no effect level in bones was ≤ 2 mg/kg/day.

Geriatric Use

Of the 450 patients with solid tumors reported from clinical studies of SUTENT[®], 115 (25.6%) were 65 and over. No overall differences in safety or effectiveness were observed between younger and older patients.

EFFECT ON ABILITY TO DRIVE AND USE MACHINE

No studies on the effects on the ability to drive or operate machinery have been performed. Patients should be advised that they may experience dizziness during treatment with sunitinib.

ADVERSE REACTIONS

Overview

Four hundred fifty (450) patients with solid tumors including 257 patients (57%) with GIST and 169 patients (38%) with cytokine-refractory MRCC have been treated in 7 completed non-randomized, open-label, single arm clinical trials and 1 randomized, double-blind, placebo-controlled clinical trial. All patients received SUTENT[®] once daily as a 50-mg oral capsule on Schedule 4/2 (see CLINICAL STUDIES). One hundred two (102) patients received placebo in the randomized, double-blind, placebo-controlled clinical trial conducted in patients with GIST.

Adverse events occurring in GIST and MRCC studies are described below.

Adverse Events in GIST Study A

Median duration of blinded study treatment was two cycles for patients on SUTENT[®] (mean 3.0, range 1-9) and one cycle (mean 1.8, range 1-6) for patients on placebo. Dose reductions occurred in 23 patients (11%) on SUTENT[®] and none on placebo. Dose interruptions occurred in 59 patients (29%) on SUTENT[®] and 31 patients (30%) on placebo. The rates of treatment-emergent, non-fatal adverse events resulting in permanent discontinuation were 7% and 6% in the SUTENT[®] and placebo groups, respectively.

Most treatment-emergent adverse events in both study arms were Grade 1 or 2 in severity. Grade 3 or 4 treatment-emergent adverse events were reported in 56% vs. 51% of patients on SUTENT[®] versus placebo, respectively. Diarrhea, hypertension, bleeding, mucositis, skin abnormalities, and altered taste were more common in patients receiving SUTENT[®]. Table 4 compares the incidence of common (>10%) treatment-emergent adverse events for patients receiving SUTENT[®] versus those on placebo.

Table 4. Treatment-Emergent Adverse Events Reported in at Least 10% of GIST Patients Who Received SUTENT or Placebo in Study A*

	GIST			
	SUTENT (n=202)		Placebo (n=102)	
Adverse Event, n (%)	All Grades	Grade 3/4 ^a	All Grades	Grade 3/4 ^b
Any		114 (56)		52 (51)
Constitutional				
Fatigue	84 (42)	17 (8)	48 (47)	8 (8)
Fever	36 (18)	3 (2)	17 (17)	1 (1)

Gastrointestinal				
Diarrhea		9 (4)		0 (0)
Nausea	81 (40)	3 (2)	27 (27)	5 (5)
	63 (31)	2 (1)	33 (32)	2 (2)
Mucositis/stomatitis	58 (29)	4 (2)	18 (18)	3 (3)
Vomiting	49 (24)	0 (0)	24 (24)	2 (2)
Constipation	41 (20)	22	14 (14)	12
Abdominal pain ^c	67 (33)	(11)	39 (38)	(12)
Cardiac				
Hypertension	31 (15)	9 (4)	11 (11)	0 (0)
Dermatology				
Rash	28 (14)	2 (1)	9 (9)	0 (0)
Skin discoloration	61 (30)	0 (0)	23 (23)	0 (0)
Hand-foot syndrome	28 (14)	9 (4)	10 (10)	3 (3)
Neurology				
Altered taste	42 (21)	0 (0)	12 (12)	0 (0)
Headache	26 (13)	3 (2)	23 (23)	0 (0)
Musculoskeletal				
Arthralgia	24 (12)	2 (1)	16 (16)	0 (0)
Back pain	23 (11)	2 (1)	16 (16)	4 (4)
Myalgia/limb pain	28 (14)	1 (1)	9 (9)	1 (1)
Respiratory				
Dyspnea	20 (10)	0 (0)	19 (19)	3 (3)
Cough	17 (8)	0 (0)	13 (13)	0 (0)
Metabolism/Nutrition				
Anorexia ^d	67 (33)	1 (1)	30 (29)	5 (5)
Asthenia	45 (22)	10 (5)	11 (11)	3 (3)
Hemorrhage/bleeding				
Bleeding, all sites	37 (18)	14 (7)	17 (17)	9 (9)

* Common Toxicity Criteria for Adverse Events (CTCAE), Version 3.0

^a Grade 4 AEs in patient on SUTENT included abdominal pain (2%) and bleeding (2%).

^b Grade 4 AEs in patients on placebo included fatigue (3%), mucositis (1%), vomiting (1%), abdominal pain (3%), back pain (1%), and bone pain (1%).

^c Includes abdominal quadrant, gastric, hypochondrial, abdominal, flank, and cancer-related pain

^d Includes decreased appetite

Oral pain other than mucositis/stomatitis occurred in 12 patients (6%) on SUTENT[®] versus 3 (3%) on placebo. Hair color changes occurred in 15 patients (7%) on SUTENT[®] versus 4 (4%) on placebo. Alopecia was observed in 10 patients (5%) on SUTENT[®] versus 2 (2%) on placebo.

Table 5 provides common (≥10%) treatment-emergent laboratory abnormalities.

Table 5. Treatment-Emergent Laboratory Abnormalities (≥10%) from Study A*

Adverse Event, n (%)	SUTENT (n = 202)		Placebo (n = 102)	
	All Grades	Grade 3/4 ^a	All Grades	Grades 3/4 ^b
Any		68 (34)		22 (22)
Gastrointestinal				
AST/ALT	78 (39)	3 (2)	23 (23)	1 (1)
Alkaline phosphatase	48 (24)	7 (4)	21 (21)	4 (4)
Total bilirubin	32 (16)	2 (1)	8 (8)	0 (0)
Indirect Bilirubin	20 (10)	0 (0)	4 (4)	0 (0)
Amylase	35 (17)	10 (5)	12 (12)	3 (3)
Lipase	50 (25)	20 (10)	17 (17)	7 (7)
Cardiac				
Decreased LVEF	21 (10)	2 (1)	3 (3)	0 (0)
Renal / Metabolic				
Creatinine	25 (12)	1 (1)	7 (7)	0 (0)
Hypokalemia	24 (12)	1 (1)	4 (4)	0 (0)
Hypernatremia	20 (10)	0 (0)	4 (4)	1 (1)
Uric acid	31 (15)	16 (8)	16 (16)	8 (8)
Hematology				
Neutropenia	107 (53)	20 (10)	4 (4)	0 (0)
Lymphopenia	76 (38)	0 (0)	16 (16)	0 (0)
Anemia	52 (26)	6 (3)	22 (22)	2 (2)
Thrombocytopenia	76 (38)	10 (5)	4 (4)	0 (0)

*Common Toxicity Criteria for Adverse Events (CTCAE), Version 3.0

^a Grade 4 AEs in patients on SUTENT included alkaline phosphatase (1%), lipase (2%), creatinine (1%), hypokalemia (1%), neutropenia (2%), anemia (2%), and thrombocytopenia (1%).

^b Grade 4 AEs in patients on placebo included amylase (1%), lipase (1%), anemia (2%), and thrombocytopenia (1%).

Grade 3 or 4 treatment-emergent laboratory abnormalities were observed in 68 (34%) versus 22 (22%) patients on SUTENT[®] and placebo, respectively. Elevated liver function tests, pancreatic enzymes, and creatinine were more common in patients treated with SUTENT[®] than placebo. Decreased LVEF and myelosuppression were also more common with SUTENT[®] treatment. Treatment-emergent electrolyte disturbances of all types were more common in patients on SUTENT[®] than on placebo, including hyperkalemia (6% vs. 4%), hypokalemia (12% vs. 4%), hypernatremia (10% vs. 4%), hyponatremia (6% vs. 1%), and hypophosphatemia (9% vs. 0%). Three SUTENT[®] patients (1.5%) had Grade 3 hypophosphatemia. Acquired hypothyroidism was noted in 8 patients (4%) on SUTENT[®] versus 1 (1%) on placebo.

Adverse Events in the MRCC Studies

The data described below reflect exposure to SUTENT[®] in 169 patients with MRCC enrolled in Studies 1 and 2. The median duration of treatment was 5.5 months (range: 0.8-11.2) for Study 1

and 7.7 months (range: 0.2-16.1) for Study 2. Dose interruptions occurred in 48 patients (45%) on Study 1 and 45 patients (71%) on Study 2; one or more dose reductions occurred in 23 patients (22%) on Study 1 and 22 patients (35%) on Study 2. Table 6 summarizes treatment emergent adverse events for at least 10 % of all patients with MRCC who received at least one 50-mg dose of SUTENT®. Hematology laboratory abnormalities are presented separately, in Table 7.

Table 6. Treatment-Emergent Adverse Events Reported in at Least 10% of MRCC Patients Treated with SUTENT*

Adverse Event, n (%)	MRCC (N=169)	
	All Grades	Grade 3**
Any	169 (100)	123 (73)
Constitutional		
Fatigue	125 (74)	19 (11)
Fever	26 (15)	2 (1)
Gastrointestinal		
Diarrhea	93 (55)	8 (5)
Nausea	92 (54)	4 (2)
Mucositis/stomatitis	90 (53)	7 (4)
Dyspepsia	77 (46)	1 (1)
Vomiting	63 (37)	7 (4)
Constipation	57 (34)	1 (1)
Abdominal pain	34 (20)	5 (3)
Glossodynia	25 (15)	0 (0)
Flatulence	24 (14)	0 (0)
Cardiac		
Hypertension	48 (28)	10 (6)
Edema, peripheral	28 (17)	1 (1)
Dermatology		
Rash	64 (38)	1 (1)
Skin discoloration	55 (33)	0 (0)
Dry skin	29 (17)	0 (0)
Hair color changes	29 (17)	0 (0)
Hand-foot syndrome	21 (12)	5 (3)
Alopecia	20 (12)	0 (0)
Neurology		
Altered taste	73 (43)	0 (0)
Headache	43 (25)	2 (1)
Dizziness	27 (16)	3 (2)

Musculoskeletal		
Arthralgia	48 (28)	2 (1)
Pain in limb	31 (18)	1 (1)
Back pain	29 (17)	1 (1)
Myalgia	29 (17)	1 (1)
Respiratory		
Dyspnea	47 (28)	8 (5)
Cough	29 (17)	1 (1)
Metabolism/Nutrition		
Anorexia	53 (31)	1 (1)
Dehydration	19 (11)	5 (3)
Hemorrhage/bleeding		
Bleeding, all sites	44 (26)	1 (1)

* Common Toxicity Criteria for Adverse Events (CTCAE), Version 3.0

** There were no Grade 4 adverse events among the events reported with a $\geq 10\%$ incidence in the MRCC population.

Other significant adverse events occurring in MRCC patients receiving SUTENT[®] included peripheral neuropathy (10%), appetite disturbance (9%), blistering of the skin (7%), periorbital edema (7%) and increased lacrimation (6%).

Table 7. Treatment-Emergent Grade 3 and 4 Hematology Laboratory Abnormalities* from Studies 1 and 2

Laboratory Test	Unit	MRCC (N=169)		
		Grade 3	Grade 4	Total (Grade 3 + 4)
Hematology, n (%)	10 ⁹ /L	54 (32)		
Neutropenia	g/L	21 (12)	4 (2)	58 (34)
Anemia	10 ⁹ /L	9 (5)	1 (1)	22 (13)
Lymphopenia	10 ⁹ /L	33 (20)	3 (2)	12 (7)
Thrombocytopenia	10 ⁹ /L	33 (20)	2 (1)	35 (21)
Leukopenia	10 ⁹ /L	5 (3)	0 (0)	5 (3)
	10 ⁹ /L	12 (7)	0 (0)	12 (7)

* Common Toxicity Criteria for Adverse Events (CTCAE), Version 3.0

Common treatment-emergent Grade 3 and 4 chemistry laboratory abnormalities in the MRCC studies included increased lipase (16%), increased amylase (5%), hypophosphatemia (10%), and hyperuricemia (10%).

Cardiovascular

See PRECAUTIONS section for information on left ventricular dysfunction.

Two patients with MRCC experienced Grade 3 myocardial ischemia, one had Grade 2 “cardiovascular toxicity” reported as an adverse event and one patient experienced a fatal myocardial infarction while on treatment.

Data from non-clinical (*in vitro* and *in vivo*) studies indicate that sunitinib has the potential to inhibit the cardiac action potential repolarization process (e.g., prolongation of QT interval). In GIST Study A, 23 patients (11%) on SUTENT[®] versus 12 (12%) on placebo had observed QT prolongation greater than 20 milliseconds from baseline. No consistent, clinically significant QTc prolongation has been observed in completed clinical studies.

Venous Thromboembolic Events

Four patients (2%) on the two MRCC studies had venous thromboembolic events reported; two patients with pulmonary embolism (both Grade 4) and two patients with deep venous thrombosis (DVT) (both Grade 3). Dose interruption occurred in one of these cases. Seven patients (3%) on SUTENT[®] and none on placebo in GIST Study A experienced venous thromboembolic events; five of the seven were Grade 3 DVTs, and two were Grade 1 or 2. Four of these seven GIST patients discontinued treatment following first observation of DVT.

Seizures

In clinical studies of SUTENT[®], seizures have been observed in subjects with radiological evidence of brain metastases. In addition, there have been rare (<1%) reports of subjects presenting with seizures and radiological evidence of reversible posterior leukoencephalopathy syndrome (RPLS). None of these subjects had a fatal outcome to the event. Patients with seizures and signs/symptoms consistent with RPLS, such as hypertension, headache, decreased alertness, altered mental functioning, and visual loss, including cortical blindness should be controlled with medical management including control of hypertension. Temporary suspension of SUTENT[®] is recommended; following resolution, treatment may be resumed at the discretion of the treating physician.

Post-marketing Experience

The following adverse reactions have been identified during post-approval use of sunitinib. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Blood and lymphatic system disorders:

Rare cases of thrombotic microangiopathy have been reported. Temporary suspension of sunitinib is recommended; following resolution, treatment may be resumed at the discretion of the treating physician.

Endocrine disorders:

Rare cases of hyperthyroidism, some followed by hypothyroidism, have been reported in clinical trials and through post-marketing experience (See also section PRECAUTION).

Immune system disorders:

Hypersensitivity reactions, including angioedema, have been reported.

Infections and infestations:

Cases of serious infection (with or without neutropenia), in some cases with fatal outcome, have been reported.

Musculoskeletal and connective tissue disorders:

Rare cases of myopathy and/or rhabdomyolysis with or without acute renal failure, have been reported. Most of these patients had pre-existing risk factors and/or were receiving concomitant medications known to be associated with these adverse reactions. Patients with signs or symptoms of muscle toxicity should be managed as per standard medical practice.

Nervous system disorders:

Taste disturbance, including ageusia, has been reported.

Renal and urinary disorders:

Cases of proteinuria and rare cases of nephrotic syndrome have been reported. Baseline urinalysis is recommended, and patients should be monitored for the development or worsening of proteinuria. The safety of continued sunitinib treatment in patients with moderate to severe proteinuria has not been systematically evaluated. Discontinue sunitinib in patients with nephrotic syndrome.

Other:

Fistula formation: Cases of fistula formation, sometimes associated with tumor necrosis and/or regression, in some cases with fatal outcome, have been reported.

Laboratory Abnormalities/Testing***Hematologic Events***

Grade 3 and 4 neutropenia were reported in 21 (13%) and 1 (1%) patients with MRCC, 19 (9%) and 3 (2%) patients with GIST on SUTENT[®], respectively. In Study A, one patient each in the SUTENT[®] and placebo groups had febrile neutropenia. Grade 3 and 4 thrombocytopenia was reported in 5 (3%) and 0 patients with MRCC, 7 (4%) and 1 (1%) patients with GIST on SUTENT[®], respectively. No GIST patients receiving placebo experienced either Grade 3 or 4 neutropenia or thrombocytopenia. The rates of dose reductions and delays for hematologic abnormalities were 4% and 2% for neutropenia, 2% and 0% for anemia, and 1% and 1% for thrombocytopenia for MRCC and GIST patients, respectively. One MRCC patient with an adverse event report of Grade 4 thrombocytopenia discontinued treatment.

Patients receiving SUTENT[®] should be monitored regularly for myelosuppression.

Hypothyroidism

Hypothyroidism was reported as an adverse event in 7 patients (4%) across the two MRCC studies. Additionally, TSH elevations were reported in 4 patients (2%). Overall, 7% of the MRCC population had either clinical or laboratory evidence of treatment-emergent hypothyroidism. Treatment-emergent acquired hypothyroidism was noted in 8 GIST patients (4%) on SUTENT[®]

versus 1 (1%) on placebo.

Patients with symptoms suggestive of hypothyroidism should have laboratory monitoring of thyroid function performed and be treated as per standard medical practice.

Pancreatic Function

Grade 3 and 4 increases in serum lipase were observed in 23 (14%) and 4 (2%), respectively, of 169 patients receiving SUTENT[®] for MRCC. Grade 3 and 4 increases in serum amylase were observed in 8 (5%) and 1 (1%) MRCC patients, respectively. Increases in lipase levels were transient and were generally not accompanied by signs or symptoms of pancreatitis in subjects with either MRCC or GIST. Pancreatitis has been observed rarely (<1%) in patients receiving SUTENT[®] for GIST or MRCC. If symptoms of pancreatitis are present, patients should have SUTENT[®] discontinued and be provided with appropriate supportive care.

OVERDOSAGE

No overdose of SUTENT[®] was reported in completed clinical studies. In non-clinical studies mortality was observed following as few as 5 daily doses of 500 mg/kg (3000 mg/m²) in rats. At this dose, signs of toxicity included impaired muscle coordination, head shakes, hypoactivity, ocular discharge, piloerection and gastrointestinal distress. Mortality and similar signs of toxicity were observed at lower doses when administered for longer durations. Treatment of overdose with SUTENT[®] should consist of general supportive measures. There is no specific antidote for overdosage with SUTENT[®]. If indicated, elimination of unabsorbed drug should be achieved by emesis or gastric lavage.

DOSAGE AND ADMINISTRATION

The recommended dose of SUTENT[®] for GIST and advanced RCC is one 50-mg oral dose taken once daily, on a schedule of 4 weeks on treatment followed by 2 weeks off (Schedule 4/2) to comprise a complete cycle of 6 weeks. SUTENT[®] may be taken with or without food.

Dose Modification

Dose increase or reduction of 12.5-mg increments is recommended based on individual safety and tolerability.

Strong CYP3A4 inhibitors such as ketoconazole may increase SUTENT[®] plasma concentrations. Selection of an alternate concomitant medication with no or minimal enzyme inhibition potential is recommended. A dose reduction for SUTENT[®] to a minimum of 37.5 mg daily should be considered if SUTENT[®] must be co-administered with a strong CYP3A4 inhibitor (see CLINICAL PHARMACOLOGY and PRECAUTIONS, Drug Interactions).

CYP3A4 inducers such as rifampin may **decrease** SUTENT[®] plasma concentrations. Selection of an alternate concomitant medication with no or minimal enzyme induction potential is recommended. A dose increase for SUTENT[®] to a maximum of 87.5 mg daily should be

considered if SUTENT[®] must be co-administered with a CYP3A4 inducer. If dose is increased, the patient should be monitored carefully for toxicity (see CLINICAL PHARMACOLOGY and PRECAUTIONS, Drug Interactions). St. John's Wort may decrease SUTENT[®] plasma concentrations unpredictably. Patients receiving SUTENT[®] should not take St. John's Wort concomitantly.

Selection of an alternate concomitant medication with no, or minimal potential to induce or inhibit CYP3A4 is recommended.

Use in pediatrics: The safety and efficacy of sunitinib in pediatric patients have not been established.

Use in the elderly: Dose adjustments are not required in elderly patients. Approximately 34% of the subjects in clinical studies of sunitinib were 65 years of age or over. No significant differences in safety or effectiveness were observed between younger and older patients.

Hepatic Insufficiency: No dose adjustment is necessary when administering sunitinib to patients with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment. Sunitinib was not studied in subjects with severe (Child-Pugh Class C) hepatic impairment (See section Pharmacokinetics Properties).

Renal Insufficiency: No clinical studies were conducted in patients with impaired renal function. Studies that were conducted excluded patients with serum creatinine > 2.0 x ULN; within this range no dosing adjustments are recommended (See section Pharmacokinetics Properties).

Storage Condition

Store below 30° C. Keep in dry place.

Supply

SUTENT[®] 12.5 mg capsules; HDPE bottle @ 30 capsules; Reg.No :DKI0754200701A1

SUTENT[®] 25 mg capsules; HDPE bottle @ 30 capsules; Reg No. :DKI0754200701B1

SUTENT[®] 50 mg capsules; HDPE bottle @ 30 capsules; Reg. No :DKI0754200701C1

HARUS DENGAN RESEP DOKTER

Manufactured by:

Pfizer Italia S.r.l., Italy

Imported by:

PT Pfizer Indonesia

PO Box 2706, Jakarta

Indonesia